

Pacific Journal of Mathematics

QUOTIENT-UNIVERSAL SEQUENTIAL SPACES

R. SIROIS-DUMAIS AND STEPHEN WILLARD

QUOTIENT-UNIVERSAL SEQUENTIAL SPACES

R. SIROIS-DUMAIS AND S. WILLARD

We produce 2^c mutually nonhomeomorphic countable sequential spaces. These are used

(1) to answer in the negative the following question of Michael and Stone [4]: is every regular T_1 -space which is a quotient of some separable metric space and a continuous image of the space \mathbf{P} of irrationals a quotient of \mathbf{P} ?

(2) to characterize c (with or without the continuum hypothesis) as the smallest cardinal κ with the property that a metric space of cardinality κ exists of which every sequential space of cardinality $\leq \kappa$ is a quotient.

1. Introduction. We let Q denote the space of rationals, \mathbf{P} the space of irrationals, \mathbf{R} the real line, and c the cardinality of \mathbf{R} . For any set X , the cardinality of X is denoted $|X|$.

We begin with the basic construction, which will be applied in the sequel in two different directions. Denote by Y the set $[Q \times (Q - \{0\})] \cup \{\infty\}$ and, for $E \subseteq \mathbf{R}$, denote by τ_E the quotient topology induced on Y by the obvious map from the subspace $[Q \times (Q - \{0\})] \cup (E \times \{0\})$ of $\mathbf{R} \times \mathbf{R}$. The set Y endowed with the topology τ_E will be denoted Y_E . Note that Y_E is a countable, regular, T_1 -space which is, by construction, the quotient of a separable metric space. (Thus, see [3], Y_E is both an \aleph_0 -space and a k -space.)

2. Quotients of \mathbf{P} . In [4], Michael and Stone establish that every metrizable continuous image of \mathbf{P} is a quotient of \mathbf{P} . The question is raised there whether this result can be extended to nonmetrizable images of \mathbf{P} , that is, whether a regular T_1 -space which is at the same time a quotient of some separable metric space and a continuous image of \mathbf{P} must be a quotient of \mathbf{P} . The construction of §1 provides the negative answer. To see this, first note that the countable discrete space (hence, every countable space) is a continuous image of \mathbf{P} (collapse each interval $(n, n + 1)$ to a point). It follows that each space Y_E is a regular T_1 -space which is a continuous image of \mathbf{P} and a quotient of some separable metric space. But:

THEOREM. *Not every space Y_E is a quotient of \mathbf{P} .*

Proof. If E and F are distinct subsets of \mathbf{R} , the topologies τ_E and τ_F on Y are different, one containing a set containing ∞ which does not belong to the other.

Now let S be the set of all surjections $f: \mathbf{P} \rightarrow Y$ such that each $f^{-1}(y)$, $y \in Y$, is closed in \mathbf{P} , and let Φ be the set of all $\phi: Y \rightarrow 2^{\mathbf{P}}$, where $2^{\mathbf{P}}$ denotes the collection of closed subsets of \mathbf{P} . Then $f \rightarrow f^{-1}$ is a one-one map from S into Φ ; since $|\Phi| = \mathfrak{c}^{\mathfrak{c}_0} = \mathfrak{c}$, we have $|S| \leq \mathfrak{c}$. Let J be the set of all T_1 topologies τ on Y such that (Y, τ) is a quotient image of \mathbf{P} . Then each $\tau \in J$ is generated by some $f \in S$, so $|J| \leq \mathfrak{c}$. Since $|\{\tau_E \mid E \subseteq \mathbf{R}\}| = 2^{\mathfrak{c}}$, and since each τ_E is T_1 , it follows that (Y, τ_E) is not a quotient of \mathbf{P} for some $E \subseteq \mathbf{R}$.

NOTES. (1) From the above, it is easily seen that there are $2^{\mathfrak{c}}$ nonhomeomorphic spaces Y_E , at most \mathfrak{c} of which can be quotients of \mathbf{P} . This result can be sharpened, with some difficulty. In fact, Y_E is a quotient of \mathbf{P} iff E is an analytic subset of \mathbf{R} .

(2) If, in the construction of Y , the set $Q \times (Q - \{0\})$ is replaced by a discrete space, say $\{(k/n, 1/n) \mid k, n \in \mathbf{N}\}$, the spaces Y_E which result still work, and have now the additional property that each has only one nonisolated point.

3. Quotient-universal sequential spaces. Let κ be an infinite cardinal and let $S(\kappa)$ denote the collection of all sequential spaces of cardinality $\leq \kappa$. A sequential space S is *quotient-universal** for $S(\kappa)$ if $S \in S(\kappa)$ and every $T \in S(\kappa)$ is a quotient of S . We are particularly interested in the existence of metrizable quotient-universal spaces for $S(\kappa)$.

Whenever $\kappa^{\mathfrak{c}_0} = \kappa$, the disjoint union of κ copies of the converging sequence will serve as a metrizable quotient-universal space for $S(\kappa)$. In particular, there is a metrizable quotient-universal space for $S(\mathfrak{c})$. In this section, we use the construction of §1 to demonstrate that, whether or not the continuum hypothesis is true, \mathfrak{c} is the smallest cardinal for which this is true. In fact, we exhibit a countable sequential space which is not a quotient of any metric space of cardinality $< \mathfrak{c}$.

LEMMA. *There exists a subset E of \mathbf{R} with $|E| = \mathfrak{c}$ which contains no uncountable closed subset of \mathbf{R} .*

Proof. Let $\{C_\alpha \mid \alpha < \mathfrak{c}\}$ be a transfinite enumeration of the \mathfrak{c} uncountable closed subsets of \mathbf{R} . Pick p_0 and q_0 in C_0 with $p_0 \neq q_0$. If p_α and q_α have been chosen in C_α for $\alpha < \beta$ so that all p_α and q_α are distinct, choose p_β and q_β in C_β so that $p_\beta \neq q_\beta$ and p_β, q_β are distinct from all p_α, q_α for $\alpha < \beta$. This is possible since any uncountable closed subset of \mathbf{R} has cardinal \mathfrak{c} so that $C_\beta - \{p_\alpha, q_\alpha \mid \alpha < \beta\} \neq \emptyset$.

* The term "universal" has been preempted by those who study spaces with a given property P which contain as subspaces every space (of appropriate cardinality or weight) having property P . See, for example, [2], [5] and [6].

Let $E = \{p_\alpha \mid \alpha < \mathfrak{c}\}$. Then $|E| = \mathfrak{c}$ and E contains no uncountable closed subset of \mathbf{R} since $q_\alpha \in C_\alpha - E$ for each α .

Let $E \subseteq \mathbf{R}$ be the set of the lemma. Let M_E denote the subspace $[Q \times (Q - \{0\})] \cup (E \times \{0\})$ of $\mathbf{R} \times \mathbf{R}$. Recall that Y_E is the quotient of M_E obtained by collapsing $E \times \{0\}$ to a single point e . Let $q: M_E \rightarrow Y_E$ be the quotient map.

Y_E is a countable sequential space, but:

THEOREM. Y_E is not the quotient of any metric space of cardinality $< \mathfrak{c}$.

Proof. Suppose there is a quotient map f of S onto Y_E , where S is a metric space and $|S| = \kappa < \mathfrak{c}$. For each $p \in E$, let $\sigma_p = (x_{p1}, x_{p2}, \dots)$ be a sequence in $Q \times (Q - \{0\})$ such that

$$|x_{pn} - (p, 0)| \leq \min \left\{ \frac{1}{n}, |x_{p(n-1)} - (p, 0)| \right\}.$$

Recall that q denotes the quotient map of M_E onto Y_E . For each n , let

$$z_{pn} = q(x_{pn})$$

and denote by η_p the sequence (z_{p1}, z_{p2}, \dots) in Y_E . Now $\eta_p \rightarrow e$. Hence, since f is a hereditary quotient map, there exists some $b_p \in f^{-1}(e)$ and a sequence $\sigma_p = (s_{p1}, s_{p2}, \dots)$ in $S - f^{-1}(e)$ such that $\sigma_p \rightarrow b_p$ and $f(\sigma_p) = \eta_p$. Let

$$f^{-1}(e) = \{x_\alpha \mid \alpha < \kappa\}$$

and, for $\alpha < \kappa$, let

$$A_\alpha = \{p \in E \mid b_p = x_\alpha\}.$$

We claim some A_α must contain a sequence (p_i) converging to some element of $\mathbf{R} - E$. For otherwise $C1_{\mathbf{R}}(A_\alpha) \subset E$ for each $\alpha < \kappa$, whence E is the union of fewer than \mathfrak{c} closed sets. But since $|E| = \mathfrak{c}$, one of these would be an uncountable closed set in E , contradicting the construction of E .

Without loss of generality, say A_1 contains a sequence (p_i) which is closed and discrete in E . Then the sequence $\eta_{p_i} = (z_{p_i1}, z_{p_i2}, \dots)$ converges to e , for each i , and the sequence $\delta_{p_i} = (s_{p_i1}, s_{p_i2}, \dots)$ converges to x_1 , for each i . A diagonal sequence $(s_{p_{i1}n_i}, s_{p_{i2}n_i}, \dots)$ with $n_k \geq k$ for each k will then converge to x_1 . Then $(z_{p_{i1}n_i}, z_{p_{i2}n_i}, \dots)$ converges to e . Hence $(x_{p_{i1}n_i}, x_{p_{i2}n_i}, \dots)$ must have a cluster point in M_E .

But $|x_{p_k n_k} - (p_k, 0)| \leq |x_{p_k k} - (p_k, 0)| \leq 1/k$, so any cluster point of $(x_{p_1 n_1}, x_{p_2 n_2}, \dots)$ in M_E would be a cluster point of $((p_1, 0), (p_2, 0), \dots)$, which is impossible by choice of the p_i .

We conclude with some observations on extension of the result above.

(1) As noted in §2, there are 2^c mutually nonhomeomorphic spaces Y_E . Since there are at most c quotients of any single countable sequential space, there can exist no quotient-universal space (metrizable or not) for $S(\aleph_0)$. It is at least consistent with the usual (Zermelo-Fraenkel) axioms for set theory (with Choice) that this result extends to all cardinals $\kappa < c$, for Martin's axiom entails $2^\kappa < 2^c$ for $\kappa < c$.

(2) Let $M(\kappa)$ denote the collection of metrizable spaces of cardinal $\leq \kappa$. The space Q of rationals is a (metrizable) quotient-universal space for $M(\aleph_0)$, while the disjoint union of c copies of the converging sequence is a quotient-universal space for $M(c)$. For cardinals κ between \aleph_0 and c little is known. Baumgartner ([1]) has shown that it is consistent with Zermelo-Fraenkel set theory with choice that all \aleph_1 -dense subsets of \mathbf{R} are order-isomorphic. (A subset A of \mathbf{R} is \aleph_1 -dense if whenever $a < b$ in \mathbf{R} , $(a, b) \cap A$ has cardinal \aleph_1 .) If this is the case, then every separable metric space M of cardinal $\leq \aleph_1$ is a quotient of the unique \aleph_1 -dense subset D of \mathbf{R} . For M is a quotient of $M \times D$, while ([7], Theorem 76) $M \times D$ is homeomorphic to a subset of \mathbf{R} and hence, by Baumgartner's result, to D .

REFERENCES

1. James E. Baumgartner, *All \aleph_1 -dense sets of reals can be isomorphic*, Fund. Math., **79** (1973), 103–106.
2. Jonsson, *Universal Relational Systems*, Math. Scand., **4** (1956), 193–208.
3. E. Michael, *\aleph_1 -spaces*, J. Math. Mech., **15** (1966), 983–1002.
4. E. Michael and A. H. Stone, *Quotients of the space of irrationals*, Pacific J. Math., **28** (1969), 629–633.
5. R. Pol, *There is no universal totally disconnected space*, Fund. Math., **79** (1973), 265–267.
6. W. Sierpinski, *Sur les espaces métriques universels*, Fund. Math., **33** (1945), 123–136.
7. ———, *General Topology*, 2nd ed., U. of Toronto Press, 1956 (translated by C. C. Krieger).

Received March 12, 1976 and in revised form May 12, 1976.

UNIVERSITY OF ALBERTA

Helen Elizabeth Adams, <i>Factorization-prime ideals in integral domains</i>	1
Patrick Robert Ahern and Robert Bruce Schneider, <i>The boundary behavior of Henkin's kernel</i>	9
Daniel D. Anderson, Jacob R. Matijevic and Warren Douglas Nichols, <i>The Krull intersection theorem. II</i>	15
Efraim Pacillas Armendariz, <i>On semiprime P.I.-algebras over commutative regular rings</i>	23
Robert H. Bird and Charles John Parry, <i>Integral bases for bicyclic biquadratic fields over quadratic subfields</i>	29
Tae Ho Choe and Young Hee Hong, <i>Extensions of completely regular ordered spaces</i>	37
John Dauns, <i>Generalized monofrom and quasi injective modules</i>	49
F. S. De Blasi, <i>On the differentiability of multifunctions</i>	67
Paul M. Eakin, Jr. and Avinash Madhav Sathaye, <i>R-endomorphisms of $R[[X]]$ are essentially continuous</i>	83
Larry Quin Eifler, <i>Open mapping theorems for probability measures on metric spaces</i>	89
Garret J. Etgen and James Pawlowski, <i>Oscillation criteria for second order self adjoint differential systems</i>	99
Ronald Fintushel, <i>Local S^1 actions on 3-manifolds</i>	111
Kenneth R. Goodearl, <i>Choquet simplexes and σ-convex faces</i>	119
John R. Graef, <i>Some nonoscillation criteria for higher order nonlinear differential equations</i>	125
Charles Henry Heiberg, <i>Norms of powers of absolutely convergent Fourier series: an example</i>	131
Les Andrew Karlovitz, <i>Existence of fixed points of nonexpansive mappings in a space without normal structure</i>	153
Gangaram S. Ladde, <i>Systems of functional differential inequalities and functional differential systems</i>	161
Joseph Michael Lambert, <i>Conditions for simultaneous approximation and interpolation with norm preservation in $C[a, b]$</i>	173
Ernest Paul Lane, <i>Insertion of a continuous function</i>	181
Robert F. Lax, <i>Weierstrass points of products of Riemann surfaces</i>	191
Dan McCord, <i>An estimate of the Nielsen number and an example concerning the Lefschetz fixed point theorem</i>	195
Paul Milnes and John Sydney Pym, <i>Counterexample in the theory of continuous functions on topological groups</i>	205
Peter Johanna I. M. De Paepe, <i>Homomorphism spaces of algebras of holomorphic functions</i>	211
Judith Ann Palagallo, <i>A representation of additive functionals on L^p-spaces, $0 < p < 1$</i>	221
S. M. Patel, <i>On generalized numerical ranges</i>	235
Thomas Thornton Read, <i>A limit-point criterion for expressions with oscillatory coefficients</i>	243
Elemer E. Rosinger, <i>Division of distributions</i>	257
Peter S. Shoenfeld, <i>Highly proximal and generalized almost finite extensions of minimal sets</i>	265
R. Sirois-Dumais and Stephen Willard, <i>Quotient-universal sequential spaces</i>	281
Robert Charles Thompson, <i>Convex and concave functions of singular values of matrix sums</i>	285
Edward D. Tymchatyn, <i>Some n-arc theorems</i>	291
Jang-Mei Gloria Wu, <i>Variation of Green's potential</i>	295