Pacific Journal of Mathematics

NTH ORDER OSCILLATIONS WITH MIDDLE TERMS OF ORDER N - 2

ATHANASSIOS G. KARTSATOS

Vol. 67, No. 2

February 1976

NTH ORDER OSCILLATIONS WITH MIDDLE TERMS OF ORDER N-2

ATHANASSIOS G. KARTSATOS

Equations of the form

(*)
$$x^{(n)} + p(t)x^{(n-2)} + H(t, x) = 0$$
, $n \ge 3$

are studied here, where $p(t) \ge 0$ and $uH(t, u) \ge 0$.

It is shown, among other things, that for n even all solutions of (*) oscillate if H is superlinear, satisfying the usual integral conditions, and u'' + p(t)u = 0 is nonoscillatory with p(t) "small" and decreasing. The case of n odd is also covered and some of the recent results of Waltman, Heidel are taken as special cases.

As far as the author knows, the only references concerning the nonlinear (*) for $p(t) \neq 0$ are those of Liossatos [9] and the author [8]. In [9] Liossatos extended several of the results of Heidel [2] to the *n*th order case. Liossatos however, considered mainly the case $p(t) \leq 0$, $uH(t, u) \leq 0$ (actually a special case of H(t, u)) and established one theorem concerning the case $p(t) \geq 0$, $uH(t, u) \geq 0$. One of the results of Liossatos was improved by the author in [8], where a result is also given for a forced equation.

In Theorem 1 below we establish a result for n = 3 which improves a result of Waltman [13, Th. 1]. The proof of this theorem contains a simpler proof of Theorem 3.8 in Heidel's paper [2]. In Theorem 2 we provide conditions ensuring the oscillation or convergence to zero of all bounded solutions of (*). Theorem 3 ensures that every solution of (*) with a zero is oscillatory if n is odd. In Theorem 4 the middle term is treated as a small perturbation, and Theorem 5 ensures oscillation or convergence to zero for functions H which are "large" compared to the middle term. For several results concerning nth order nonlinear equations the reader is referred to the survey paper [8].

2. Preliminaries. In what follows, T will denote a fixed nonnegative number, $R_T = [T, \infty)$, $R_+ = [0, \infty)$, $R = (-\infty, \infty)$. For the equation

$$(*)$$
 $x^{(n)} + p(t)x^{(n-2)} + H(t, x) = 0$, $n \ge 3$

we shall assume that the functions $p: R_T \to R_+$, $H: R_T \times R \to R$ are continuous on their domains and such that $uH(t, u) \ge 0$ for every $(t, u) \in R_T \times R$. A "solution" of (*) will be any function x(t) which

is defined, n times continuously differentiable, and satisfies (*) on some infinite subinterval of R_T . Only nontrivial solutions will be considered here, i.e., solutions which do not become identically equal to zero for all large t. A function $f:[a, \infty) \to R$, $a \ge T$, is said to be "oscillatory" if it has an unbounded set of zeros in $[a, \infty)$. The equation (*) is said to be "D-oscillatory", if, for n even, every solution of (*) is oscillatory, and, for n odd, every solution of (*) is oscillatory, or tends monotonically to zero as $t \to \infty$. The equation (*) is said to be "BD-oscillatory" if the above definition holds true, but for the bounded solutions of (*). A stronger result than the following auxiliary theorem was proven by the author in [6] but for n even.

THEOREM A. Consider the equation

(2.1)
$$x^{(n)} + H(t, x) = 0$$
, $t \ge T$

and the inequality

$$(2.2) u^{(n)} + H(t, u) \leq 0, \quad t \geq T,$$

where H(t, u) is nonnegative and increasing in u for every $u \in R$. Then the existence of a positive (bounded and positive) solution of (2.2) implies the existence of a positive (bounded and positive) solution of (2.1). Furthermore, if all positive (bounded and positive) solutions of (2.1) tend monotonically to zero, then the same is true for all positive (bounded and positive) solutions of (2.2).

Proof. For *n* even the theorem is a particular case of Lemma 2.1 in [6], where H(t, u) > 0 was considered but its proof carries over to the present case without modifications. For *n* odd, assume that all positive solutions of (2.1) tend monotonically to zero, and let u(t) be a positive solution of (2.2) such that u(t) > 0, u'(t) > 0 for $t \ge t_1 \ge T$. Then, from the proof of Lemma 2.1 in [6], it follows immediately, that (2.1) has also a solution x(t) with $x(t_1) = u(t_1)$ and x'(t) > 0, $t \ge t_1$. This is however a contradiction to the assumption $\lim_{t\to\infty} x(t) = 0$.

For a functional version of the above theorem, the reader is referred to Onose [11].

THEOREM B. In the equation (*) assume that x(t) is a nonoscillatory solution and that u'' + p(t)u is nonoscillatory. Then $x(t)x^{(n-2)}(t) \ge 0$ or $x(t)x^{(n-2)}(t) \le 0$ for all large t. If, moreover, uH(t, u) > 0 for every $(t, u) \in R_T \times R$ with $u \ne 0$, then these inequalities are strict. REMARK. The second assertion of this theorem is slightly stronger than the corresponding result of Heidel [2, Th. 3.6].

Proof. We partly follow the steps of Heidel's Theorem 3.6 in [2]. Let x(t) be a nonoscillatory solution of (*) and assume that $x(t) > 0, t \ge t_1 \ge T$. A similar proof covers the case x(t) < 0. Now let $x^{(n-2)}(t)$ be oscillatory, let $x^{(n-2)}(t_2) = 0$ for some $t_2 > t_1$, and assume that $x^{(n-2)}(t_3) < 0$ for some $t_3 > t_2$. Now let

$$egin{aligned} L &= \sup \left\{ t \in [t_2,\,t_3);\, x^{(n-2)}(t) = 0
ight\} \,, \ M &= \inf \left\{ t \in (t_3,\,\infty);\, x^{(n-2)}(t) = 0
ight\} \,. \end{aligned}$$

Then, by continuity, $x^{(n-2)}(L) = 0$, $x^{(n-2)}(M) = 0$, $L < t_3 < M$ and $x^{(n-2)}(t) < 0$, $t \in (L, M)$. Now multiply (*) by $x^{(n-2)}(t)$ and integrate between L, M to obtain

(2.3)
$$\int_{L}^{M} p(s) [x^{(n-2)}(s)]^{2} ds + \int_{L}^{M} H(s, x(s)) x^{(n-2)}(s) ds$$
$$= \int_{L}^{M} [x^{(n-1)}(s)]^{2} ds .$$

An application of Nehari's lemma [10, p. 431] now yields:

(2.4)
$$\int_{L}^{M} p(s) [x^{(n-2)}(s)]^{2} ds < \int_{L}^{M} [x^{(n-1)}(s)]^{2} ds ,$$

which implies

(2.5)
$$\int_{L}^{M} H(s, x(s)) x^{(n-2)}(s) ds > 0 ,$$

a contradiction to the negativeness of $x^{(n-2)}(t)$ on (L, M). Consequently, if $x^{(n-2)}(t)$ is oscillatory, we must have $x^{(n-2)}(t) \ge 0$ for all large t. This proves our assertion. To show the second conclusion, it suffices to observe that if H(t, x(t)) > 0 for $t \ge t_1$, then $x^{(n)}(t) < 0$ at each zero of $x^{(n-2)}(t)$. This concavity property of $x^{(n-2)}(t)$ implies that either $x^{(n-2)}(t)$ crosses the axis—impossible by the above argument—or $x^{(n-2)}(t)$ has a cusp at each one of its zeros—impossible, because it is differentiable there. Consequently, $x^{(n-2)}(t)$ cannot be oscillatory in this case. This completes the proof.

3. Main results. The following theorem improves Theorem 1 in Waltman's paper [13].

THEOREM 1. Let n = 3. Moreover, let p(t) be decreasing and H be increasing in u. Then if

$$\int_{T}^{\infty} H(t, \pm k) dt = \pm \infty$$

for every k > 0, every solution of (*) with a zero is oscillatory.

Proof. Let x(t) be a solution of (*) which has a zero at $t_1 \ge T$ and is such that x(t) > 0, $t \in (t_1, \infty)$. Assume further that $x(t) \ge L > 0$, $t \ge t_2 > t_1$, where L is a fixed constant. Then integration of (*) from t_2 to $t \ge t_2$ gives

$$egin{aligned} x''(t) &- x''(t_2) + p(t)x(t) - p(t_2)x(t_2) - \int_{t_2}^t x(s)d[p(s)] \ &\leq -\int_{t_2}^t H(s,\,L)ds \longrightarrow -\infty \quad ext{as} \quad t \longrightarrow \infty \ . \end{aligned}$$

Thus, $x''(t) \to -\infty$ as $t \to \infty$, a contradiction to the positivity of x(t). It follows that $\lim_{t\to\infty} \inf x(t) = 0$. The proof now follows exactly as in Waltman's theorem in [13], because we have just obtained that since $x(t_1) = 0$ and $\lim_{t\to\infty} \inf x(t) = 0$, there is a number $t_2 > t_1$ such that $x'(t_2) = 0$. We omit the rest of the proof.

Waltman assumed in [13] that p'(t) exists and is nonpositive and

eventually, for any constants A, B and any number $t_1 \ge T$, where $H(t, u) \equiv q(t)u^r$ with $Q(t) \ge 0$ and r = the quotient of two odd positive integers. Obviously, (3.1) implies the integral condition

$$\int_{t_1}^{\infty} q(s) ds = +\infty$$
 ,

which is actually equivalent to the integral assumption in Theorem 1 if H is of the above type. For another improvement of Waltman's theorem see Heidel's Corollary 3.4 in [2].

THEOREM 2. Let uH(t, u) > 0 for all $(t, u) \in R_r \times R$ with $u \neq 0$, and let H be increasing in u. Let the equation u'' + p(t)u = 0 be nonoscillatory with $[t^{\alpha}p(t)]$ decreasing in R_r , where $0 \leq \alpha \leq n - 1$. Then if

$$\int_{T}^{\infty} t^{\alpha} H(t, \pm k) dt = \pm \infty$$

for every k > 0, Equation (*) is BD-oscillatory.

Proof. Let x(t) be a bounded nonoscillatory solution of (*). Assume without loss of generality that x(t) > 0, $t \ge t_1 \ge T$. Then Theorem B implies that $x(t)x^{(n-2)}(t) \le 0$ or $x(t)x^{(n-2)}(t) \ge 0$ for all large t. Assume that $x^{(n-2)}(t) \ge 0$, $t \ge t_2 \ge t_1$. Then from (*) we obtain

(3.2)
$$x^{(n)}(t) + H(t, x(t)) \leq 0$$
, $t \geq t_2$.

Since

$$\int_{T}^{\infty} t^{n-1} H(t, \pm k) dt = \pm \infty$$
 ,

it follows from Theorem 2.9 of the author in [8], and the analogous theorem for n odd, that the equation (2.1) is *BD*-oscillatory. Theorem A implies now that x(t) cannot be eventually positive for n even, and that $\lim_{t\to\infty} x(t) = 0$ for n odd. However, this last conclusion is also impossible. In fact, since $x^{(n-2)}(t) \ge 0$, $t \ge t_2$ and n-2 is an odd integer, we must have $x'(t) \ge 0$ eventually. This contradicts $\lim_{t\to\infty} x(t) = 0$. Consequently, $x^{(n-2)}(t) \le 0$ for all large t, say for every $t \ge t_2 \ge \max\{1, t_1\}$. Now we multiply (*) by t^{α} and integrate from t_2 to t to obtain

$$(3.3) t^{\alpha} x^{(n-1)}(t) - t_{2}^{\alpha} x^{(n-1)}(t_{2}) - \alpha \int_{t_{2}}^{t} s^{\alpha-1} x^{(n-1)}(s) ds \\ + t^{\alpha} p(t) x^{(n-3)}(t) - t_{2}^{\alpha} p(t_{2}) x^{(n-3)}(t_{2}) \\ - \int_{t_{2}}^{t} x^{(n-3)}(s) d[s^{\alpha} p(s)] = - \int_{t_{2}}^{t} s^{\alpha} H(s, x(s)) ds$$

Now if $x^{(n-3)}(t_3) < 0$ for some $t_3 \ge t_2$, then $x^{(n-3)}(t) \le x^{(n-3)}(t_3) < 0$ for every $t \ge t_3$. This implies $\lim_{t\to\infty} x(t) = -\infty$, a contradiction. Thus, $x^{(n-3)}(t) \ge 0$ for $t \ge t_2$. Now since x(t) is bounded, it follows that $(-1)^j x^{(j)}(t) \le 0$ for n even and $(-1)^j x^{(j)}(t) \ge 0$ for n odd, and for $j = 1, 2, \dots, n-2, t \in [t_2, \infty)$. Assume that for some constant L > 0, $x(t) \ge L, t \in [t_2, \infty)$. Then from (3.3) we obtain

(3.4)
$$t^{\alpha}x^{(n-1)}(t) - \alpha \int_{t_2}^t s^{\alpha-1}x^{(n-1)}(s)ds \leq C - \int_{t_2}^t t^{\alpha}H(s, L)ds$$
.

Taking limits as $t \to \infty$ in (3.4) we obtain

(3.5)
$$\lim_{t\to\infty}\left[t^{\alpha}x^{(n-1)}(t)-\alpha\int_{t_2}^ts^{\alpha-1}x^{(n-1)}(s)ds\right]=-\infty.$$

Now let

(3.6)
$$\varphi(t) \equiv \int_{t_2}^t s^{\alpha-1} x^{(n-1)}(s) ds , \quad t \ge t_2 .$$

Then we have

$$(3.7) t\varphi'(t) - \alpha\varphi(t) = q(t) , t \ge t_2$$

where $\lim_{t\to\infty} q(t) = -\infty$. Now we apply Lemma 1 of Staikos and Sficas [12] to obtain

(3.8)
$$\int_{t_2}^{\infty} s^{\alpha-1} x^{(n-1)}(s) ds = \pm \infty .$$

The above integral cannot be $-\infty$ because then (3.5) would imply $x^{(n-1)}(t) \leq 0$ for all large t_1 which combined with $x^{(n-2)}(t) \leq 0$ gives $\lim_{t\to\infty} x(t) = -\infty$, a contradiction. Consequently, since $t_2 \geq 1$, we get

(3.9)
$$\int_{t_2}^{\infty} s^{\alpha-1} x^{(n-1)}(s) ds \leq \int_{t_2}^{\infty} s^{n-2} x^{(n-1)}(s) ds = + \infty .$$

Now the argument follows the steps of Theorem 1 in [4] and we omit the rest of it, which leads to the contradictions

(3.10)
$$\int_{t_2}^{\infty} x'(t) dt = \begin{cases} +\infty & n \text{ even} \\ -\infty & n \text{ odd} \end{cases}.$$

Consequently, no positive x(t) exists for n even and $x(t) \rightarrow 0$ monotonically as $t \rightarrow \infty$ for n odd. Dual arguments cover the assumption x(t) < 0. This completes the proof.

The following theorem characterizes further the solutions of (*) under slightly stronger assumptions than those of Theorem 2 and for n odd.

THEOREM 3. Let the assumptions of Theorem 2 be satisfied with p(t) decreasing in R_T and uH(t, u) > 0 for $u \neq 0$. Then, for n odd, every bounded solution of (*) with a zero is oscillatory.

Proof. Let x(t) be a bounded solution of (*) such that $x(t_1) = 0$ for $t_1 \ge T$ and x(t) > 0, $t \in (t_1, \infty)$. Theorem B implies that $x(t)x^{(n-2)}(t) > 0$ 0 or $x(t)x^{(n-2)}(t) < 0$ for all large t, and from the proof of Theorem 2 it follows that the second of these inequalities holds. Now since $x^{(n-2)}(t) < 0$ for all large t, say $t \ge t_2 \ge t_1$, we must have $x^{(n-3)}(t) > 0$ for all t $\geq t_2$. In fact, if $x^{(n-3)}(\overline{t}) \leq 0$ for some $\overline{t} \geq t_2$, then $x^{(n-3)}(t) < 0$ $x^{(n-3)}(\overline{t})$ for all $t \ge \overline{t}$, which implies the contradiction $x(t) \to -\infty$ as $t \rightarrow \infty$. Now $x^{(n-3)}(t)$ must have a zero to the right of t_1 . If this was not true then $x^{(n-3)}(t) > 0$ for $t \ge t_1$. This, along with $(-1)^j x^{(j)}(t) \ge 1$ 0, $j = 1, 2, \dots, n-2$ (cf. proof of Theorem 2), shows that none of the derivatives $x^{(j)}(t)$, $j = 1, 2, \dots, n-3$ can have a zero after t_1 . Thus, x'(t) < 0 for $t \in (t_1, \infty)$, a contradiction to the fact that $x(t_1) = 0$ and x(t) > 0, $t \in (t_1, \infty)$. Consequently, $x^{(n-3)}(t_3) = 0$ for some $t_3 > t_1$. We are planning to show that $x^{(n-3)}(t)$ is oscillatory. Suppose this is not true. Then let $t_4 \ge t_3$ be the last zero of $x^{(n-3)}(t)$. We must now have $x^{(n-3)}(t) > 0$, $t > t_4$. Now integrate (*) from t_4 to $t \ge t_4$, after multiplication by $x^{(n-3)}(t)$, to obtain

$$(3.11) x^{(n-1)}(t)x^{(n-3)}(t) + \frac{[x^{(n-2)}(t_4)]^2}{2} - \frac{[x^{(n-2)}(t)]^2}{2}$$

$$\begin{array}{l} + \ p(t) \ \frac{[x^{(n-3)}(t)]^2}{2} - \int_{t_4}^t \frac{[x^{(n-3)}(s)]^2}{2} d[p(s)] \\ + \ \int_{t_4}^t H(s, \ x(s)) x^{(n-3)}(s) ds = 0 \ . \end{array}$$

Now we either have $x^{(n-1)}(t) > 0$ for all large t, or $x^{(n-1)}(t)$ is oscillatory. The first of these possibilities combined with (3.11) implies that $\lim_{t\to\infty} x^{(n-2)}(t) = L < 0$, which is impossible because it implies $x^{(n-3)}(t) < 0$ for all large t. Now if $x^{(n-1)}(t)$ is oscillatory, then for some sequence of maxima $x^{(n-2)}(t_m)$ (with $t_m \to \infty$ as $m \to \infty$) we have $\lim_{m\to\infty} x^{(n-2)}(t_m) = 0$ and $x^{(n-1)}(t_m) = 0$. Letting $t = t_m$ in (3.11) we obviously get a contradiction because the limit on the left exists and is finite positive or $+\infty$ (p(t) is bounded and $\lim_{t\to\infty} x^{(n-3)}(t) = 0$). Consequently, $x^{(n-3)}(t)$ has to be oscillatory. However, we have already established that $x^{(n-2)}(t) < 0$ for all large t. This obviously implies a contradiction. Consequently, x(t) cannot be positive after it has a zero. A similar argument holds true if we assume that x(t) is negative for all large t. This completes the proof.

Results are expected of course to hold for all solutions of (*) under stronger assumptions on the function H(t, u). It turns out however that the method developed by the author in [4] does not apply in the present case due to the fact that we do not have enough information concerning the sign of $x^{(n-1)}(t)$ for a positive or negative solution of (*). Nevertheless, if the function p(t) is sufficiently small, we can treat the term $p(t)x^{(n-2)}(t)$ as a small perturbation because we can show that, under the conditions of Theorem B with uH(t, u) >0 for $u \neq 0$, the decrease of p(t) for all large t suffices to ensure that $\lim_{t\to\infty} x^{(n-1)}(t) = 0$ for any nonoscillatory solution x(t) of (*). This is the content of the following

THEOREM 4. Let the equation u'' + p(t)u = 0 by nonoscillatory. Moreover, let p(t) be decreasing in R_T and such that

$$\int_{T}^{\infty} t^{n} p(t) dt < +\infty$$
 .

Then (*) is D-oscillatory (BD-oscillatory) if the same is true for (2.1).

Proof. Let us first remark that $\lim_{t\to\infty} p(t) = 0$. In fact, if this is not true, then

which, by the classical Wintner's criterion [14], implies that all

solutions of u'' + p(t)u = 0 oscillate, a contradiction. Now let x(t) be an eventually positive solution of (*). Then the conclusion of Theorem B implies the existence of some $t_1 \ge T$ such that $x^{(n-2)}(t)x(t) > 0$ or $x^{(n-2)}(t)x(t) < 0$ for $t \ge t_1$. Now let the first of these inequalities hold, (2.1) be *D*-oscillatory and *n* even. Then it follows that x(t) is an eventually positive solution of

(3.12)
$$x^{(n)} + H(t, x) \leq 0$$

which, according to Theorem A, is a contradiction. Consequently, $x^{(n-2)}(t) < 0$ and x(t) > 0, $t \ge t_1$. Thus, we must have $x^{(n-3)}(t) > 0$, $t \ge t_1$. This happens because $x^{(n-3)}(t) < 0$ for all large t implies $x(t) \to -\infty$ as $t \to \infty$, and if $x^{(n-3)}(\tilde{t}) = 0$ for some $\tilde{t} \ge t_1$, then $x^{(n-3)}(\tilde{t}) < 0$ for all $t \ge \tilde{t}$, a contradiction again. Now integrate (*) from t_1 to $t \ge t_1$ to obtain

$$x^{(n-1)}(t) - x^{(n-1)}(t_1) + \int_{t_1}^t p(s) x^{(n-2)}(s) ds + \int_{t_1}^t H(s, x(s)) ds = 0$$
,

or

$$(3.13) x^{(n-1)}(t) - x^{(n-1)}(t_1) = -p(t)x^{(n-3)}(t) + p(t_1)x^{(n-3)}(t_1) + \int_{t_1}^t x^{(n-3)}(s)d[p(s)] - \int_{t_1}^t H(s, x(s))ds .$$

Now, as we noticed above, $\lim_{t\to\infty} p(t) = 0$, and $x^{(n-3)}(t)$ is bounded. Consequently, the first term on the right of (3.13) tends to zero as $t\to\infty$, and the two last terms are nonpositive and decreasing to finite limits, otherwise $x^{(n-1)}(t) \to -\infty$ as $t\to\infty$, a contradiction. It follows that $\lim_{t\to\infty} x^{(n-1)}(t) = L$ exists. If L > 0, then $x^{(n-2)}(t) \to +\infty$ as $t\to\infty$, a contradiction. If L < 0, then $x^{(n-2)}(t) \to -\infty$ as $t\to\infty$, a contradiction. Consequently,

$$(3.14) |x^{(n-2)}(t)| \leq |x^{(n-2)}(t_1)| + \int_{t_1}^t |x^{(n-1)}(s)| ds$$
$$\leq Mt$$

for all $t \ge t_1$ and some positive constant *M*. Now let $q(t) \equiv -p(t)x^{(n-2)}(t)$. Then the integral

$$-\int_t^\infty rac{(t-s)^{n-1}}{(n-1)!}q(s)ds < +\infty$$
 ,

and this follows easily from the integral assumption on p(t). If we let

$$P(t) \equiv -\int_t^{\infty} rac{(t-s)^{n-1}}{(n-1)!} q(s) ds$$
 ,

then $P^{(n)}(t) = q(t)$. Using the transformation $W(t) \equiv x(t) - P(t)$, $t \ge t_1$, we obtain

$$W^{(n)} + H(t, W + P(t)) = 0$$
, $t \ge t_1$.

Now the rest of the proof follows as in [6, Th. 2.1] for n even and we omit it. The case of odd n can be treated similarly and its proof is omitted. Similar considerations cover the case of bounded solutions under the assumption of *BD*-oscillation of (2.1), and we omit these arguments as well. This completes the proof.

The above theorem is intimately related to but does not contain any of the results of the third section of Heidel's paper [2]. We remark that Theorem 3 does not make use of all the hypotheses of Theorem 3, and that the conclusions in Theorem 3 can be deduced for any solution of (*) that has a zero if (2.1) is *D*-oscillatory. The proof of this last statement follows exactly as in Theorem 3 and we omit it.

In the following result we actually take into consideration the growth of the function H(t, u) with respect to the middle term.

THEOREM 5. Let (*) satisfy the following assumptions:

(i) u'' + p(t)u = 0 is nonoscillatory and p(t) is decreasing in R_{τ} ;

(ii) For every $\lambda > 0$, $\mu \ge 0$, there are no eventually positive (negative) solutions u(t) of the equation

(3.15)
$$u^{(n)} + H(t, u) - \mu t p(t) = 0$$
$$(u^{(n)} + H(t, u) + \mu t p(t) = 0)$$

such that $u(t) \geq \lambda$ ($u(t) \leq -\lambda$) and $H(t, u(t)) - \mu t p(t) \geq 0$ ($H(t, u(t)) - \mu t p(t) \leq 0$) for all large t. Moreover, if for a sequence $\{t_m\}, m = 1, 2, \cdots$ we have $\lim_{m\to\infty} t_m = \infty$ and $H(t_m, u_m) < \mu t_m p(t_m)(H(t_m, u_m) > - \mu t_m p(t_m))$ where $u_m > 0$ ($u_m < 0$) $m = 1, 2, \cdots$, then $\lim_{m\to\infty} \inf |u_n| = 0$. Then (*) is D-oscillatory.

Proof. Assume that there exists a solution x(t) of (*) which is eventually bounded below by a constant $\lambda > 0$. Then $x^{(n-2)}(t) \leq 0$ for all large t, and $\lim_{t\to\infty} x^{(n-1)}(t) = 0$. Consequently, there exists $t_1 \geq T$ such that

$$x(t) \geq \lambda > 0 \;, \;\;\; x^{(n-2)}(t) \leq 0 \;, \;\;\; -x^{(n-2)}(t) \leq \mu t$$

for every $t \geq t_1$. Consequently,

(3.16)
$$0 = x^{(n)}(t) + H(t, x(t)) + p(t)x^{(n-2)}(t)$$
$$\geq x^{(n)} + H(t, x(t)) - \mu t p(t) .$$

It follows from our assumptions that we cannot have $H(t, x(t)) \ge \mu tp(t)$ for all large t. If this was true, then Theorem A would imply that (3.15) has a positive solution such that $u'(t) \ge 0$ for all large t (again, this last assertion follows from Lemma 2.1 in [6] taking into consideration the class B(T, k) considered there). This is a contradiction. Thus, for some sequence $\{t_m\}$, $n = 1, 2, \cdots$ with $\lim_{m\to\infty} t_m = \infty$, we must have

$$H(t_m, x(t_m)) \leq \mu t_m p(t_m)$$
, $m = 1, 2, \cdots$,

which, by our assumptions, implies

(3.17)
$$\lim_{m\to\infty}\inf x(t_m)=\lim_{t\to\infty}\inf x(t)=0.$$

Now, taking into consideration that $x^{(n-2)}(t) \leq 0$, it follows that (3.17) is impossible for n even, and it implies $\lim_{t\to\infty} x(t) = 0$ for n odd. Similar consideration cover the case of a negative x(t). This completes the proof.

The conditions of the above theorem seem rather stringent at first glance. They are not however, and they imply oscillation criteria for all three important cases; sublinear, superlinear, linear, as the following corollary indicates. The method of proof of the above theorem was employed by the author in [7] where oscillations with perturbations Q(t, x) have been studied. A similar argument for a similar problem can be found in the paper [1] of Graef and Spikes.

COROLLARY 1. Consider the differential equation

$$(3.18) x^{(n)} + p(t)x^{(n-2)} + q(t)|x|^{\alpha} \operatorname{sgn} x = 0, \quad t \in R_T,$$

where p(t) is decreasing and such that the equation u'' + p(t)u = 0is nonoscillatory. Moreover, q(t) > 0, $t \in R_T$ and

(3.19)
$$\lim_{t \to \infty} [t p(t)/q(t)] = 0.$$

Then if α is a positive constant (3.18) is D-oscillatory provided one of the following conditions holds:

(i)
$$\int_{T}^{\infty} t^{n-1-\varepsilon}q(t)dt = +\infty$$
 for $\alpha = 1$ and some $\varepsilon > 0$;
(ii) $\int_{T}^{\infty} t^{\alpha(n-1)}q(t)dt = +\infty$ for $\alpha < 1$;
(iii) $\int_{T}^{\infty} t^{n-1}q(t)dt = +\infty$ for $\alpha > 1$.

One can easily see now that the conditions of Theorem 5 are

486

satisfied, in connection with the result of the author in [3]. Another corollary of the proof of Theorem 5 provides an appraisal of the nonoscillatory solutions of (*):

COROLLARY 2. Let the assumptions of Corollary 1 hold with (3.19) replaced by

$$[tp(t)/q(t)] \leq L$$
 (constant)

and the integral conditions replaced by the same conditions where $\pm q(t) - ktp(t)$ (for any k > 0) replaces q(t) and $\pm \infty$ replaces the second members. Then for every nonoscillatory solution x(t) of (*) we have

$$\lim_{t o\infty} |x(t)| = \lim_{t o\infty} \inf |x(t)| = K \leq (kL)^{1/lpha}$$

for some constant k > 0 such that $|x^{(n-2)}(t)| \leq kt$ eventually.

In Theorems 2-5 we made the basic assumption that u'' + p(t)u = 0is nonoscillatory. Let us first notice that if this equation is oscillatory, then $x^{(n-2)}(t)x(t) \ge 0$ $(x^{(n-2)}(t)x(t) \le 0)$ is impossible for a positive (negative) nonoscillatory solution of (*) and all large t. In fact, assume that x(t) > 0 for all large t. Then from (*) it follows that

$$(3.20) u'' + p(t)u \leq 0$$

for all large t, where $u(t) \equiv x^{(n-2)}(t)$. If we assume the first of the above inequalities, then (3.20) has a positive solution. By Theorem A this is impossible. Similarly, the second of the above inequalities is impossible for a negative x(t). Results can be now formulated analogous to several third order results in the third section of Heidel's paper [2]. We shall undertake this task in a future paper.

It would be very interesting to have some results ensuring the fact that $x^{(n-1)}(t)$ is of fixed sign for any nonoscillatory solution x(t) of (*). Such a result would definitely permit relaxation of the conditions imposed in Theorem 5 and would make possible the extension of Theorem 2 to all solutions of (*).

References

^{1.} J. R. Graef, P. W. Spikes, Asymptotic properties of solutions of functional differential equations of arbitrary order, J. Math. Anal. Appl., to appear.

^{2.} J. W. Heidel, Qualitative behaviour of solutions of a third order nonlinear differential equation, Pacific J. Math., 27 (1968), 507-526.

^{3.} A. G. Kartsatos, Criteria for oscillation of solutions of differential equations of arbitrary order, Proc. Japan Acad., 44 (1968), 599-602.

^{4.} _____, On oscillations of even order nonlinear differential equations, J. Diff. Equations, **6** (1969), 232-237.

5. _____, Contributions to the research of the oscillation and the asymptotic behaviour of solutions of ordinary differential equations, Bull. Soc. Math. Grèce, **10** (1969), 1-48 (Greek).

6. ____, On nth-order differential inequalities, J. Math. Anal. Appl., 52 (1975), 1-9.

7. _____, Oscillation and nonoscillation for perturbed differential equations, to appear.

8. _____, Recent results on oscillation of solutions of forced and perturbed nonlinear differential equations of even order, Proc. NSF-CBMS Reg. Conference, Stability of Dynamical Systems, Mississippi State University, 1975, to appear.

9. G. E. Liossatos, Qualitative behaviour of solutions of nonlinear differential equations of order n, Dissertation, Univ. of Athens, Doct. Athens, Greece, 1973.

10. Z. Nehari, Oscillation criteria for second order linear differential equations, Trans, Amer. Math. Soc., 2 (1957), 428-445.

11. H. Onose, A comparison theorem and the forced oscillation, Bull. Austral. Math. Soc., 13 (1975), 13-19.

12. V. A. Staikos, Y. G. Sficas, Forced oscillations for differential equations of arbitrary order, J. Diff. Equations, 17 (1975), 1-11.

13. P. Waltman, Oscillation criteria for third order nonlinear differential equations, Pacific J. Math., **18** (1966), 385-389.

14. A. Wintner, A criterion of oscillatory stability, Quart. Appl. Math., 7 (1949), 115-117.

Received November 25, 1975.

UNIVERSITY OF SOUTH FLORIDA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor) University of California Los Angeles, California 90024

R. A. BEAUMONT

University of Washington Seattle, Washington 98105 J. Dugundji

Department of Mathematics University of Southern California Los Angeles, California 90007

D. GILBARG AND J. MILGRAM Stanford University Stanford, California 94305

ASSOCIATE EDITORS

E.F. BECKENBACH

B. H. NEUMANN

F. WOLF K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF HAWAII UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON * * *

AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),

8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics Manufactured and first issued in Japan

Pacific Journal of Mathematics Vol. 67, No. 2 February, 1976

Patricia Andresen and Marvin David Marcus, Weyl's inequality and quadratic forms on the Grassmannian	277
George Bachman and Alan Sultan, <i>Regular lattice measures: mappings and spaces</i>	291
David Geoffrey Cantor, On certain algebraic integers and approximation by rational functions with integral coefficients	323
James Richard Choike, On the value distribution of functions meromorphic in the unit disk with a spiral asymptotic value	339
David Earl Dobbs, <i>Divided rings and going-down</i>	353
Mark Finkelstein and Robert James Whitley, Integrals of continuous	
functions	365
Ronald Owen Fulp and Joe Alton Marlin, Integrals of foliations on	
manifolds with a generalized symplectic structure	373
Cheong Seng Hoo, Principal and induced fibrations	389
Wu-Chung Hsiang and Richard W. Sharpe, Parametrized surgery and	
isotopy	401
Surender Kumar Jain, Surjeet Singh and Robin Gregory Symonds, <i>Rings</i> whose proper cyclic modules are quasi-injective	461
Pushpa Juneja, On extreme points of the joint numerical range of commuting normal operators.	473
Athanassios G. Kartsatos, <i>Nth order oscillations with middle terms of order</i> N-2	477
John Keith Luedeman, The generalized translational hull of a semigroup	489
Louis Jackson Ratliff. Jr., <i>The altitude formula and DVR's</i>	509
Ralph Gordon Stanton, C. Sudler and Hugh C. Williams, <i>An upper bound</i>	
for the period of the simple continued fraction for \sqrt{D}	525
David Westreich <i>Global analysis and periodic solutions of second order</i>	
systems of nonlinear differential equations	537
David Lee Armacost Correction to: "Compactly cogenerated LCA	
groups"	555
Jerry Malzan, Corrections to: "On groups with a single involution"	555
David Westreich, Correction to: "Bifurcation of operator equations with	
unbounded linearized part"	555
ν Γ	