

Pacific Journal of Mathematics

ON PUNCTURED BALLS IN MANIFOLDS

WOLFGANG H. HEIL

ON PUNCTURED BALLS IN MANIFOLDS

WOLFGANG HEIL

E. Brown showed that for any map f of a punctured disc B_n with n holes into a 2-manifold M that is an embedding of ∂B_n , there is an embedding g of a punctured disk B_k into M such that $g(\partial B_k)$ is a subcollection of $f(\partial B_n)$. In this paper E. Brown's approach is extended to show that a similar result holds for maps of punctured q -balls into certain q -manifolds ($q \geq 3$).

Let $PC(q)$ denote the collection of (topological) q -manifolds M^q with the property that if h is an embedding of $S^{q-1} \times [0, 1]$ into M^q that is null homotopic, then $h(S^{q-1} \times \frac{1}{2})$ bounds a topological q -cell in M^q .

Note that $PC(1)$ and $PC(2)$ consist of all 1-manifolds and 2-manifolds, respectively. It is well-known that $PC(3)$ consists of all 3-manifolds provided the Poincaré conjecture is true in dimension 3. Since the generalized Poincaré conjecture holds for dimensions ≥ 5 , [2] we are led to conjecture that $PC(q)$ consists of all (topological) q -manifolds for $q \geq 5$, particularly since, from the proposition below, if $h: S^{q-1} \rightarrow \partial M^q$ is an embedding such that $h(S^{q-1})$ is null-homotopic in M^q , then M^q is indeed a q -cell ($q \geq 5$). However, C. McA. Gordon, whom I would like to thank most sincerely for providing the proof of the following proposition, informs me that C. T. C. Wall and John Morgan have counter examples for $q > 4$.

PROPOSITION. *Let $C \cong S^{q-1}$ be a boundary component of a compact q -manifold M . If $[C] = 0$ in $\pi_{q-1}(M)$, then M is contractible.*

Proof. Let $q \geq 3$. By the Whitehead and Hurewicz Theorems it suffices to show that $\pi_1(M) = 1$ and $H_*(M) = 0$. Now $\partial M = C$ since otherwise $[C] \neq 0$ in $H_{q-1}(M)$. Also, M is orientable since otherwise for the orientation cover M' of M we have $\partial M' = C' \cup C''$ (copies over C) and $[C'] = 0$ in $\pi_{q-1}(M')$, a contradiction.

There is a map $f: (B^q, S^{q-1}) \rightarrow (M, \partial M)$ such that $f|S^{q-1}$ is a homeomorphism. Orient M so that f has degree 1. Then for the fundamental classes z_q, w_q in $H_q(B^q, S^{q-1}), H_q(M, \partial M)$, resp., we have $f^*(z_q) = w_q$ and a commutative diagram

$$\begin{array}{ccc}
 H^{q-k}(B^q, S^{q-1}) & \xleftarrow{f^*} & H^{q-k}(M, \partial M) \\
 \downarrow \cap z_q & & \downarrow \cap w_q \\
 H_k(B^q) & \xrightarrow{f_*} & H_k(M)
 \end{array}$$

By Lefschetz duality, the vertical maps are isomorphisms. Therefore $f_*(- \cap z_q)f_*$ is an isomorphism. It follows that f_* is onto and hence that $H_*(M) = 0$.

To show that $\pi_1(M) = 1$, let $p: \tilde{M} \rightarrow M$ be the universal covering. Then f lifts to $\tilde{f}: (B^q, S^{q-1}) \rightarrow (\tilde{M}, \partial\tilde{M})$. But $1 = \deg(f) = \deg(p \circ \tilde{f}) = (\deg p)(\deg \tilde{f})$, hence $\deg p = \pm 1$ and $\pi_1(M) = 1$.

For $q \geq 2$, $n \geq 1$, let B_n^q be a punctured q -ball with $n - 1$ holes, i.e., B_n^q is obtained from S^q by removing the interiors of n mutually disjoint q -balls.

For a bicollared $S^{q-1} \subset M^q$ let $N \approx S^{q-1} \times I$ be a neighborhood of S^{q-1} and let $M' = \text{cl}(M - N) \cup B' \cup B''$, where the boundaries of the q -balls B' , B'' are attached to the boundary components $S^{q-1} \times 0$ and $S^{q-1} \times 1$ of $\text{cl}(M - N)$. We say M' is obtained from M by *surgery along* S^{q-1} . Let X be the space obtained from M' by identifying B' and B'' under a homeomorphism. Note that X can be obtained from M^q by attaching a q -ball B to S^{q-1} along its boundary and $X - B = M' - (B' \cup B'') = M - S^{q-1}$.

LEMMA. *Let S be a $(q - 1)$ -sphere in $X - B$. If $S \approx 0$ in X , then $S \approx 0$ in M' .*

Proof. Suppose S^{q-1} separates M into M_1 and M_2 ; then $M' = M'_1 \cup M'_2$, where $M'_1 = M_1 \cup B'$, $M'_2 = M_2 \cup B''$. Let X'_i be obtained from M_i by collapsing S^{q-1} to a point. The projection $p: X \rightarrow X'_1 \vee X'_2$ is a homotopy equivalence which sends S into X'_1 , say. This can be seen as follows: Identify a neighborhood of S^{q-1} with $N = S^{q-1} \times [-1, 1]$, where $S^{q-1} = S^{q-1} \times \{0\}$. Let w be the "centerpoint" of B and for $y \in S^{q-1}$ let $r(y)$ be the "radius" in B from y to w . In $X'_1 \vee X'_2$ we identify $p(N) = (S^{q-1} \times I)/(S^{q-1} \times \{0\})$ with the cones over $S^{q-1} \times \{-1\}$ and $S^{q-1} \times \{1\}$ wedged together at their vertices to a vertex v . Let $g: X'_1 \vee X'_2 \rightarrow X$ be the map that is the identity outside $p(N)$ and which sends the join of x and v (for $x \in S^{q-1} \times \{-1\}$, respectively $S^{q-1} \times \{1\}$) linearly to $x \times [-1, 0] \cup r(x \times \{0\})$, resp. $x \times [0, 1] \cup r(x \times 0)$. Then it is clear that g is a homotopy inverse of p . But since X'_1 is a retract of $X'_1 \vee X'_2$ it follows that $S \approx 0$ in X'_1 already and hence in $M'_1 \approx X'_1$.

If S^{q-1} does not separate M , let $\tilde{X} \rightarrow X$ be the infinite cyclic covering of X determined by B : the q -ball B lifts to q -balls $\cdots B_{-1}, B_0, B_1, \cdots$ and each component of $\tilde{X} - \bigcup_{i=-\infty}^{\infty} B_i$ maps homeomorphically onto $X - B$. For each i , let X'_i be obtained from M' by collapsing B' and B'' to single points. There is a projection $\tilde{X} \rightarrow \bigvee_{i=-\infty}^{\infty} X'_i$ that is a homotopy equivalence and hence $\pi_{q-1}(X'_j)$ injects into $\pi_{q-1}(\tilde{X})$, for each j . Let \tilde{S} be a lift of S to \tilde{X} . Then \tilde{S} lies in a component of $\tilde{X} - \bigcup B_i$ and is mapped into a factor X'_j of $\bigvee X'_i$. It follows that $\tilde{S} \approx 0$ in X'_j , hence $S \approx 0$ in M' .

THEOREM. *Let $f: B_n^q \rightarrow M^q$ be a map such that $f|_{\partial B_n^q}$ is a bicollared embedding, $f(\partial B_n^q) = S_1 \cup \cdots \cup S_n$. Suppose that the manifold M' obtained from M^q by surgery along S_i ($i = 2, \cdots, n$) belongs to $PC(q)$. Then some subcollection of $\{S_1, \cdots, S_n\}$ contains S_1 and bounds an embedded punctured q -ball in M .*

Proof. By Brown's result we can assume that $q \geq 3$. Let X be obtained from M by attaching q -balls B_i to S_i ($i = 2, \cdots, n$) along their boundaries. Then $X - \bigcup_{i=2}^n B_i = M' - \bigcup_{i=2}^n B'_i \cup \bigcup_{i=2}^n B''_i$, where B'_i, B''_i are the balls used for surgery on S_i . Now $S_1 \approx 0$ in X . By the lemma, $S_1 \approx 0$ in M' . Since $M' \in PC(q)$, S_1 bounds a q -ball B_* in M' . Let E be the component of $B_* - \bigcup_{i=2}^n (B'_i \cup B''_i)$ which has S_1 on its boundary. If for each $i = 2, \cdots, n$ only one of $\partial B'_i, \partial B''_i \subset \partial E$, then E is the desired punctured ball in M bounded by S_1 and some of the S_i 's. In fact, this is the only case that can happen. For suppose for some i , $\partial B'_i$ and $\partial B''_i \subset \partial E$. Then let k be a simple arc in E from a point of $\partial B'_i$ to a point on $\partial B''_i$ such that k misses the other ∂B_j 's and such that k corresponds to a simple closed curve in M that intersects S_i in one point and misses the other S_j 's. In M , the intersection numbers $\#(k, S_i) = \pm 1$, but $\#(k, \Sigma_{j \neq i} S_j) = 0$, which is impossible since $S_i \sim \bigcup_{j \neq i} S_j$.

REFERENCES

1. E. M. Brown, *A note on punctured disks in a 2-manifold*, Proc. Amer. Math. Soc., **22** (1969), 471.
2. M. H. A. Newman, *The engulfing theorem for topological manifolds*, Ann. of Math., **84** (1966), 555-571.

Received February 6, 1976 and in revised form July 28, 1976.

FLORIDA STATE UNIVERSITY
TALLAHASSEE, FL 32306

Pacific Journal of Mathematics

Vol. 68, No. 1

March, 1977

Richard Julian Bagby, <i>On L^p, L^q multipliers of Fourier transforms</i>	1
Robert Beauwens and Jean-Jacques Van Binnebeek, <i>Convergence theorems in Banach algebras</i>	13
James Cyril Becker, <i>Skew linear vector fields on spheres in the stable range</i>	25
Michael James Beeson, <i>Continuity and comprehension in intuitionistic formal systems</i>	29
James K. Deveney, <i>Generalized primitive elements for transcendental field extensions</i>	41
Samuel S. Feder, Samuel Carlos Gitler and K. Y. Lam, <i>Composition properties of projective homotopy classes</i>	47
Nathan Jacob Fine, <i>Tensor products of function rings under composition</i>	63
Benno Fuchssteiner, <i>Iterations and fixpoints</i>	73
Wolfgang H. Heil, <i>On punctured balls in manifolds</i>	81
Shigeru Itoh, <i>A random fixed point theorem for a multivalued contraction mapping</i>	85
Nicolas P. Jewell, <i>Continuity of module and higher derivations</i>	91
Roger Dale Konyndyk, <i>Residually central wreath products</i>	99
Linda M. Lesniak and John A. Roberts, <i>On Ramsey theory and graphical parameters</i>	105
Vo Thanh Liem, <i>Some cellular subsets of the spheres</i>	115
Dieter Lutz, <i>A perturbation theorem for spectral operators</i>	127
P. H. Maserick, <i>Moments of measures on convex bodies</i>	135
Stephen Joseph McAdam, <i>Unmixed 2-dimensional local domains</i>	153
D. B. McAlister and Norman R. Reilly, <i>E-unitary covers for inverse semigroups</i>	161
William H. Meeks, III and Julie Patrusky, <i>Representing codimension-one homology classes by embedded submanifolds</i>	175
Premalata Mohapatro, <i>Generalised quasi-Nörlund summability</i>	177
Takahiko Nakazi, <i>Superalgebras of weak-*Dirichlet algebras</i>	197
Catherine Louise Olsen, <i>Norms of compact perturbations of operators</i>	209
William Henry Ruckle, <i>Absolutely divergent series and isomorphism of subspaces. II</i>	229
Bernard Russo, <i>On the Hausdorff-Young theorem for integral operators</i>	241
Arthur Argyle Sagle and J. R. Schumi, <i>Anti-commutative algebras and homogeneous spaces with multiplications</i>	255
Robert Evert Stong, <i>Stiefel-Whitney classes of manifolds</i>	271
D. Suryanarayana, <i>On a theorem of Apostol concerning Möbius functions of order k</i>	277
Yoshio Tanaka, <i>On closedness of C- and C^*-embeddings</i>	283
Chi Song Wong, <i>Characterizations of certain maps of contractive type</i>	293