

Pacific Journal of Mathematics

ON RAMSEY THEORY AND GRAPHICAL PARAMETERS

LINDA M. LESNIAK AND JOHN A. ROBERTS

ON RAMSEY THEORY AND GRAPHICAL PARAMETERS

LINDA LESNIAK-FOSTER AND JOHN ROBERTS

A graph G is said to have a factorization into the subgraphs G_1, \dots, G_k if the subgraphs are spanning, pairwise edge-disjoint, and the union of their edge sets equals the edge set of G . For a graphical parameter f and positive integers n_1, n_2, \dots, n_k ($k \geq 1$), the f -Ramsey number $f(n_1, n_2, \dots, n_k)$ is the least positive integer p such that for any factorization $K_p = \bigcup_{i=1}^k G_i$, it follows that $f(G_i) \geq n_i$ for at least one i , $1 \leq i \leq k$. In the following, we present two results involving f -Ramsey numbers which hold for various vertex and edge partition parameters, respectively. It is then shown that the concept of f -Ramsey number can be generalized to more than one vertex partition parameter, more than one edge partition parameter, and combinations of vertex and edge partition parameters. Formulas are presented for these generalized f -Ramsey numbers and specific illustrations are given.

1. Introduction. A subgraph H of a graph G is called *spanning* if H has the same vertex set as G . A graph G is said to have a *factorization* into the subgraphs G_1, G_2, \dots, G_k , written $G = \bigcup_{i=1}^k G_i$, if the subgraphs are spanning, pairwise edge-disjoint, and the union of their edge sets equals the edge set of G . It is permissible for a subgraph G_i to be empty; i.e., have no edges.

Let f be a graphical parameter, and let n_1, n_2, \dots, n_k , ($k \geq 1$) be positive integers. In [2], Chartrand and Polimeni defined the f -Ramsey number $f(n_1, n_2, \dots, n_k)$ as the least positive integer p such that for any factorization $K_p = \bigcup_{i=1}^k G_i$ of the complete graph of order p , it follows that $f(G_i) \geq n_i$ for at least one subgraph G_i , $1 \leq i \leq k$. If $\omega(G)$ is the maximum order among the complete subgraphs of G , then the ω -Ramsey number is the ordinary Ramsey number (see [3; p. 16]) in k variables.

The chromatic number $\chi(G)$ of a graph G is the minimum number of colors which may be assigned to the vertices of G so that adjacent vertices are assigned different colors. The vertex-arboricity $a(G)$ of G is the minimum number of subsets into which the vertex set of G may be partitioned so that each subset induces an acyclic subgraph. Chartrand and Polimeni [2] gave formulas for the χ -Ramsey numbers and the a -Ramsey numbers. We present a result which holds for several "partition" parameters (including chromatic number and vertex-

arboricity as special cases). Furthermore, it is shown that the concept of f -Ramsey number can be generalized to more than one parameter. Formulas are presented for these generalized f -Ramsey numbers, and specific illustrations are given involving chromatic number, edge chromatic number, vertex-arboricity, and arboricity.

2. Vertex partition parameters. A graphical property ρ will be called *co-hereditary* if (1) every subgraph of a graph having property ρ has property ρ and (2) the graph consisting of disjoint graphs, each having property ρ , has property ρ .

Let ν be a graphical property which the trivial graph K_1 possesses. We define the *vertex partition number* $\nu(G)$ of a graph G as the minimum number of subsets into which the vertex set of G can be partitioned so that each subset induces a subgraph having property ν . Clearly, $\nu(G) = 1$ if and only if G has property ν . The *limit* $\lim \nu$ of a vertex partition parameter ν is defined as $\lim \nu = \lim_{n \rightarrow \infty} \nu(K_n)$, provided this limit exists. We write $\lim \nu = \infty$ if $\nu(K_n) \rightarrow \infty$ as $n \rightarrow \infty$. We assume that all properties ν under discussion are co-hereditary and that $\lim \nu = \infty$. It is a consequence of the definitions that $\nu(H) \leq \nu(G)$ if H is a subgraph of G , for such properties ν .

For positive integers n_1, n_2, \dots, n_k and vertex partition parameter ν , the ν -*Ramsey number* $\nu(n_1, n_2, \dots, n_k)$ is the least positive integer p such that given any factorization $K_p = \bigcup_{i=1}^k G_i$, it follows that $\nu(G_i) \geq n_i$ for at least one i , $1 \leq i \leq k$. Since $\lim \nu = \infty$, for each i , $1 \leq i \leq k$, there exists a positive integer m_i such that $\nu(K_{m_i}) \geq n_i$. Hence, since ν is co-hereditary, $\nu(n_1, n_2, \dots, n_k)$ exists and is bounded above by $r(m_1, m_2, \dots, m_k)$, the Ramsey number in the k variables m_1, m_2, \dots, m_k . We also note that $\nu(n_1, n_2, \dots, n_k)$ is symmetric in n_1, n_2, \dots, n_k .

There are properties ν which are not co-hereditary and for which $\lim \nu \neq \infty$ such that $\nu(n_1, n_2, \dots, n_k)$ does not exist for certain positive integers n_1, n_2, \dots, n_k . For example, if ν denotes the property of being connected, then $\nu(G)$ is the number of components of the graph G . Then $\nu(3, 3)$ does not exist since for every positive integer p , there exists a factorization $K_p = G_1 \cup G_2$ such that neither G_1 nor G_2 has more than two components.

For a vertex partition parameter ν and positive integer k , let $\bar{\nu}(k)$ denote the largest integer p for which there exists a factorization $K_p = \bigcup_{i=1}^k G_i$ such that $\nu(G_i) = 1$ for $i = 1, 2, \dots, k$. Then we have the following lemma.

LEMMA 1. *If ν is a vertex partition parameter for which $\lim \nu = \infty$ and the corresponding property ν is co-hereditary, then $\bar{\nu}(k)$ exists for every positive integer k .*

Proof. Since $\nu(K_1) = 1$ and $K_1 = \bigcup_{i=1}^k K_i$, it follows that $\bar{\nu}(k) \geq 1$ for each positive integer k . Since $\lim \nu = \infty$, there exists a positive integer p such that $\nu(K_p) \geq 2$. Hence, if we consider an arbitrary factorization $K_m = \bigcup_{i=1}^k G_i$, where $m = r(p, p, \dots, p)$ is the Ramsey number in k variables, then G_i contains K_p as a subgraph for at least one i , $1 \leq i \leq k$, say $i = j$. Therefore, $\nu(G_j) \geq \nu(K_p) \geq 2$, which implies that $\bar{\nu}(k)$ exists and, in fact, $\bar{\nu}(k) < m$.

We can now present a formula for any ν -Ramsey number.

THEOREM 1. *Let n_1, n_2, \dots, n_k ($k \geq 1$) be positive integers, and let ν be a vertex partition parameter for which the corresponding property ν is co-hereditary and $\lim \nu = \infty$. Then*

$$\nu(n_1, n_2, \dots, n_k) = 1 + \bar{\nu}(k) \cdot \prod_{i=1}^k (n_i - 1).$$

Proof. If $n_i = 1$ for some $i = 1, 2, \dots, k$, then $\nu(n_1, n_2, \dots, n_k) = 1$ and the theorem follows. Thus, we assume that $n_i \geq 2$ for each i , $1 \leq i \leq k$, and let

$$p = 1 + \bar{\nu}(k) \cdot \prod_{i=1}^k (n_i - 1).$$

First, we verify the inequality $\nu(n_1, n_2, \dots, n_k) \leq p$. Assume that this is not the case. Then there exists a factorization $K_p = \bigcup_{i=1}^k G_i$ such that $\nu(G_i) \leq n_i - 1$ for $i = 1, 2, \dots, k$. For each G_i , $i = 1, 2, \dots, k$, let the vertex set $V(G_i)$ be partitioned into $\nu(G_i)$ classes so that for each " ν -class" α , $\nu(\langle \alpha \rangle) = 1$, where $\langle \alpha \rangle$ denotes the subgraph induced by the class α . Then G_1 has a ν -class α_1 containing at least $1 + \bar{\nu}(k) \prod_{i=2}^k (n_i - 1)$ vertices, G_2 has a ν -class α_2 containing at least $1 + \bar{\nu}(k) \cdot \prod_{i=3}^k (n_i - 1)$ vertices of α_1 , and, in general, for $1 \leq l < k$, if $\alpha_1, \alpha_2, \dots, \alpha_l$ are ν -classes, respectively, of G_1, G_2, \dots, G_l for which

$$\left| \bigcap_{i=1}^l \alpha_i \right| \geq 1 + \bar{\nu}(k) \cdot \prod_{i=l+1}^k (n_i - 1),$$

then G_{l+1} has a ν -class α_{l+1} such that

$$\left| \bigcap_{i=1}^{l+1} \alpha_i \right| \geq 1 + \bar{\nu}(k) \cdot \prod_{i=l+2}^k (n_i - 1).$$

Hence, each G_i , $1 \leq i \leq k$, has a ν -class α_i such that $|\bigcap_{i=1}^k \alpha_i| \geq 1 + \bar{\nu}(k)$. Let \mathcal{U} be a set of $1 + \bar{\nu}(k)$ vertices in $\bigcap_{i=1}^k \alpha_i$, and define H_i to be the subgraph in G_i induced by \mathcal{U} for $i = 1, 2, \dots, k$. Then

$$K_{1+\bar{\nu}(k)} = H_1 \cup H_2 \cup \dots \cup H_k,$$

where $\nu(H_i) = 1$ for $1 \leq i \leq k$ since ν is co-hereditary. This, however, is impossible. Therefore, $\nu(n_1, n_2, \dots, n_k) \leq p$.

In order to show that $\nu(n_1, n_2, \dots, n_k) = p$, it suffices to exhibit a factorization $K_{p-1} = \bigcup_{i=1}^k G_i$ such that $\nu(G_i) \leq n_i - 1$ for $i = 1, 2, \dots, k$. Let $r = \prod_{i=1}^k (n_i - 1)$, and consider r pairwise disjoint copies of $K_{\bar{\nu}(k)}$, labeled $K_{\bar{\nu}(k)}^1, K_{\bar{\nu}(k)}^2, \dots, K_{\bar{\nu}(k)}^r$. By definition of $\bar{\nu}(k)$, there exists a factorization $K_{\bar{\nu}(k)} = \bigcup_{i=1}^k F_i$ such that $\nu(F_i) = 1$ for $i = 1, 2, \dots, k$. We denote by F_{il} the F_i contained in $K_{\bar{\nu}(k)}^l$, $l = 1, 2, \dots, r$ and $i = 1, 2, \dots, k$. With each of the r k -tuples (c_1, c_2, \dots, c_k) , $1 \leq c_j \leq n_j - 1$ and $1 \leq j \leq k$, we identify a complete graph $K_{\bar{\nu}(k)}^l$, $l = 1, 2, \dots, r$, in such a way that the identification is one-to-one. Then, for each $i = 1, 2, \dots, k$ and $l = 1, 2, \dots, r$, we associate with F_{il} the k -tuple identified with $K_{\bar{\nu}(k)}^l$. Let the graph G_i ($i = 1, 2, \dots, k$) consist of the graphs $F_{i1}, F_{i2}, \dots, F_{ir}$ where a vertex of F_{is} is adjacent to each vertex of F_{it} if and only if the i th coordinate is the first coordinate in which their associated k -tuples differ. It then follows that $K_{p-1} = \bigcup_{i=1}^k G_i$. For each $i = 1, 2, \dots, k$, define $V_{i,j}$ to consist of the set of all vertices v such that v is a vertex of an F_{il} whose associated k -tuple (c_1, c_2, \dots, c_k) has $c_i = j$; $j = 1, 2, \dots, n_i - 1$. Then $\{V_{i1}, V_{i2}, \dots, V_{i,n_i-1}\}$ is a partition of $V(G_i)$ for which the subgraph $\langle V_{i,j} \rangle$ consists of $r/(n_i-1)$ pairwise disjoint copies of $F_{i,j}$, $j = 1, 2, \dots, n_i - 1$. Hence, $\nu(V_{i,j}) = 1$ for each such j , which implies that $\nu(G_i) \leq n_i - 1$ for $i = 1, 2, \dots, k$.

For the chromatic number χ , it follows that $\bar{\chi}(k) = 1$ for all $k \geq 1$. Hence, we obtain an immediate corollary.

COROLLARY 1a. (Chartrand and Polimeni [2]). *If n_1, n_2, \dots, n_k are positive integers, then*

$$\chi(n_1, n_2, \dots, n_k) = 1 + \prod_{i=1}^k (n_i - 1).$$

The *edge-arboricity* $a_1(G)$ of a graph G is the minimum number of subsets in a partition of the edge set of G such that each subset induces an acyclic subgraph. For the *vertex-arboricity* $a(G)$ of G , we have another corollary.

COROLLARY 1b. (Chartrand and Polimeni [2]). *If n_1, n_2, \dots, n_k are positive integers, then*

$$a(n_1, n_2, \dots, n_k) = 1 + 2k \cdot \prod_{i=1}^k (n_i - 1).$$

Proof. Again, it suffices to evaluate $\bar{a}(k)$. First, we observe that if there is a factorization $K_p = \bigcup_{i=1}^k G_i$, where $a(G_i) = 1$ for $i = 1, 2, \dots, k$, then $k \geq a_1(K_p) = \lfloor p/2 \rfloor$. Since $p = 2k$ is the largest such integer, we have $\bar{a}(k) = 2k$ and the desired result.

As one further illustration of Theorem 1, we consider the 2-chromatic number $\chi^{(2)}(G)$ of a graph G (see 1), defined as the least number of subsets in any partition of $V(G)$ such that the subgraph induced by each subset contains no path of length two. Also, we define the edge chromatic number $\chi_1(G)$ of G as the least number of colors needed to color the edges of G so that adjacent edges are colored differently.

COROLLARY 1c. For positive integers n_1, n_2, \dots, n_k ,

$$\chi^{(2)}(n_1, n_2, \dots, n_k) = 1 + 2 \left\lfloor \frac{k}{2} \right\rfloor \cdot \prod_{i=1}^k (n_i - 1).$$

Proof. To determine $\bar{\chi}^{(2)}(k)$, it is equivalent to determine the largest integer n such that $\chi_1(K_n) \leq k$. Since $\chi_1(K_p) = p$ if p is odd and $\chi_1(K_p) = p - 1$ if p is even, it follows that $n = 2\lfloor k/2 \rfloor$, which gives the desired result.

The concept of the ν -Ramsey number can be generalized. Let $\nu_1, \nu_2, \dots, \nu_k$ be vertex partition parameters where again we assume the corresponding properties are co-hereditary and $\lim \nu_i = \infty$ for each i , $1 \leq i \leq k$. Then we define the $(\nu_i)_i^k$ -Ramsey number $(\nu_i)_i^k(n_1, n_2, \dots, n_k)$ as the least positive integer p such that given any factorization $K_p = \bigcup_{i=1}^k G_i$, it follows that $\nu_i(G_i) \geq n_i$ for at least one i , $1 \leq i \leq k$. Following an earlier argument we note that $(\nu_i)_i^k(n_1, n_2, \dots, n_k)$ exists since each ν_i is co-hereditary and $\lim \nu_i = \infty$. In this case, we do not have symmetry in the k -variables n_1, n_2, \dots, n_k ; however, it does follow that

$$(\nu_{i_j})_{j=1}^k(n_{i_1}, n_{i_2}, \dots, n_{i_k}) = (\nu_i)_i^k(n_1, n_2, \dots, n_k),$$

where i_1, i_2, \dots, i_k is any permutation of $1, 2, \dots, k$.

For vertex partition parameters $\nu_1, \nu_2, \dots, \nu_k$, we define $(\bar{\nu}_i)_i^k(k) = (\bar{\nu})_i^k$ to be the largest integer p such that there exists a factorization $K_p = \bigcup_{i=1}^k G_i$ with $\nu_i(G_i) = 1$ for $i = 1, 2, \dots, k$. Using an argument similar to that given in Lemma 1, one can show that $(\bar{\nu}_i)_i^k$ exists, and moreover, a technique analogous to that employed in the proof of Theorem 1 can be used to verify the following generalization of Theorem 1.

THEOREM 2. *Let n_1, n_2, \dots, n_k ($k \geq 1$) be positive integers, and let $\nu_1, \nu_2, \dots, \nu_k$ be vertex partition parameters for which the corresponding properties ν_i are co-hereditary and $\lim \nu_i = \infty$ for $1 \leq i \leq k$. Then*

$$(\nu_i)_1^k(n_1, n_2, \dots, n_k) = 1 + (\bar{\nu}_i)_1^k \cdot \prod_{i=1}^k (n_i - 1).$$

By setting $\nu_i = \nu$ for $i = 1, 2, \dots, k$ in the statement of Theorem 2, we obtain Theorem 1. We present two specific illustrations of Theorem 2.

COROLLARY 2a. *Let n_1, n_2, \dots, n_k ($k \geq 1$) be positive integers, and let $\nu_1, \nu_2, \dots, \nu_k$ be parameters such that $\nu_i = a$ for $1 \leq i \leq t$, where $1 \leq t \leq k$, and $\nu_i = \chi$ for all other ν_i . Then*

$$(\nu_i)_1^k(n_1, n_2, \dots, n_k) = 1 + 2t \cdot \prod_{i=1}^k (n_i - 1).$$

Proof. In this case, $(\bar{\nu}_i)_1^k$ is the largest integer p such that there exists a factorization $K_p = \bigcup_{i=1}^k G_i$ with $a(G_i) = 1$ for $1 \leq i \leq t$ and $\chi(G_i) = 1$ for all other i such that $i \leq k$. This is clearly equal to $\bar{a}(t)$, which has the value $2t$.

Similarly, since $\bar{\chi}^{(2)}(t) = 2\{t/2\}$ for each positive integer t , we have the following.

COROLLARY 2b. *Let n_1, n_2, \dots, n_k ($k \geq 1$) be positive integers, and let $\nu_1, \nu_2, \dots, \nu_k$ be parameters such that $\nu_i = \chi^{(2)}$ for $1 \leq i \leq t$, where $1 \leq t \leq k$ and $\nu_i = \chi$ for all other ν_i . Then*

$$(\nu_i)_1^k(n_1, n_2, \dots, n_k) = 1 + 2\{t/2\} \cdot \prod_{i=1}^k (n_i - 1).$$

3. Edge partition parameters. Let ϵ denote a graphical property which the graph K_2 possesses. We then define the *edge partition number* $\epsilon(G)$ of a nonempty graph G as the least number of elements E_i in a partition of the edge set $E(G)$ of G such that each induced subgraph E_i has property ϵ . It is clearly equivalent to say that $\epsilon(G)$ is the minimum positive k for which there exists a factorization $G = \bigcup_{i=1}^k G_i$ such that $\epsilon(G_i) = 1$ for $i = 1, 2, \dots, k$. For an empty graph G , we define $\epsilon(G) = 0$. In this section, we shall henceforth assume that ϵ is a co-hereditary property (so that H is a subgraph of G implies that

$$\epsilon(H) \leq \epsilon(G), \text{ and that } \lim \epsilon = \infty \text{ (i.e., } \lim_{n \rightarrow \infty} \epsilon(K_n) = \infty).$$

Our next lemma, presented without proof, is an immediate consequence of the definitions for edge partition number and factorization.

LEMMA 2. *If $G = \bigcup_{i=1}^k G_i$ and ϵ is an edge partition parameter, then*

$$\epsilon(G) \leq \sum_{i=1}^k \epsilon(G_i).$$

Let ϵ be an edge partition parameter, and let n_1, n_2, \dots, n_k ($k \geq 1$) be nonnegative integers. The ϵ -Ramsey number $\epsilon(n_1, n_2, \dots, n_k)$ is the least positive integer p such that for any factorization $K_p = \bigcup_{i=1}^k G_i$, it follows that $\epsilon(G_i) \geq n_i$ for at least one i , $1 \leq i \leq k$. If $n_i = 0$ for some i , $1 \leq i < k$, then clearly $\epsilon(n_1, n_2, \dots, n_k) = 1$; hence, we henceforth assume that $n_i > 0$ for $i = 1, 2, \dots, k$. Using an argument analogous to those used earlier, one can verify that $\epsilon(n_1, n_2, \dots, n_k)$ exists since ϵ is co-hereditary and $\lim \epsilon = \infty$. In this case also, $\epsilon(n_1, n_2, \dots, n_k)$ is symmetric in n_1, n_2, \dots, n_k .

THEOREM 3. *Let n_1, n_2, \dots, n_k ($k \geq 1$) be positive integers, and let ϵ be an edge partition parameter such that the corresponding property ϵ is co-hereditary and $\lim \epsilon = \infty$. Then $\epsilon(n_1, n_2, \dots, n_k) = N$ where*

$$N = 1 + \max \left\{ p \mid \epsilon(K_p) \leq \sum_{i=1}^k (n_i - 1) \right\}.$$

Proof. Since $\lim \epsilon = \infty$ and $\epsilon(K_1) = 0$, N exists and $N \geq 2$. Without loss of generality, we assume that $n_1 \leq n_2 \leq \dots \leq n_k$. The theorem clearly follows if $n_k = 1$. Thus we assume that $n_k \geq 2$.

First, we establish the inequality $\epsilon(n_1, n_2, \dots, n_k) \leq N$. Let $\bigcup_{i=1}^k G_i$ be a factorization of K_N . It follows from Lemma 2 and the definition of N that

$$\sum_{i=1}^k \epsilon(G_i) \geq \epsilon(K_N) > \sum_{i=1}^k (n_i - 1).$$

Thus, $\epsilon(G_i) \geq n_i$ for at least one i , $1 \leq i \leq k$.

In order to show that $\epsilon(n_1, n_2, \dots, n_k) \geq N$, we exhibit a factorization $K_{N-1} = \bigcup_{i=1}^k G_i$, where $\epsilon(G_i) \leq n_i - 1$ for $1 \leq i \leq k$. Let m be the least integer such that $n_m \geq 2$. By the way in which N and m were chosen,

$$\epsilon(K_{N-1}) \leq \sum_{i=1}^k (n_i - 1) = \sum_{i=m}^k (n_i - 1).$$

This implies that there exists a factorization

$$K_{N-1} = \bigcup_{i=m}^k \bigcup_{j=1}^{n_i-1} H_j^i, \text{ where } \epsilon(H_j^i) \leq 1$$

($m \leq i \leq k$ and $1 \leq j \leq n_i - 1$). For $m \leq i \leq k$, let $G_i = \bigcup_{j=1}^{n_i-1} H_j^i$. For $1 \leq i \leq m - 1$ (if such i exists), let $G_i = \bar{K}_{N-1}$, the complement of K_{N-1} . Then $K_{N-1} = \bigcup_{i=1}^k G_i$, where $\epsilon(G_i) \leq n_i - 1$ for $1 \leq i \leq k$.

Using the fact that the edge chromatic number $\chi_1(K_p)$ of a nontrivial complete graph K_p is p if p is odd and $p - 1$ if p is even, we obtain the following corollary.

COROLLARY 3a. *Let n_1, n_2, \dots, n_k be positive integers. If $n_1 = n_2 = \dots = n_k = 1$, then $\chi_1(n_1, n_2, \dots, n_k) = 2$. Otherwise, $\chi_1(n_1, n_2, \dots, n_k) = 2[(L + 1)/2] + 1$ where $L = \sum_{i=1}^k (n_i - 1)$.*

COROLLARY 3b. *Let n_1, n_2, \dots, n_k be positive integers, and let a_1 denote the (edge) arboricity parameter. If $n_1 = n_2 = \dots = n_k = 1$, then $a_1(n_1, n_2, \dots, n_k) = 2$. Otherwise,*

$$a_1(n_1, n_2, \dots, n_k) = 1 + 2 \sum_{i=1}^k (n_i - 1).$$

Proof. The result follows from the fact that for $p \geq 2$, $a_1(K_p) = \{p/2\}$.

Let n_1, n_2, \dots, n_k be nonnegative integers, and let $\epsilon_1, \epsilon_2, \dots, \epsilon_k$ be edge partition parameters where again we assume the corresponding properties are co-hereditary and $\lim \epsilon_i = \infty$ for each i , $1 \leq i \leq k$. Then we may define the $(\epsilon_i)_i^k$ -Ramsey number $(\epsilon_i)_i^k(n_1, n_2, \dots, n_k)$ as the least positive integer p such that for any factorization $K_p = \bigcup_{i=1}^k G_i$, it follows that $\epsilon_i(G_i) \geq n_i$ for at least one i where $1 \leq i \leq k$. Again there is no loss of generality in assuming each $n_i > 0$ since $(\epsilon_i)_i^k(n_1, n_2, \dots, n_k) = 1$ if $n_i = 0$ for some i , $1 \leq i \leq k$. If we let M denote the largest integer p for which there exists a factorization $K_p = \bigcup_{i=1}^k G_i$ such that $\epsilon_i(G_i) \leq n_i - 1$ for $i = 1, 2, \dots, k$, then it follows that M exists and that $(\epsilon_i)_i^k(n_1, n_2, \dots, n_k) = 1 + M$; however, it is not possible to give such a compact expression for $(\epsilon_i)_i^k(n_1, n_2, \dots, n_k)$ as for one edge partition parameter (Theorem 3) or k vertex partition parameters.

As an illustration of the foregoing, we consider $(a_1, \chi_1)(m, n)$, for positive integers $m \geq 2$ and $n \geq 2$, defined as the least integer p such that for any factorization $K_p = G_1 \cup G_2$, either $a_1(G_1) \geq m$ or $\chi_1(G_2) \geq n$.

First we show that for every two such positive integers, we have $(a_1, \chi_1)(m, n) \leq 2m + n - 2$. If this is not the case, then there exists a factorization $K_{2m+n-2} = G_1 \cup G_2$ such that $a_1(G_1) \leq m - 1$ and $\chi_1(G_2) \leq$

$n - 1$. This implies that G_1 has at most $(m - 1)(2m + n - 3)$ edges while G_2 has at most $(n - 1) \cdot [(2m + n - 2)/2]$ edges. However, this implies that K_{2m+n-2} has less than $(2m + n - 2)(2m + n - 3)/2$ edges, thereby producing a contradiction.

Next, we note for every positive integer $m \geq 2$ and every odd positive integer $n \geq 3$, that $(a_1, \chi_1)(m, n) = 2m + n - 2$. Here, it suffices to produce a factorization $K_{2m+n-3} = F_1 \cup F_2$ for which $a_1(F_1) \leq m - 1$ and $\chi_1(F_2) \leq n - 1$. Since $2m + n - 3$ is even, there exists a factorization $K_{2m+n-3} = \bigcup_{i=1}^k P_i$, where $k = (2m + n - 3)/2$ and P_i is a spanning path, $i = 1, 2, \dots, k$ (see [3, p. 91]). Let $F_1 = \bigcup_{i=1}^{m-1} P_i$. For $m \leq i \leq k$, we can write $P_i = P_{i,1} \cup P_{i,2}$ where no two edges of $P_{i,j}$, $j = 1, 2$, have adjacent edges. If we let $F_2 = \bigcup_{i=m}^k [P_{i,1} \cup P_{i,2}]$, then we see that $\chi_1(F_2) \leq n - 1$. Since $a_1(F_1) \leq m - 1$, we have a suitable factorization.

Based on the previous observations, we offer the following conjecture.

CONJECTURE. For every two positive integers $m \geq 2$ and $n \geq 2$,

$$(a_1, \chi_1)(m, n) = 2m + n - 2.$$

4. Vertex and edge partition parameters. Let $k = k_1 + k_2$, where k_i , $i = 1, 2$, is a positive integer. Denote by $\rho_1, \rho_2, \dots, \rho_{k_1}$ vertex partition parameters, and denote by $\rho_{k_1+1}, \rho_{k_1+2}, \dots, \rho_k$ edge partition parameters, for which the corresponding properties ρ_i ($1 \leq i \leq k$) are co-hereditary and $\lim \rho_i = \infty$. For positive integers n_1, n_2, \dots, n_{k_1} and nonnegative integers $n_{k_1+1}, n_{k_1+2}, \dots, n_k$, we define the $(\rho_i)_i^k$ -Ramsey number $(\rho_i)_i^k(n_1, n_2, \dots, n_k)$ as the least positive integer ρ such that for any factorization $K_\rho = \bigcup_{i=1}^k G_i$, $\rho_i(G_i) \geq n_i$ for at least one i , $1 \leq i \leq k$. Here we also have that if i_1, i_2, \dots, i_k is a permutation of $1, 2, \dots, k$, then

$$(\rho_{i_j})_{j=1}^k(n_{i_1}, n_{i_2}, \dots, n_{i_k}) = (\rho_i)_i^k(n_1, n_2, \dots, n_k).$$

Let $\overline{(\rho_i(n_i))_i^k}$ denote the largest positive integer p for which there exists a factorization $K_p = \bigcup_{i=1}^k G_i$, where $\rho_i(G_i) \leq n_i$ for $i = 1, 2, \dots, k$. An argument similar to that used in the proof of Lemma 1 guarantees the existence of $\overline{(\rho_i(n_i))_i^k}$ and a straightforward extension of the proof of Theorem 1 can be used to demonstrate the following result.

THEOREM 4. Let k_1 and k_2 be positive integers, where $k = k_1 + k_2$. For $i = 1, 2, \dots, k_1$, let ρ_i be a vertex partition parameter and for $i = k_1 + 1, k_1 + 2, \dots, k$, let ρ_i be an edge partition parameter such that the corresponding properties ρ_i , $1 \leq i \leq k$, are co-hereditary and $\lim \rho_i = \infty$, $1 \leq i \leq k$. Then for positive integers n_1, n_2, \dots, n_k ,

$$(\rho_i)_1^k(n_1, n_2, \dots, n_k) = 1 + \overline{(\rho_i(m_i))}_1^k \cdot \prod_{i=1}^{k_1} (n_i - 1),$$

where $m_i = 1$ for $1 \leq i \leq k_1$, and $m_i = n_i = 1$ for $k_1 + 1 \leq i \leq k$.

As an illustration of Theorem 4, we present the following corollary.

COROLLARY 4a. *Let $k = k_1 + k_2$, where k_1 and k_2 are positive integers, and let n_1, n_2, \dots, n_k be positive integers. If $\rho_i = \chi$ for $1 \leq i \leq k_1$ and $\rho_i = \chi_1$ for $k_1 + 1 \leq i \leq k$, then*

$$(\rho_i)_1^k(n_1, n_2, \dots, n_k) = 1 + [\chi_1(n_{k_1+1}, n_{k_1+2}, \dots, n_k) - 1] \cdot \prod_{i=1}^{k_1} (n_i - 1).$$

Proof. By Theorem 4, it suffices to evaluate $\overline{(\rho_i(m_i))}_1^k$, where $m_i = 1$ for $1 \leq i \leq k_1$ and $m_i = n_i - 1$ for $k_1 + 1 \leq i \leq k$. However, since $\rho_i = \chi$ for $1 \leq i \leq k_1$, it suffices to consider $\overline{(\rho_i(m_i))}_{k_1+1}^k$, which equals $\chi_1(n_{k_1+1}, n_{k_1+2}, \dots, n_k) - 1$.

In a similar manner, we obtain the following result concerning chromatic number χ and edge arboricity a_1 .

COROLLARY 4b. *Let $k = k_1 + k_2$, where k_1 and k_2 are positive integers, and let n_1, n_2, \dots, n_k be positive integers. If $\rho_i = \chi$ for $1 \leq i \leq k_1$ and $\rho_i = a_1$ for $k_1 + 1 \leq i \leq k$, then*

$$(\rho_i)_1^k(n_1, n_2, \dots, n_k) = 1 + [a_1(n_{k_1+1}, n_{k_1+2}, \dots, n_k) - 1] \cdot \prod_{i=1}^{k_1} (n_i - 1).$$

REFERENCES

1. G. Chartrand, D. Geller, and S. Hedetniemi, *A generalization of the chromatic number*, Proc. Comb. Phil. Soc., **64** (1968), 265-271.
2. G. Chartrand and A. D. Polimeni, *Ramsey theory and chromatic numbers*, Pacific J. Math., (to appear).
3. F. Harary, *Graph Theory*, Addison-Wesley, Reading, Mass. (1969).

Received June 9, 1975 and in revised form July 23, 1976.

LOUISIANA STATE UNIVERSITY
AND
UNIVERSITY OF LOUISVILLE
LOUISVILLE, KY 40208

Pacific Journal of Mathematics

Vol. 68, No. 1

March, 1977

Richard Julian Bagby, <i>On L^p, L^q multipliers of Fourier transforms</i>	1
Robert Beauwens and Jean-Jacques Van Binnebeek, <i>Convergence theorems in Banach algebras</i>	13
James Cyril Becker, <i>Skew linear vector fields on spheres in the stable range</i>	25
Michael James Beeson, <i>Continuity and comprehension in intuitionistic formal systems</i>	29
James K. Deveney, <i>Generalized primitive elements for transcendental field extensions</i>	41
Samuel S. Feder, Samuel Carlos Gitler and K. Y. Lam, <i>Composition properties of projective homotopy classes</i>	47
Nathan Jacob Fine, <i>Tensor products of function rings under composition</i>	63
Benno Fuchssteiner, <i>Iterations and fixpoints</i>	73
Wolfgang H. Heil, <i>On punctured balls in manifolds</i>	81
Shigeru Itoh, <i>A random fixed point theorem for a multivalued contraction mapping</i>	85
Nicolas P. Jewell, <i>Continuity of module and higher derivations</i>	91
Roger Dale Konyndyk, <i>Residually central wreath products</i>	99
Linda M. Lesniak and John A. Roberts, <i>On Ramsey theory and graphical parameters</i>	105
Vo Thanh Liem, <i>Some cellular subsets of the spheres</i>	115
Dieter Lutz, <i>A perturbation theorem for spectral operators</i>	127
P. H. Maserick, <i>Moments of measures on convex bodies</i>	135
Stephen Joseph McAdam, <i>Unmixed 2-dimensional local domains</i>	153
D. B. McAlister and Norman R. Reilly, <i>E-unitary covers for inverse semigroups</i>	161
William H. Meeks, III and Julie Patrusky, <i>Representing codimension-one homology classes by embedded submanifolds</i>	175
Premalata Mohapatro, <i>Generalised quasi-Nörlund summability</i>	177
Takahiko Nakazi, <i>Superalgebras of weak-*Dirichlet algebras</i>	197
Catherine Louise Olsen, <i>Norms of compact perturbations of operators</i>	209
William Henry Ruckle, <i>Absolutely divergent series and isomorphism of subspaces. II</i>	229
Bernard Russo, <i>On the Hausdorff-Young theorem for integral operators</i>	241
Arthur Argyle Sagle and J. R. Schumi, <i>Anti-commutative algebras and homogeneous spaces with multiplications</i>	255
Robert Evert Stong, <i>Stiefel-Whitney classes of manifolds</i>	271
D. Suryanarayana, <i>On a theorem of Apostol concerning Möbius functions of order k</i>	277
Yoshio Tanaka, <i>On closedness of C- and C^*-embeddings</i>	283
Chi Song Wong, <i>Characterizations of certain maps of contractive type</i>	293