

# Pacific Journal of Mathematics

***E*-UNITARY COVERS FOR INVERSE SEMIGROUPS**

D. B. MCALISTER AND NORMAN R. REILLY

## E-UNITARY COVERS FOR INVERSE SEMIGROUPS

D. B. McALISTER AND N. R. REILLY

An inverse semigroup is called *E-unitary* if the equations  $ea = e = e^2$  together imply  $a^2 = a$ . In a previous paper, the first author showed that every inverse semigroup has an *E-unitary* cover. That is, if  $S$  is an inverse semigroup, there is an *E-unitary* inverse semigroup  $P$  and an idempotent separating homomorphism of  $P$  onto  $S$ . The purpose of this paper is to consider the problem of constructing *E-unitary* covers for  $S$ .

Let  $S$  be an inverse semigroup and let  $F$  be an inverse semigroup, with group of units  $G$ , containing  $S$  as an inverse subsemigroup and suppose that, for each  $s \in S$ , there exists  $g \in G$  such that  $s \leq g$ . Then  $\{(s, g) \in S \times G : s \leq g\}$  is an *E-unitary* cover of  $S$ . The main result of §1 shows that every *E-unitary* cover of  $S$  can be obtained in this way. It follows from this that the problem of finding *E-unitary* covers for  $S$  can be reduced to an embedding problem. A further corollary to this result is the fact that, if  $P$  is an *E-unitary* cover of  $S$  and  $P$  has maximal group homomorphic image  $G$ , then  $P$  is a subdirect product of  $S$  and  $G$  and so can be described in terms of  $S$  and  $G$  alone. The remainder of this paper is concerned with giving such a description.

**1. *E-unitary* covers.** An inverse semigroup is called *E-unitary* if the equations  $ea = e = e^2$  together imply  $a^2 = a$ . It was shown in [4] that every inverse semigroup  $S$  has an *E-unitary* cover in the sense that there is an *E-unitary* inverse semigroup  $P$  together with an idempotent separating homomorphism  $\theta$  of  $P$  onto  $S$ . It was further shown in [5] that every *E-unitary* inverse semigroup is isomorphic to a  $P(G, \mathcal{X}, \mathcal{Y})$  where  $\mathcal{X}$  is a down directed partially ordered set with  $\mathcal{Y}$  an ideal and subsemilattice of  $\mathcal{X}$  and where  $G$  acts on  $\mathcal{X}$  by order automorphisms in such a way that  $\mathcal{X} = G\mathcal{Y}$ ; see [5] for details. The group  $G$  in  $P = P(G, \mathcal{X}, \mathcal{Y})$  is isomorphic to the maximum group homomorphic image  $P/\sigma$  of  $P$  where

$$\sigma = \{(a, b) \in P \times P : ea = eb \text{ for some } e^2 = e \in P\}.$$

**DEFINITION 1.1.** Let  $S$  be an inverse semigroup and let  $G$  be a group. Then an *E-unitary* inverse semigroup  $P$  is an *E-unitary cover of  $S$  through  $G$*  if

- (i)  $P/\sigma \approx G$
- (ii) there is an idempotent separating homomorphism  $\theta$  of  $P$  onto  $S$ .

Thus, if  $P = P(G, \mathcal{X}, \mathcal{Y})$  is an  $E$ -unitary cover of  $S$  then  $P$  is an  $E$ -unitary cover of  $S$  through  $G$ . As stated, the problem of finding  $E$ -unitary covers of an inverse semigroup  $S$  consists of finding homomorphisms onto  $S$ . The main result of this section shows that this problem can be replaced by an embedding problem.

**DEFINITION 1.2.** [2] Let  $S = S^1$  be an inverse semigroup, with group of units  $G$ . Then  $S$  is a *factorizable* inverse semigroup if and only if, for each  $a \in S$  there exists  $g \in G$  such that  $a \leq g$ .

Chen and Hsieh showed in [1] that every inverse semigroup  $S$  can be embedded in a factorizable inverse semigroup. Indeed, let  $\theta$  be a homomorphism of  $S$  into the symmetric inverse semigroup  $\mathcal{I}_X$  on a set  $X$ . Let  $Y = X$  if  $X$  is finite and  $Y = X \cup X'$ , with  $X \cap X' = \emptyset$ ,  $|X| = |X'|$ , otherwise. Then  $F = \{\alpha \in \mathcal{I}_Y : \alpha \leq \gamma \text{ for some permutation } \gamma \text{ of } Y\}$  is a factorizable inverse semigroup which contains  $S\theta$ .

**PROPOSITION 1.3.** *Let  $F$  be a factorizable inverse semigroup with group of units  $G$  and let  $\theta$  be a one-to-one homomorphism of an inverse semigroup  $S$  into  $F$ . Suppose that for each  $g \in G$  there exists  $s \in S$  such that  $s\theta \leq g$ . Then*

$$P = \{(s, g) \in S \times G : s\theta \leq g\}$$

*is an  $E$ -unitary cover of  $S$  through  $G$ .*

*Proof.* It is clear that  $P$  is an inverse subsemigroup of  $S \times G$  and, because  $F$  is factorizable, that the first projection  $\pi_1: P \rightarrow S$  is an idempotent separating homomorphism of  $P$  onto  $S$ . Likewise, because of the condition on the homomorphism  $\theta: S \rightarrow F$ , the second projection  $\pi_2: P \rightarrow G$  is a homomorphism of  $P$  onto  $G$ .

Now  $(s, g), (t, h) \in P$  with  $(s, g)\pi_2 = (t, h)\pi_2$  implies  $g = h$  thus  $(st^{-1})\theta = s\theta(t\theta)^{-1} \leq gh^{-1} = 1$ . It follows from this that  $st^{-1}$  is idempotent so that  $es = et$ , for some idempotent  $e$  in  $S$ . But then  $(e, 1) \in P$  and  $(e, 1)(s, g) = (e, 1)(t, h)$  so that  $(s, g)\sigma(t, h)$ . This shows that  $\pi_2 \circ \pi_1^{-1} \subseteq \sigma$ . On the other hand, since  $\pi_2$  is a homomorphism onto a group,  $\sigma \subseteq \pi_2 \circ \pi_1^{-1}$ . Hence  $P$  has maximum group homomorphic image  $G$ .

Finally, since  $(s, 1) \in P$  implies  $s\theta \leq 1$ , and so  $s^2 = s$ , if  $(s, g)\pi_2 = 1$  then  $(s, g)$  is idempotent. Hence  $P$  is  $E$ -unitary.

It follows from the remarks before Proposition 1.3 that every inverse semigroup has an  $E$ -unitary cover. The main result of this section shows that every  $E$ -unitary cover of  $S$  through  $G$  is constructed as in Proposition 1.3 for some factorizable inverse semigroup  $F$ , with group of

units  $G$ , containing  $S$  as an inverse subsemigroup. In order to prove this, we need some lemmas.

LEMMA 1.4. *Let  $\theta: P(G, \mathcal{X}, \mathcal{Y}) \rightarrow S$  be an idempotent separating homomorphism of an  $E$ -unitary inverse semigroup  $P(G, \mathcal{X}, \mathcal{Y})$  onto  $S$ . For each  $A \in \mathcal{Y}$  set  $N_A = \{g \in G: g^{-1}A \in \mathcal{Y} \text{ and } (A, g)\theta = (A, 1)\theta\}$ ; if  $X = gA$  with  $g \in G, A \in \mathcal{Y}$ , set  $N_X = gN_Ag^{-1}$ . Then  $N_X$  is well defined. Further, the relation  $\pi$  on  $\mathcal{X} \times G$  defined by*

$$(A, g)\pi(B, h) \text{ if and only if } A = B \text{ and } gh^{-1} \in N_A$$

*is an equivalence on  $\mathcal{X} \times G$  inducing  $\theta \circ \theta^{-1}$  on  $P(G, \mathcal{X}, \mathcal{Y})$ .*

*Proof.* It was shown in [5] that the subgroups  $N_A, A \in \mathcal{Y}$  satisfy the following three conditions

- (i)  $N_A \triangleleft C_A = \{g \in G: gB = B \text{ for all } B \leq A\}$  for  $A \in \mathcal{Y}$ ;
- (ii)  $A, gA \in \mathcal{Y}$  implies  $N_{gA} = gN_Ag^{-1}$ ;
- (iii)  $A \leq B \in \mathcal{Y}$  implies  $N_B \subseteq N_A$ .

We use (ii), to show that  $N_X$  is well defined. Suppose  $gA = hB, A, B \in \mathcal{Y}$ . Then  $B = h^{-1}gA$  so that,  $N_B = (h^{-1}g)N_A(h^{-1}g)^{-1}$  by (ii). Thus  $hN_Bh^{-1} = gN_Ag^{-1}$ . When the  $N_X, X \in \mathcal{X}$  are defined in this way, it is easy to see that they obey the analogs of (i), (ii), (iii) and the remainder of the lemma follows easily.

For each  $C \in \mathcal{X}, g \in G$ , we shall denote by  $[C, g]$  the  $\pi$ -class containing  $(C, g)$ . Further, we shall denote by  $\mathcal{Z}$  the set  $(\mathcal{X} \times G)/\pi \cup G$ .

LEMMA 1.5. *For each  $s \in S$  such that  $s = (A, g)\theta$ , set*

$$\Delta\rho_s = \{[C, h]: h^{-1}C \leq A\}$$

$$\text{and } [C, h]\rho_s = [C, hg] \text{ for each } [C, h] \in \Delta\rho_s.$$

*Then  $\rho: s \rightarrow \rho_s$  is a faithful representation of  $S$  by one-to-one partial transformations of  $\mathcal{Z}$ .*

*Proof.* This follows straightforwardly, using the fact that  $\{N_X: X \in \mathcal{X}\}$  obeys (i), (ii), (iii) of Lemma 1.4.

For each  $g \in G$ , define  $\alpha_g: \mathcal{Z} \rightarrow \mathcal{Z}$  by

$$h\alpha_g = hg \text{ and } [C, h]\alpha_g = [C, hg]$$

for each  $h \in G, [C, h] \in (\mathcal{X} \times G)/\pi$ . Then, as in Lemma 1.5, it follows that  $\alpha: h \rightarrow \alpha_h$  is a faithful representation of  $G$  by permutations of  $\mathcal{Z}$ . Let  $F = \{\gamma \in \mathcal{I}_{\mathcal{Z}}: \gamma \leq \alpha_g \text{ for some } g \in G\}$ .

PROPOSITION 1.6. *F is a factorizable inverse semigroup which contains  $Sp$ . Further, F has group of units  $\{\alpha_g : g \in G\}$  and for each  $\alpha_g$  there exists  $\rho_s \leq \alpha_g$ ; also*

$$P(G, \mathcal{X}, \mathcal{Y}) \approx \{(s, g) \in S \times G : s \leq \alpha_g\}.$$

*Proof.* The only part requiring verification is that  $P(G, \mathcal{X}, \mathcal{Y}) \approx P$  where  $P = \{(s, g) \in S \times G : s \leq \alpha_g\}$ .

Define  $\phi : P(G, \mathcal{X}, \mathcal{Y}) \rightarrow P$  by  $(A, g)\phi = ((A, g)\theta, \alpha_g)$ . Then, since  $\theta$  is idempotent separating and  $\alpha$  is faithful,  $\phi$  is one-to-one. Further

$$\begin{aligned} (A, g)\phi(B, h)\phi &= ((A, g)\theta, \alpha_g)((B, h)\theta, \alpha_h) \\ &= ((A \wedge gB, gh)\theta, \alpha_g\alpha_h) \\ &= ((A, g)(B, h))\phi \end{aligned}$$

so  $\phi$  is a homomorphism.

Finally, if  $(s, \alpha_g) \in P$ , where  $s = (B, h)\theta$ , then  $\rho_s \leq \alpha_g$  implies  $[C, 1]\rho_s = [C, 1]\alpha_g$  for each  $C \leq B$ ; that is,  $[C, h] = [C, g]$  for each  $C \leq B$ . In particular,  $[B, h] = [B, g]$  so that  $s = (B, h)\theta = (B, g)\theta$ . Hence  $(s, \alpha_g) = (B, g)\phi$  so that  $\phi$  is onto.

Summing up, we have the following theorem.

THEOREM 1.7. *Let G be a group and let S be an inverse semigroup. Let F be a factorizable inverse semigroup with group of units G which contains S as an inverse subsemigroup. Suppose that, for each  $g \in G$ , there exists  $s \in S$  such that  $s \leq g$ . Then*

$$\{(s, g) \in S \times G : s \leq g\}$$

*is an E-unitary cover of S through G. Conversely, each E-unitary cover is isomorphic to a cover obtained in this way.*

COROLLARY 1.8. *Let P be an E-unitary cover of S through G. Then P is a subdirect product of S and G.*

Theorem 1.7 shows that the problem of finding E-unitary covers of S through G is equivalent to finding an embedding of S into a factorizable inverse semigroup. Such embeddings are hard to classify as they may be much larger than S and G. On the other hand, Corollary 1.8 shows that every E-unitary cover of S through G is a subdirect-product of S and G and therefore depends only on S and G. In the next section, we turn to the problem of constructing those subdirect products of S and G which are E-unitary covers of S through G.

## 2. Subdirect products of inverse semigroups.

DEFINITION 2.1. Let  $S$  and  $T$  be inverse semigroups. Then a mapping  $\phi: S \rightarrow 2^T$  is a *subhomomorphism* of  $S$  into  $T$  if

- (i)  $\phi(s) \neq \square$  for each  $s \in S$ ;
- (ii)  $\phi(s)\phi(t) \subseteq \phi(st)$  for all  $s, t \in S$ ;
- (iii)  $\phi(s^{-1}) = \phi(s)^{-1}$  for each  $s \in S$ ,

where, for any  $A \subseteq T$ ,  $A^{-1} = \{a^{-1}: a \in A\}$ .

The set  $\phi(S) = \{t \in T: t \in \phi(s) \text{ for some } s \in S\}$  is, from (ii) and (iii), an inverse subsemigroup of  $T$ . We say that  $\phi$  is *surjective* if  $T = \phi(S)$ .

PROPOSITION 2.2. *Let  $S$  and  $T$  be inverse semigroups and let  $\phi$  be a surjective subhomomorphism of  $S$  into  $T$ . Then*

$$\Pi(S, T, \phi) = \{(s, t) \in S \times T: t \in \phi(s)\}$$

*is an inverse semigroup which is a subdirect product of  $S$  and  $T$ .*

*Conversely, suppose that  $V$  is an inverse semigroup which is a subdirect product of  $S$  and  $T$  and let  $\psi$  be the induced homomorphism of  $V$  into  $S \times T$ . Then  $\phi$  defined by*

$$\phi(s) = \{t \in T: (s, t) \in V\psi\}$$

*is a surjective subhomomorphism of  $S$  into  $T$ . Further*

$$V\psi = \Pi(S, T, \phi).$$

*Proof.* This is straightforward.

Proposition 2.2 shows that every  $E$ -unitary cover of  $S$  through  $G$  is determined by a subhomomorphism of  $S$  into  $G$ ; and, dually, by a subhomomorphism of  $G$  into  $S$ . We shall consider these two approaches and the relationships between them in the later sections of the paper.

**3. Subhomomorphisms into a group.** In this section we shall describe the subhomomorphisms of an inverse semigroup  $S$  into a group  $G$ . Note, however, that not every subhomomorphism  $\phi$  of  $S$  into  $G$  gives an  $E$ -unitary cover of  $S$  through  $G$ .

DEFINITION 3.1. Let  $S$  be an inverse semigroup and let  $G$  be a group. Then a subhomomorphism  $\phi: S \rightarrow G$  is *unitary* if

$$1 \in \phi(s) \text{ implies } s^2 = s.$$

Note that, if  $s$  is an idempotent then

$$\phi(s)\phi(s)^{-1} = \phi(s)\phi(s^{-1}) \subseteq \phi(ss^{-1}) = \phi(s)$$

so that  $1 \in \phi(s)$ .

**PROPOSITION 3.2.** *Let  $S$  be an inverse semigroup and let  $G$  be a group. Suppose that  $\phi$  is a surjective unitary subhomomorphism of  $S$  into  $G$ . Then  $\Pi(S, G, \phi)$  is an  $E$ -unitary cover of  $S$  through  $G$ . Conversely, let  $P$  be an  $E$ -unitary cover of  $S$  through  $G$  with  $\psi$  the induced homomorphism  $P \rightarrow S \times G$ . Then  $\phi$  defined by*

$$\phi(s) = \{g \in G : (s, g) \in P\psi\}$$

*is a surjective unitary subhomomorphism of  $S$  into  $G$ .*

*Proof.* Suppose that  $\phi$  is unitary. Then, since the idempotents of  $P = \Pi(S, G, \phi)$  are the elements  $(e, 1)$  with  $e^2 = e$  in  $S$ , the projection  $\pi_S$  of  $P$  onto  $S$  is idempotent separating.

The projection  $\pi_G: P \rightarrow G$  is onto, so, to prove that  $G \approx P/\sigma$ , we need only show that  $\pi_G \circ \pi_G^{-1} = \sigma$ . In fact, since  $\pi_G$  is a homomorphism onto a group, so that  $\sigma \subseteq \pi_G \circ \pi_G^{-1}$ , we need only show that  $\pi_G \circ \pi_G^{-1} \subseteq \sigma$ .

Suppose that  $(s, g)\pi_G = (t, h)\pi_G$  so that  $g = h$ . Then  $(s, g), (t, g) \in P$  implies  $(st^{-1}, 1) \in P$ . That is,  $1 \in \phi(st^{-1})$ . Since  $\phi$  is unitary, this implies  $st^{-1}$  is idempotent so that  $es = et$  for some idempotent  $e$  in  $S$ . But then  $(e, 1)(s, g) = (e, 1)(t, g) = (e, 1)(t, h)$  so that  $(s, g)\sigma(t, h)$ . Thus  $\pi_G \circ \pi_G^{-1} \subseteq \sigma$ .

Finally, if  $(s, g) \in P$  and  $(e, 1)(s, g) = (e, 1)$  then  $g = 1$  so that  $1 \in \phi(s)$ . Since  $\phi$  is unitary, this requires  $s^2 = s$ ; whence  $(s, g)$  is idempotent. Hence  $P$  is an  $E$ -unitary cover of  $S$  through  $G$ .

Conversely, suppose that  $P$  is an  $E$ -unitary cover of  $S$  through  $G$  and let  $1 \in \phi(s)$ . Then  $(s, 1) = p\psi$  for some  $p \in P$ . But  $1 = p\psi\pi_G = p\sigma^{\sharp}$  implies  $p^2 = p$ , since  $P$  is  $E$ -unitary, so that  $s = p\psi\pi_s = p\theta$ , where  $\theta$  is the idempotent separating homomorphism  $P \rightarrow S$ , is also idempotent. Hence  $\phi$  is unitary.

The proof of Proposition 3.2 is strikingly reminiscent of that of Proposition 1.3. This is because Proposition 1.3 is a special case of Proposition 3.2. For, let  $\theta$  be a homomorphism of  $S$  into a factorizable inverse semigroup  $F$ , with group of units  $G$ , and set

$$\phi(s) = \{g \in G : s\theta \leq g\}.$$

Then  $g \in \phi(s), h \in \phi(t)$  implies  $s\theta \leq g, t\theta \leq h$  so that  $s\theta t\theta \leq gh$ . Thus

$(st)\theta \leq gh$ ; that is,  $gh \in \phi(st)$ . Hence  $\phi(s)\phi(t) \subseteq \phi(st)$ . Further  $s\theta \leq g$  if and only if  $s\theta^{-1} \leq g^{-1}$ ; that is  $s^{-1}\theta \leq g^{-1}$ . Hence  $\phi(s^{-1}) = \phi(s)^{-1}$  so that  $\phi$  is a subhomomorphism. If  $\theta$  is one-to-one, then clearly  $\phi$  is unitary so that Proposition 1.3 follows.

In order to obtain a subhomomorphism  $\phi$ , as above, from a mapping  $\theta: S \rightarrow F$  one need not assume that  $\theta$  is a homomorphism. Only that  $\theta$  is a  $v$ -prehomomorphism in the sense of the following definition.

DEFINITION 3.4. Let  $S, T$  be inverse semigroups then a mapping  $\theta: S \rightarrow T$  is a  $v$ -prehomomorphism if it obeys the following two conditions.

- (i)  $(st)\theta \leq s\theta t\theta$  for each  $s, t \in S$ ;
- (ii)  $(s^{-1})\theta = (s\theta)^{-1}$  for each  $s \in S$ .

If  $S$  and  $T$  are semilattices then a  $v$ -prehomomorphism is just an isotone mapping of  $S$  into  $T$ .

The results in the next lemma follow straightforwardly from the definitions.

LEMMA 3.5. Let  $S$  be an inverse semigroup and let  $F$  be a factorizable inverse semigroup with group of units  $G$ . Suppose that  $\theta$  is a  $v$ -prehomomorphism of  $S$  into  $F$ . Then  $\phi$  defined by

$$\phi(s) = \{g \in G: s\theta \leq g\}$$

is a subhomomorphism of  $S$  into  $G$ . It is surjective if and only if, for each  $g \in G$ , there exists  $s \in S$  such that  $s\theta \leq g$ ; it is unitary if and only if  $\theta$  is idempotent determined in the sense that  $a\theta$  idempotent implies  $a$  idempotent.

Lemma 3.5 shows that  $v$ -prehomomorphisms of  $S$  into  $F$  give rise to subhomomorphisms from  $S$  into  $G$ . On the other hand, Proposition 1.6 shows that surjective unitary subhomomorphisms of  $S$  into  $G$  can be obtained from embeddings of  $S$  into factorizable inverse semigroups with groups of units  $G$ . To end this section, we show that every subhomomorphism of  $S$  into  $G$  is determined by a  $v$ -prehomomorphism  $\theta$  of  $S$  into a factorizable inverse semigroup  $\mathcal{K}(G)$  which depends only on  $G$ .

It follows from this that every subdirect product of  $S$  and  $G$ , in particular every  $E$ -unitary cover of  $S$  through  $G$ , is determined by a  $v$ -prehomomorphism of  $S$  into  $\mathcal{K}(G)$ . The problem of constructing  $v$ -prehomomorphisms between inverse semigroups is considered, in detail, in [6].

Let  $G$  be a group. Then we shall denote by  $\mathcal{K}(G)$  the set of all cosets  $X = Ha$  of  $G$  modulo subgroups of  $G$ . The following simple



lemma characterizes the members of  $\mathcal{K}(G)$  among the nonempty subsets of  $G$ .

LEMMA 3.6 [3]. *Let  $G$  be a group and let  $X$  be a nonempty subset of  $G$ . Then  $X \in \mathcal{K}(G)$  if and only if  $X = XX^{-1}X$ .*

It follows from Lemma 3.6 that any nonempty intersection of cosets is again a coset. We may thus define a binary operation  $*$  on  $\mathcal{K}(G)$  as follows: for  $X, Y \in \mathcal{K}(G)$ ,

$$X * Y = \text{smallest coset that contains } XY.$$

PROPOSITION 3.7 [8]. *Let  $G$  be a group. Then  $\mathcal{K}(G)$  is a factorizable inverse semigroup with group of units isomorphic to  $G$ . The idempotents are the subgroups of  $G$ . Further, for  $X, Y \in \mathcal{K}(G)$ ,  $X \leq Y$  if and only if  $X \supseteq Y$ .*

PROPOSITION 3.8. *Let  $S$  be an inverse semigroup and let  $G$  be a group. Suppose that  $\phi$  is a subhomomorphism of  $S$  into  $G$ . Then  $\theta$  defined by*

$$a\theta = \phi(a) \text{ considered as a member of } \mathcal{K}(G)$$

*is a  $v$ -prehomomorphism of  $S$  into  $\mathcal{K}(G)$ .*

*Proof.* Let  $X = \phi(a)$ ; then  $X \subseteq XX^{-1}X$ . On the other hand, if  $g_1, g_2, g_3 \in X$  then

$$g_1 g_2^{-1} g_3 \in \phi(a)\phi(a)^{-1}\phi(a) = \phi(a)\phi(a^{-1})\phi(a) \subseteq \phi(aa^{-1}a) = \phi(a),$$

since  $\phi$  is a subhomomorphism. Hence  $XX^{-1}X \subseteq X$  and so  $X = XX^{-1}X$ . This shows  $X \in \mathcal{K}(G)$ , so that  $\theta$  is a mapping into  $\mathcal{K}(G)$ .

Next, since  $\phi$  is a subhomomorphism,  $a\theta b\theta \subseteq (ab)\theta$  for each  $a, b \in S$ . But  $a\theta * b\theta$  is the smallest coset containing  $a\theta b\theta$  so this implies  $a\theta * b\theta \subseteq (ab)\theta$ . That is, by Lemma 3.7,  $a\theta * b\theta \geq (ab)\theta$ . Finally, again since  $\phi$  is a subhomomorphism,  $(a^{-1})\theta = \phi(a^{-1}) = \phi(a)^{-1} = (a\theta)^{-1}$  for each  $a \in S$ . Hence  $\theta$  is a  $v$ -prehomomorphism of  $S$  into  $\mathcal{K}(G)$ .

It follows from Proposition 3.8 that the subdirect products of  $S$  and  $G$  are determined by  $v$ -prehomomorphisms of  $S$  into  $\mathcal{K}(G)$ . More precisely, we have the following theorem, which sums up the results of this section. It should be pointed out however that it may be easier to find subhomomorphisms of  $S$  into  $G$  directly than to find  $v$ -prehomomorphisms of  $S$  into  $\mathcal{K}(G)$ .

THEOREM 3.9. *Let  $S$  be an inverse semigroup and let  $G$  be a group.*

(A). Let  $\theta$  be a  $v$ -prehomomorphism of  $S$  into  $\mathcal{K}(G)$  and suppose that, for each  $g \in G$ , there exists  $s \in S$  such that  $s\theta \leq \{g\}$ ; i.e.  $g \in s\theta$ . Then  $\{(s, g): g \in s\theta\}$  is an inverse semigroup which is a subdirect product of  $S$  and  $G$ . Every subdirect product of  $S$  and  $G$  is of this form for some  $v$ -prehomomorphism of  $S$  into  $\mathcal{K}(G)$ .

(B). With  $\theta$  as in (A),  $\{(s, g): g \in s\theta\}$  is an  $E$ -unitary cover of  $S$  through  $G$  if and only if  $\theta$  is idempotent determined. Every  $E$ -unitary cover of  $S$  through  $G$  is of this form for some idempotent determined  $v$ -prehomomorphism of  $S$  into  $\mathcal{K}(G)$ .

**4. Subhomomorphisms from a group.** In §3, we characterized the  $E$ -unitary covers of an inverse semigroup  $S$ , through a group  $G$ , as subdirect products  $\Pi(S, G, \phi)$  with  $\phi$  a subhomomorphism of  $S$  into  $G$ . They can also be described in the form  $\Pi(G, S, \phi)$  with  $\phi$  a subhomomorphism of  $G$  into  $S$ . In this section, we give such a description. As might be expected the results obtained are, in a sense, dual to those in §3.

DEFINITION 4.1. Let  $S$  and  $T$  be inverse semigroups. Then a mapping  $\theta: S \rightarrow T$  is a  $\wedge$ -prehomomorphism of  $S$  into  $T$  if it obeys the following two conditions.

- (i)  $a\theta b\theta \leq (ab)\theta$  for each  $a, b \in S$ ;
- (ii)  $(a^{-1})\theta = (a\theta)^{-1}$  for each  $a \in S$ .

PROPOSITION 4.2. Let  $S$  be an inverse semigroup and let  $G$  be a group. Suppose that  $T$  is an inverse semigroup containing  $S$  and let  $\theta$  be a  $\wedge$ -prehomomorphism of  $G$  into  $T$ . Then  $\phi$  defined by

$$\phi(g) = \{s \in S: s \leq g\theta\}$$

is a subhomomorphism of  $G$  into  $S$ ;  $\phi$  is surjective if and only if, for each  $s \in S$  there exists  $g \in G$  such that  $s \leq g\theta$ .

The semigroup  $\Pi(G, S, \phi)$  is  $E$ -unitary. It is an  $E$ -unitary cover of  $S$  through  $G$  if  $\phi$  is surjective.

*Proof.* The fact that  $\phi$  is a subhomomorphism and the statement about the surjectivity of  $\phi$  are readily verified.

Let  $a = 1\theta$ ; then, since  $\theta$  is a  $\wedge$ -prehomomorphism

$$a = aa^{-1}a = aaa \leq (1.1)\theta a = a^2 \leq a$$

so that  $a$  is idempotent. Let  $(g, s) \in \Pi(G, S, \phi)$  and suppose that  $(g, s)(1, e) = (1, e)$  where  $e$  is idempotent. Then  $g = 1$  so that  $s \leq 1\theta = a$ ; thus  $s$  is idempotent. Hence  $\Pi(G, S, \phi)$  is  $E$ -unitary.

Suppose that  $\phi$  is surjective. Then, if we identify  $S \times G$  with  $G \times S$ ,  $\Pi(G, S, \phi) = \Pi(S, G, \phi^*)$  where  $g \in \phi^*(s)$  if and only if  $s \in \phi(g)$ . Since  $1 \in \phi^*(s)$  implies  $s \leq 1\theta = a$ , which is idempotent,  $\phi^*$  is unitary. Hence, by Proposition 3.2,  $\Pi(G, S, \phi)$  is an  $E$ -unitary cover of  $S$  through  $G$ .

Proposition 4.2 is analogous to Proposition 3.2. The next proposition is similar to Proposition 3.8; it shows that every  $E$ -unitary cover of  $S$  through  $G$  is determined by a  $\wedge$ -prehomomorphism of  $G$  into a semigroup  $C(S)$  depending only on  $S$ .

DEFINITION 4.3 [9]. Let  $S$  be an inverse semigroup. Then a nonempty subset  $H$  of  $S$  is called *permissible* if

- (i)  $a \in H$ ,  $b \leq a$  implies  $b \in H$ ;
- (ii)  $a, b \in H$  implies  $ab^{-1}$ ,  $a^{-1}b$  idempotent.

Schein [9] shows that the set  $C(S)$  of permissible subsets of  $S$  forms an inverse semigroup under subset multiplication. Further  $S$  can be embedded in  $C(S)$  by means of the homomorphism  $\eta$  given by

$$a\eta = \{x \in S : x \leq a\}$$

for each  $a \in S$ .

PROPOSITION 4.4. Let  $S$  be an inverse semigroup and let  $G$  be a group. Suppose that  $\phi$  is a surjective subhomomorphism of  $G$  into  $S$  such that  $\Pi(G, S, \phi)$  is an  $E$ -unitary cover of  $S$  through  $G$ . Then  $\phi(g)$  is permissible for each  $g \in G$  and  $\theta$  defined by

$$g\theta = \phi(g) \quad \text{considered as a member of } C(S)$$

is a  $\wedge$ -prehomomorphism of  $G$  into  $C(S)$ . Further

$$\Pi(G, S, \phi) = \{(g, s) \in G \times S : s \leq g\theta\};$$

here we identify  $S$  with  $S\eta$ .

*Proof.* Suppose  $a \in \phi(g)$ ,  $b \leq a$ ; thus  $b = ea$  for some  $e^2 = e \in S$ . Then  $(g, a) \in P = \Pi(G, S, \phi)$  and  $(1, e) \in P$  so that  $(1, e)(g, a) = (g, b) \in P$ . Hence  $b \in \phi(g)$ . Next, suppose  $a, c \in \phi(g)$  then  $(g, a), (g, c) \in P$  so that  $(1, a^{-1}c) \in P$ . Since  $P$  is an  $E$ -unitary cover of  $S$  through  $G$ , with  $\pi_G \circ \pi_G^{-1} = \sigma$ , where  $\pi_G$  denotes the projection of  $P$  onto  $G$ , this implies that  $a^{-1}c$  is an idempotent. Similarly  $ac^{-1}$  is an idempotent. Hence  $\phi(g)$  is permissible.

It is now easy to show that  $\theta$  is a  $\wedge$ -prehomomorphism and, because  $X \leq Y$  in  $C(S)$  if and only if  $X \subseteq Y$ , that  $P = \{(g, s) \in G \times S : s \leq g\theta\}$ .

If we combine the results of Propositions 4.2 and 4.4, then we obtain the following dual to Theorem 3.9.

**THEOREM 4.5.** *Let  $S$  be an inverse semigroup and let  $G$  be a group. Let  $\theta$  be a  $\wedge$ -prehomomorphism of  $G$  into  $C(S)$  such that, for each  $s \in S$  there exists  $g \in G$  with  $s \leq g\theta$ . Then*

$$\{(g, s) \in G \times S : s \leq g\theta\}$$

*is an  $E$ -unitary cover of  $S$  through  $G$ . Conversely, each  $E$ -unitary cover of  $S$  through  $G$  has this form for some  $\wedge$ -prehomomorphism  $\theta$  of  $G$  into  $C(S)$ .*

## 5. Examples

5.1. *Free group covers.* Let  $S$  be an inverse semigroup and let  $X$  be a set of generators for  $S$  as an inverse semigroup. Let  $X^{-1}$  be a set in one to one correspondence with, but disjoint from,  $X$ . Then there is a homomorphism  $\theta$  from the free semigroup  $F_{X \cup X^{-1}}$ , on  $X \cup X^{-1}$ , onto  $S$  such that  $x\theta^{-1} = x^{-1}\theta$  for each  $x \in X$ . Similarly, there is a homomorphism  $\psi: F_{X \cup X^{-1}} \rightarrow FG_X$ , the free group on  $X$  such that  $x\psi^{-1} = x^{-1}\psi$  for each  $x \in X$ . Define  $\phi: S \rightarrow 2^{FG_X}$  by

$$w \in \phi(s) \text{ if and only if } w = u\psi \text{ for some } u \in F_{X \cup X^{-1}}$$

with  $u\theta = s$ ; that is  $\phi(s) = s\theta^{-1}\psi$  for some  $s \in S$ .

**PROPOSITION 5.1.**  *$\phi$  is a surjective unitary subhomomorphism of  $S$  into  $FG_X$ .*


*Proof.* It is straightforward to show that  $\phi$  is a surjective subhomomorphism of  $S$  into  $FG_X$ . Suppose that  $1 \in \phi(s)$ . Then there exists  $w \in F_{X \cup X^{-1}}$  such that  $w\theta = s$ ,  $w\psi = 1$ .

Let  $\eta$  be the canonical homomorphism from  $F_{X \cup X^{-1}}$  into the free inverse semigroup  $FI_X$  on  $S$ . Then both  $\theta$  and  $\psi$  can be factored through  $\eta$ . Since  $w\psi = 1$  and  $FI_X$  is  $E$ -unitary [7] it follows that  $w$ , regarded as an element of  $FI_X$ , is idempotent. Hence  $s = w\theta$  is an idempotent of  $S$ . This shows that  $\phi$  is unitary.

As a result of the freeness of  $FG_X$ ,  $\Pi(S, FG_X, \phi)$ , with  $\phi$  as above, has a weak universal property.

**PROPOSITION 5.2.** *Let  $S$  be an inverse semigroup and let  $P = \Pi(S, FG_X, \phi)$  as above with  $\alpha$  the homomorphism  $P \rightarrow S$ . Suppose that  $Q$  is an  $E$ -unitary cover of  $S$  through  $G$  with homomorphism  $\beta: Q \rightarrow S$ . Then there is a homomorphism  $\gamma: P \rightarrow Q$  such that  $\alpha = \gamma\beta$ .*

*Proof.* With the notation above, we have the following diagram of maps



For each  $x \in X$ , choose  $y \in Q$  such that  $y\beta = x\theta$ . Then there is a homomorphism  $\nu: F_{X \cup X^{-1}} \rightarrow Q$  such that  $x\nu\beta = x\theta$  and  $(x\nu)^{-1} = x^{-1}\nu$  for each  $x \in X$ . Then  $\nu\sigma^\#$  is a homomorphism of  $F_{X \cup X^{-1}}$  into  $G$  and can be factored through  $\psi$ . That is, there is a homomorphism  $\delta: FG_X \rightarrow G$  such that  $\nu\sigma^\# = \psi\delta$ .

From the definition of  $P$ ,  $P = \{(w\theta, w\psi): w \in F_{X \cup X^{-1}}\}$ . Define  $\gamma: P \rightarrow Q$  by  $(w\theta, w\psi)\gamma = w\nu$ . Then  $w\theta = u\theta$ ,  $w\psi = u\psi$  implies  $w\psi\delta = u\psi\delta$ , that is  $w\nu\sigma^\# = u\nu\sigma^\#$  and  $w\nu\beta = u\nu\beta$ . Since  $Q$  is an  $E$ -unitary cover of  $S$  through  $G$ , Corollary 1.8 shows that  $u\nu = w\nu$ . Hence  $\gamma$  is well defined; it is clearly a homomorphism. Further, from the definition,

$$(w\theta, w\psi)\gamma\beta = w\nu\beta = w\theta = (w\theta, w\psi)\alpha,$$

for each  $(w\theta, w\psi) \in P$ . Hence  $\alpha = \gamma\beta$ .

5.2. *The Preston-Vagner cover.* Let  $\rho: S \rightarrow \mathcal{I}_S$  be the Preston-Vagner representation of an inverse semigroup  $S$  and let  $Y = S$  if  $S$  is finite, if not  $Y = S \cup S'$  with  $S \cap S' = \emptyset$ ,  $|S| = |S'|$ . Then  $F = \{\alpha \in \mathcal{I}_Y: \alpha \leq \gamma \text{ for some permutation } \gamma \text{ of } Y\}$  is a factorizable inverse semigroup containing  $S\rho$ . It gives rise to the subhomomorphism  $\phi$  where, for each  $s \in S$ ,

$$\begin{aligned}
 \phi(s) &= \{\alpha: \rho_s \leq \alpha, \alpha \text{ a permutation of } Y\} \\
 &= \{\alpha: x\alpha = xs \text{ for each } x \in Sss^{-1}, \alpha \text{ a permutation of } Y\}.
 \end{aligned}$$

This subhomomorphism gives an  $E$ -unitary cover of  $S$  through  $K$  where

$$K = \{\alpha \in S_Y: (xe)\alpha = x(e\alpha) \text{ for all } x \in S \text{ and some } e^2 = e \in S\},$$

where  $S_Y$  denotes the symmetric group on  $Y$ .

5.3. *E-unitary covers of bisimple inverse semigroups.* A construction is given in [6] for the  $v$ -prehomomorphisms  $\theta$  of a bisimple inverse semigroup  $S$  into an inverse semigroup  $T$ . When applied to the semigroup  $\mathcal{K}(G)$  of cosets of a group  $G$ , this construction specializes to give the following construction for the  $E$ -unitary covers of  $S$  through  $G$ .

Let  $H$  be a subgroup of  $G$  and let  $S(H) = \{a \in G : aHa^{-1} \subseteq H\}$ . Then  $S(H)$  is a subsemigroup of  $G$  and  $(G/H, S(H)/H)$  is a partial semigroup under the multiplication  $*$ :

$$X * Y = XY \quad \text{for each } X \in S(H)/H, Y \in G/H.$$

Pick an idempotent  $e \in S$  and set  $R_e = \{a \in S : aa^{-1} = e\}$ ,  $P_e = R_e \cap eSe$  and let  $\theta: R_e \rightarrow G/H$  be a one-to-one mapping such that the following hold

- (i)  $a\theta \in S(H)/H$  if  $a \in P_e$
- (ii)  $a\theta b\theta = (ab)\theta$  for  $a \in P_e, b \in R_e$
- (iii)  $G = \cup \{a\theta^{-1}b\theta : a, b \in R_e\}$ .

Then  $\{(s, g) \in S \times G : g \in a\theta^{-1}b\theta \text{ where } s = a^{-1}b\}$  is an  $E$ -unitary cover of  $S$  through  $G$ . Conversely, each such has this form for some  $\theta: R_e \rightarrow G/H$  as above.

5.4. *E-unitary covers for semilattices of groups.* A construction is given in [6] for the  $v$ -prehomomorphisms  $\theta$  of a semilattice of groups  $S$  into an inverse semigroup  $T$ . When applied to the semigroup  $\mathcal{K}(G)$  of cosets of a group  $G$ , this construction specializes to give the following description of the  $E$ -unitary covers of  $S$  through  $G$ .

Let  $E$  be a semilattice and let  $\theta$  be an anti-isotone mapping of  $E$  into the lattice of subgroups of  $G$ . For each  $e \in E$  set  $G_e = e\theta$  and  $C_e = \{a \in G : aG_f a^{-1} = G_f \text{ for each } f \preceq e\}$ . Then  $G_e$  is a normal subgroup of  $C_e$  and the groups  $K_e = C_e/G_e$  form a semilattice of groups  $SL(E, \theta, \mathcal{K}(G))$  with linking homomorphisms  $\phi_{e,f}: K_e \rightarrow K_f$  given by

$$X\phi_{e,f} = G_f X \quad \text{for each } X \in K_e, e \succeq f.$$

Suppose that  $S$  is a semilattice of groups with semilattice of idempotents  $E$ . Suppose that  $\theta$  is an anti-isotone mapping of  $E$  into the lattice of subgroups of  $G$  and let  $\phi$  be an idempotent determined homomorphism of  $S$  into  $SL(E, \theta, \mathcal{K}(G))$  such that  $G = \cup \{a\phi : a \in S\}$ . Then

$$\{(s, g) \in S \times G : g \in s\phi\}$$

is an  $E$ -unitary cover of  $S$  through  $G$ . Conversely, each such has this form for some  $\theta: E \rightarrow \mathcal{K}(G)$  and  $\phi: S \rightarrow SL(E, \theta, \mathcal{K}(G))$ .

## REFERENCES

1. S. Y. Chen and S. C. Hsieh, *Factorizable inverse semigroups*, Semigroup Forum, **8** (1974), 283–297.
2. A. H. Clifford and G. B. Preston, *Algebraic theory of semigroups*, Vols. 1 and 2. Math. Surveys of the American Math. Soc. (Providence, R. I., Vol. 1 (1961), Vol. 2 (1967)).
3. P. Dubreil, *Contribution à la théorie des demi-groupes*, Mèm. Acad. Sci. Inst. France, **63** (1941), 1–52.
4. D. B. McAlister, *Groups, semilattices and inverse semigroups*, Trans. Amer. Math. Soc., **192** (1974), 227–244.
5. ———, *Groups, semilattices and inverse semigroups, II*, Trans. Amer. Math. Soc., **196** (1974), 351–370.
6. ———,  *$v$ -Prehomomorphisms on inverse semigroups*.
7. H. E. Scheiblich, *Free inverse semigroups*, Proc. Amer. Math. Soc., **38** (1973), 1–7.
8. B. M. Schein, *Semigroups of strong subsets* (Russian), Volžskii Matem. Sbornik, Kuibyšev, **4** (1966), 180–186.
9. ———, *Completions, translational hulls and ideal extensions of inverse semigroups*, Czechoslovak Math. J., **23** (1973), 575–609.

Received May 24, 1976. This research was partly supported by a grant from the National Science Foundation and a grant from the National Research Council.

NORTHERN ILLINOIS UNIVERSITY  
DEKALB, IL 60115

AND

SIMON FRASER UNIVERSITY  
BURNABY, BRITISH COLUMBIA, CANADA





Richard Julian Bagby, <i>On <math>L^p</math>, <math>L^q</math> multipliers of Fourier transforms</i> .....	1
Robert Beauwens and Jean-Jacques Van Binnebeek, <i>Convergence theorems in Banach algebras</i> .....	13
James Cyril Becker, <i>Skew linear vector fields on spheres in the stable range</i> .....	25
Michael James Beeson, <i>Continuity and comprehension in intuitionistic formal systems</i> .....	29
James K. Deveney, <i>Generalized primitive elements for transcendental field extensions</i> .....	41
Samuel S. Feder, Samuel Carlos Gitler and K. Y. Lam, <i>Composition properties of projective homotopy classes</i> .....	47
Nathan Jacob Fine, <i>Tensor products of function rings under composition</i> .....	63
Benno Fuchssteiner, <i>Iterations and fixpoints</i> .....	73
Wolfgang H. Heil, <i>On punctured balls in manifolds</i> .....	81
Shigeru Itoh, <i>A random fixed point theorem for a multivalued contraction mapping</i> .....	85
Nicolas P. Jewell, <i>Continuity of module and higher derivations</i> .....	91
Roger Dale Konyndyk, <i>Residually central wreath products</i> .....	99
Linda M. Lesniak and John A. Roberts, <i>On Ramsey theory and graphical parameters</i> .....	105
Vo Thanh Liem, <i>Some cellular subsets of the spheres</i> .....	115
Dieter Lutz, <i>A perturbation theorem for spectral operators</i> .....	127
P. H. Maserick, <i>Moments of measures on convex bodies</i> .....	135
Stephen Joseph McAdam, <i>Unmixed 2-dimensional local domains</i> .....	153
D. B. McAlister and Norman R. Reilly, <i>E-unitary covers for inverse semigroups</i> .....	161
William H. Meeks, III and Julie Patrusky, <i>Representing codimension-one homology classes by embedded submanifolds</i> .....	175
Premalata Mohapatro, <i>Generalised quasi-Nörlund summability</i> .....	177
Takahiko Nakazi, <i>Superalgebras of weak-<math>*</math>Dirichlet algebras</i> .....	197
Catherine Louise Olsen, <i>Norms of compact perturbations of operators</i> .....	209
William Henry Ruckle, <i>Absolutely divergent series and isomorphism of subspaces. II</i> .....	229
Bernard Russo, <i>On the Hausdorff-Young theorem for integral operators</i> .....	241
Arthur Argyle Sagle and J. R. Schumi, <i>Anti-commutative algebras and homogeneous spaces with multiplications</i> .....	255
Robert Evert Stong, <i>Stiefel-Whitney classes of manifolds</i> .....	271
D. Suryanarayana, <i>On a theorem of Apostol concerning Möbius functions of order <math>k</math></i> .....	277
Yoshio Tanaka, <i>On closedness of <math>C</math>- and <math>C^*</math>-embeddings</i> .....	283
Chi Song Wong, <i>Characterizations of certain maps of contractive type</i> .....	293