

Pacific Journal of Mathematics

**REPRESENTING CODIMENSION-ONE HOMOLOGY CLASSES
BY EMBEDDED SUBMANIFOLDS**

WILLIAM H. MEEKS, III AND JULIE PATRUSKY

REPRESENTING CODIMENSION-ONE HOMOLOGY CLASSES BY EMBEDDED SUBMANIFOLDS

WILLIAM H. MEEKS III AND JULIE PATRUSKY

In this paper we prove a general theorem on representing codimension-one homology classes on compact manifolds. Our theorem gives an elementary proof of the classical result that a one-dimensional homology class on a compact orientable surface can be represented by an embedded circle precisely when the class is primitive. We will call a homology class primitive if it is the zero class or if it is not a nontrivial multiple of another class.

Our representation theorem is motivated by our earlier work [2] on the classical situation. For compact orientable surfaces we developed a simple algorithm for representing primitive homology classes by embedded circles. This algorithm is also useful for proving other results related to two and three dimensional topology which are not implied by the general result here. Recently Mark Meyerson [3] has given another proof of the two dimensional case by applying Lickorish "twist" homeomorphisms to a fixed homology class.

The following theorem deals with the decomposition of a n -dimensional manifold M^n by an embedded submanifold N^{n-1} which represents a homology class $\delta \in H_{n-1}(M, Z)$. We will say that such a representation N is minimal if there is no other representation N' of δ having fewer path components.

REPRESENTATION THEOREM. *Suppose M is a compact orientable piecewise linear n -dimensional manifold and $\gamma \in H_{n-1}(M, Z)$ is a primitive nonzero class.*

1. *The class $\delta = \kappa\gamma$ has a minimal representation by a submanifold.*
2. *If N is a minimal representation for $\delta = \kappa\gamma$ then each path component of $M - N$ has two ends and the number of path components of $N = |N|$ is κ .*

Proof. The Poincaré dual to $\delta = \kappa\gamma$ can be represented by a piecewise linear mapping $P(\delta): M \rightarrow S^1$. The standard duality theorems imply that wherever $r \in S^1$ is a regular value for $P(\delta)$, then $P^{-1}(r) \subset M$ is an embedded submanifold representing δ . Hence any $\delta \in H_{n-1}(M, Z)$ has a minimal representative.

Suppose now that N is an oriented path-connected submanifold of M with $[N] \neq 0 \in H_{n-1}(M, Z)$. Since N is path connected and nontrivial

on homology then $M - N$ is path connected. Hence there is an embedded circle $\sigma \subset M$ with $[\sigma] \cap [N] = +1$, where \cap is the intersection pairing on homology. This implies that $[N]$ is primitive.

DEFINITION. If $N \subset M$ is an embedded submanifold, and U is a path component of $M - N$, then $T = \text{end closure}$ of U is formed by placing a compact boundary on each end of U .

Let N be an embedded submanifold representing $\delta \in H_{n-1}(M, Z)$. Suppose T is the end closure of an oriented path component of $M - N$ with at least three ends, E_1, E_2, E_3 , coming from cuts along distinct path components N_1, N_2, N_3 of N . Orient the end E_i as the respective component N_i is oriented. Let γ_1 be a path joining $p_1 \in E_1$ to $p_2 \in E_2$, and let γ_2 be a path joining $p_2 \in E_2$ to $p_3 \in E_3$ with $\gamma_1 \cap \gamma_2 = \{p_2\}$. If $\text{sgn}(\gamma_1 \cap E_1) \neq \text{sgn}(\gamma_1 \cap E_2)$, then we can take the connected sum of N_1 and N_2 in M along γ_1 , and similarly, if $\text{sgn}(\gamma_2 \cap E_2) \neq \text{sgn}(\gamma_2 \cap E_3)$, we can connect N_2 and N_3 in M along γ_2 . If neither of these cases hold, form the composite path $\gamma_3 = \gamma_1 \circ \gamma_2^{-1}$. Clearly, $\text{sgn}(\gamma_3 \cap E_1) \neq \text{sgn}(\gamma_3 \cap E_3)$. Since the normal bundle to E_2 is trivial, we may push γ_3 off of E_2 and take the connected sum of N_1 and N_3 in M along this variation of γ_3 .

A slight change in the above argument shows that the condition that E_1, E_2, E_3 come from cuts along distinct path components of N is not needed. Hence, if N is a minimal representative of δ , then each path component of $M - N$ has two ends. This implies that $[N] = |N| \cdot [N_1]$, where N_1 is a path component of N . Since N_1 is primitive, the theorem is proved.

REFERENCES

1. D. R. J. Chillingworth, *Winding numbers on surfaces*, II., Math. Ann., **199** (1972), 142-153.
2. W. Meeks and J. Patrusky, *Representing homology classes by embedded circles on a compact surface*. To be published in Illinois J. Math.
3. M. D. Meyerson, *Representing homology classes of closed orientable surfaces*. To be published in Proc. A.M.S.
4. B. L. Reinhart, *Algorithms for Jordan curves on compact surfaces*, Ann. Math., **75** (1962), 209-222.

Received December 12, 1974, and in revised form July 23, 1976. This work was supported under NSF Grant No. MCS 76-07147.

UNIVERSITY OF CALIFORNIA, LOS ANGELES
AND
UNIVERSITY OF CALIFORNIA, BERKELEY

Pacific Journal of Mathematics

Vol. 68, No. 1

March, 1977

Richard Julian Bagby, <i>On L^p, L^q multipliers of Fourier transforms</i>	1
Robert Beauwens and Jean-Jacques Van Binnebeek, <i>Convergence theorems in Banach algebras</i>	13
James Cyril Becker, <i>Skew linear vector fields on spheres in the stable range</i>	25
Michael James Beeson, <i>Continuity and comprehension in intuitionistic formal systems</i>	29
James K. Deveney, <i>Generalized primitive elements for transcendental field extensions</i>	41
Samuel S. Feder, Samuel Carlos Gitler and K. Y. Lam, <i>Composition properties of projective homotopy classes</i>	47
Nathan Jacob Fine, <i>Tensor products of function rings under composition</i>	63
Benno Fuchssteiner, <i>Iterations and fixpoints</i>	73
Wolfgang H. Heil, <i>On punctured balls in manifolds</i>	81
Shigeru Itoh, <i>A random fixed point theorem for a multivalued contraction mapping</i>	85
Nicolas P. Jewell, <i>Continuity of module and higher derivations</i>	91
Roger Dale Konyndyk, <i>Residually central wreath products</i>	99
Linda M. Lesniak and John A. Roberts, <i>On Ramsey theory and graphical parameters</i>	105
Vo Thanh Liem, <i>Some cellular subsets of the spheres</i>	115
Dieter Lutz, <i>A perturbation theorem for spectral operators</i>	127
P. H. Maserick, <i>Moments of measures on convex bodies</i>	135
Stephen Joseph McAdam, <i>Unmixed 2-dimensional local domains</i>	153
D. B. McAlister and Norman R. Reilly, <i>E-unitary covers for inverse semigroups</i>	161
William H. Meeks, III and Julie Patrusky, <i>Representing codimension-one homology classes by embedded submanifolds</i>	175
Premalata Mohapatro, <i>Generalised quasi-Nörlund summability</i>	177
Takahiko Nakazi, <i>Superalgebras of weak-*Dirichlet algebras</i>	197
Catherine Louise Olsen, <i>Norms of compact perturbations of operators</i>	209
William Henry Ruckle, <i>Absolutely divergent series and isomorphism of subspaces. II</i>	229
Bernard Russo, <i>On the Hausdorff-Young theorem for integral operators</i>	241
Arthur Argyle Sagle and J. R. Schumi, <i>Anti-commutative algebras and homogeneous spaces with multiplications</i>	255
Robert Evert Stong, <i>Stiefel-Whitney classes of manifolds</i>	271
D. Suryanarayana, <i>On a theorem of Apostol concerning Möbius functions of order k</i>	277
Yoshio Tanaka, <i>On closedness of C- and C^*-embeddings</i>	283
Chi Song Wong, <i>Characterizations of certain maps of contractive type</i>	293