

Pacific Journal of Mathematics

**ON THE HAUSDORFF-YOUNG THEOREM FOR INTEGRAL
OPERATORS**

BERNARD RUSSO

ON THE HAUSDORFF-YOUNG THEOREM FOR INTEGRAL OPERATORS

BERNARD RUSSO

A sharp inequality of Hausdorff-Young type is proved for integral operators. Applications are made in operator theory and in harmonic analysis on locally compact groups.

In recent work the author stated an inequality of Hausdorff-Young type for integral operators which proved to be useful in obtaining L^p estimates on certain locally compact unimodular groups. The present paper is devoted to a closer analysis of that inequality together with some applications to operator theory and to L^p -Fourier analysis on locally compact groups.

In the first place, the proof given previously in [15 I], for the inequality is incomplete, so this paper will begin in §2 with a correct proof of the inequality (Theorem 1). Also shown in §2 is the nonexistence of extremal functions in a particular instance (Prop. 8). In §3 the results of §2 are applied to obtain estimates for the norm of the L^p -Fourier transform on certain unimodular groups. Here some of the machinery from [10] is used in the examples, one class of which (Prop. 13) does not depend on Theorem 1. This has happened before, see [15 I: §3]. For certain members of this class however, it is shown that a better estimate can be obtained using Theorem 1 (Prop. 15). In §4 the study of Hausdorff-Young inequalities on nonunimodular groups is initiated. In view of the recent work on Plancherel formulas for nonunimodular groups such inequalities with constant 1 might be considered routine. It is shown here using Theorem 1, that the natural Hausdorff-Young inequality on the " $ax + b$ " group has a constant less than 1 (Prop. 19). In §5 an operator valued analog of the Fourier transform on Abelian groups is introduced, which is motivated by preceding sections, and it is shown, using Theorem 1, that it behaves in some respects like the Fourier transform (Prop. 20).

2. The Hausdorff-Young Theorem for integral operators. Let X be a σ -finite measure space, $k \in L^2(X \times X)$, and let K be the integral operator with kernel k , i.e.,

$$(1) \quad Kf(x) = \int k(x, y)f(y)dy, \quad f \in L^2(X), \quad x \in X.$$

For $1 \leq p, q < \infty$ the mixed norms (cf. [3]) are defined by

$$(2) \quad \|k\|_{p,q} = \left(\int \left(\int |k(x, y)|^p dx \right)^{q/p} dy \right)^{1/q},$$

with the obvious meaning if p or q is ∞ , and k^* is defined by $k^*(x, y) = \overline{k(y, x)}$. For any bounded linear operator T on $L^2(X)$

$$(3) \quad \|T\|_r = \{\text{tr}(T^*T)^{r/2}\}^{1/r}, \quad 1 \leq r < \infty,$$

and $\|T\|_\infty =$ the operator bound of T . Then c_p denotes the set of operators T with $\|T\|_p < \infty$.

THEOREM 1. *Let $1 < p < 2$, $p' = p/(p-1)$ and let $k \in L^2(X \times X)$ where X is a σ -finite measure space. If K is the integral operator with kernel k , then*

$$(4) \quad \|K\|_{p'} \leq (\|k\|_{p,p} \|k^*\|_{p,p'})^{1/2}.$$

The proof will be preceded by five lemmas, the first three of which are proved by standard measure theoretic techniques.

LEMMA 2. *If $p, q \in [1, \infty)$ and $\|k\|_{p,q} < \infty$, then there exist simple functions $\{s_n\}$ on $X \times X$ (independent of p and q) such that $\|k - s_n\|_{p,q} \rightarrow 0$ and $\|k^* - s_n^*\|_{p,q} \rightarrow 0$.*

LEMMA 3. *Let E be a measurable subset of $X \times X$ such that $\|\chi_E\|_{p,q} < \infty$ for some $p, q \in [1, \infty)$. Then*

$$\lim_{\substack{|F| \rightarrow 0 \\ F \in \mathcal{C}_E}} \|\chi_F\|_{p,q} = 0$$

where $|F|$ is the measure of F and χ_F is the characteristic function of F .

LEMMA 4. *For each measurable subset E of $X \times X$ of finite measure and each $\delta > 0$, there exists a set F which is a finite disjoint union of measurable rectangles such that $|E \Delta F| < \delta$.*

For the convenience of the reader it is remarked that the sequence $\{s_n\}$ in Lemma 2 is just the usual one [7: (11.35)], Lemma 3 is a consequence of dominated convergence, and Lemma 4 can be reduced to the case of a finite measure space whence the proof follows by showing that the collection of sets with the desired property is a monotone class.

LEMMA 5. Let $Z = X \times X$ and define $\varphi \in L^2(Z \times Z)$ by $\varphi(z, w) = k(z_1, w_1)k^*(z_2, w_2)$ for $z = (z_1, z_2), w = (w_1, w_2) \in Z$. Let Φ be the integral operator on $L^2(Z)$ with kernel φ . Then

- (1) $\|\varphi\|_{p, q} = \|\varphi^*\|_{p, q} = \|k\|_{p, q} \|k^*\|_{p, q}$ for $p, q \in [1, \infty]$.
- (2) $\|\Phi\|_r = \|K\|_r^2$ for $1 \leq r \leq \infty$.

Proof. (1) is immediate from the definition of the mixed norms and (2) is immediate once it is noticed that $\Phi = K \otimes K^*$.

REMARK 6. If $K = \sum_i \lambda_i \xi_i \otimes \bar{\Psi}_i, \lambda_i > 0, \{\xi_i\}, \{\Psi_i\}$ ON sets in $L^2(X)$, is the "spectral" decomposition of K in the sense of [16], then $\Phi = \sum_{(i, j)} \lambda_i \lambda_j (\xi_i \times \Psi_j) \otimes (\bar{\Psi}_i \times \bar{\xi}_j)$ is the "spectral" decomposition of Φ .

The following crude estimate was known to the author in 1972 and appears in [14].

LEMMA 7. Let $k = \sum_{i=1}^m \alpha_i \chi_{A_i \times B_i}$, where $\alpha_i \in \mathbb{C}$ and $\{A_i \times B_i\}_{i=1}^m$ are pairwise disjoint measurable rectangles in $X \times X$. Then for $1 < p < 2, p' = p/(p - 1)$

$$(5) \quad \|K\|_{p'} \leq 2^{1/p'} \max(\|k\|_{p, p'}, \|k^*\|_{p, p'}).$$

The proof of (5) is an adaptation of a standard interpolation argument inspired by [3: §7] and based on the validity of (4) at the endpoints $p = 1, 2$. The crudeness of the result and the necessary modifications are due to the circumstance that although $\|k\|_{1, \infty}$ (resp. $\|k^*\|_{1, \infty}$) and $\|k\|_{2, 2}$ interpolate to $\|k\|_{p, p'}$ (resp. $\|k^*\|_{p, p'}$) it is problematical whether $\max(\|k\|_{1, \infty}, \|k^*\|_{1, \infty})$ and $\|k\|_{2, 2}$ interpolate to $O(\max(\|k\|_{p, p'}, \|k^*\|_{p, p'}))$.

Proof of Lemma 7. Assume that $\|k\|_{p, p'} \leq 1$ and $\|k^*\|_{p, p'} \leq 1$. For $\xi \in \mathbb{C}$ define a function k_ξ by the rule

$$(6) \quad k_\xi(x, y) = |k(x, y)|^{(2-\xi)p/2} \operatorname{sgn} k(x, y) M(x, y)^{-p+\xi(p'+p)/2}$$

when $k(x, y) \neq 0$ and $k_\xi(x, y) = 0$ if $k(x, y) = 0$ where $M(x, y) = \max(\|k(x, \cdot)\|_p, \|k(\cdot, y)\|_p)$. The function k_ξ is supported on a set of finite measure and bounded for ξ in the strip $\Omega: 0 \leq \operatorname{Re} \xi \leq 1$. Thus $k_\xi \in L^2(X \times X)$ for $\xi \in \Omega$ and so defines an integral operator K_ξ which is at least Hilbert Schmidt. Let A be a finite rank operator on $L^2(X)$ with polar decomposition $A = V|A|$ and assume $\|A\|_p \leq 1$. Set

$$A_\xi = V|A|^{(2-\xi)p/2} \quad \text{for} \quad \xi \in \mathbb{C}$$

so that A_ξ is Hilbert Schmidt and $K_\xi A_\xi$ is trace class for $\xi \in \Omega$. Thus $F(\xi) = \text{tr}(K_\xi A_\xi)$ exists for $\xi \in \Omega$ and we have

$$(7) \quad |F(\xi)| \leq \|K_\xi\|_p \|A_\xi\|_p, \quad \xi \in \Omega, \quad 1 \leq p \leq \infty.$$

It is easily verified that F is analytic in the interior of Ω and continuous and bounded on Ω . For t real, $\|A_{it}\|_1 = \|A\|_p \leq 1$; $\|k_{it}\|_{1,\infty} \leq 1$, and $\|(k_{it})^*\|_{1,\infty} \leq 1$. Thus (7) with $p = 1$, $\xi = it$ and the case $p = 1$ of (4) imply $|F(it)| \leq 1$. Again for t real, $\|A_{1+it}\|_2^2 = \|A\|_p^2 \leq 1$, and $\|k_{1+it}\|_2^2 \leq 2$, the latter since

$$\|k_{1+it}\|^2 = |k|^p M^{p'-p}, \quad \|k_{1+it}\|_2^2 = \left(\int \int_S + \int \int_T \right) |k_{1+it}|^2 dx dy$$

where $S = \{(x, y) \in X \times X: \|k(x, \cdot)\|_p \geq \|k(\cdot, y)\|_p\}$ and $T = X \times X - S$, so that

$$\begin{aligned} \|k_{1+it}\|_2^2 &\leq \int \int_S |k(x, y)|^p \|k(x, \cdot)\|_p^{p'-p} dx dy \\ &\quad + \int \int_T |k(x, y)|^p \|k(\cdot, y)\|_p^{p'-p} dx dy \\ &\leq \|k^*\|_{p,p'} + \|k\|_{p,p'} \leq 2. \end{aligned}$$

Finally (7) with $p = 2$, $\xi = 1 + it$ and the case $p = 2$ of (4) imply $|F(1 + it)| \leq \sqrt{2}$. By the three lines theorem $|F(s + it)| \leq (\sqrt{2})^s$ for every real s , $0 \leq s \leq 1$. Putting $s = 2/p'$ and $t = 0$ results in $|\text{tr}(KA)| = |F(2/p')| \leq 2^{1/p'}$ so that $\|K\|_{p'} = \sup\{|\text{tr}(KA)|: \|A\|_p \leq 1, A \text{ finite rank}\} \leq 2^{1/p'}$ and (5) is proved.

Proof of Theorem 1. By an approximation argument using Lemmas 2, 3, 4 (and described in [15-I: p. 301]) (5) holds for arbitrary $k \in L^2(X \times X)$. For a given k define $X_0 = X \times X$, $k_0 = k$ and for $n \geq 1$, $X_n = X_{n-1} \times X_{n-1}$, $k_n(z, w) = k_{n-1}(z_1, w_1) k_{n-1}^*(z_2, w_2)$ for $z = (z_1, z_2)$ and $w = (w_1, w_2) \in X_n$. Let K_n be the integral operator on $L^2(X_n)$ with kernel k_n . By Lemma 5

$$\begin{aligned} \|K_n\|_{p'} &= \|K_{n-1}\|_{p'}^2, & n = 1, 2, \dots \\ \|k_n\|_{p,p'} &= \|k_n^*\|_{p,p'} = \|k_{n-1}\|_{p,p'} \|k_{n-1}^*\|_{p,p'} & n = 1, 2, \dots \end{aligned}$$

and by Lemma 7

$$\|K_n\|_{p'} \leq 2^{1/p'} \max(\|k_n\|_{p,p'}, \|k_n^*\|_{p,p'}) \quad n = 0, 1, 2, \dots$$

It follows that

$$\|K\|_{p'} \leq (2^{1/p'})^{2-n} (\|k\|_{p,p'} \|k^*\|_{p,p'})^{1/2} \quad n = 1, 2, \dots$$

and (4) follows by letting $n \rightarrow \infty$.

For the first application consider kernels of the following sort: let f and g be measurable functions on \mathbf{R}^n and let $k(x, y) = f(x - y)g(y)$ so that the corresponding operator K has the form $K = L_f M_g$, $L_f =$ convolution by f , $M_g =$ multiplication by g . The following inequality, valid for $1 < p < 2$, $1/p + 1/p' = 1$, can be obtained from Theorem 1 or by using a known interpolation theorem [11: Th. 3]:

$$(8) \quad \|L_f M_g\|_{p'} \leq \|f\|_p \|g\|_{p'}.$$

Although it is not known if 1 is the best constant in (8) the following result seems to indicate that it is not.

PROPOSITION 8. *If the two sides of (8) are equal and finite, then $f = 0$ a.e., or $g = 0$ a.e.*

Proof. Note first that $\|k\|_{p,p'} = \|f\|_p \|g\|_{p'}$ and by a change of variable and Minkowski's integral inequality, that

$$(9) \quad \|k^*\|_{p,p'} \leq \left(\int \left(\int |f(y)|^{p'} |g(x - y)|^p dx \right)^{p/p'} dy \right)^{1/p} = \|k\|_{p,p'}.$$

By assumption $\|K\|_{p'} = \|k\|_{p,p'}$ so by Theorem 1 $\|K\|_{p'} \leq \|k^*\|_{p,p'} \leq \|k\|_{p,p'} = \|K\|_{p'}$ and so there is equality in (9), whence [5: p. 148] $|f(y)|^{p'} |g(x - y)|^p = \varphi(x)\Psi(y)$ a.e. (x, y) . This forces $f = 0$ a.e., or $g = 0$ a.e.

REMARK 9. Since it is not known if extremal functions exist in general or even in this example it cannot be asserted at this point that $\|\mathcal{F}_p(\Gamma_3)\| < A_p^2$.

3. Unimodular groups. In this section L^p estimates are obtained for some examples of unimodular groups whose Plancherel formulas were derived in [10].

If G is a locally compact separable group and ω is a multiplier on G , then the set of equivalence classes of irreducible ω -representations of G is denoted by \hat{G}^ω . If the group $G(\bar{\omega})$ has a type I regular representation and if G is unimodular then there is a measure $\mu = \mu_{G,\omega}$ on \hat{G}^ω such that

$$(10) \quad \int_{\hat{G}^\omega} \|\pi_\xi(f)\|_2^2 d\mu(\xi) = \int_G |f(x)|^2 dx, \quad f \in L^1(G) \cap L^2(G),$$

where $\pi_\xi \in \xi$ for μ -almost all ξ . This is the unimodular version of the projective Plancherel theorem of Kleppner and Lipsman [10 I: Th. 7.1]. As an immediate consequence of (10) and [12: Th. 3] one has:

PROPOSITION 10. *Let G be a locally compact separable unimodular group with normalized multiplier ω such that $\widehat{G}(\bar{\omega})$ has a Type I regular representation. Then for $1 < p < 2$ there is a constant $\|\mathcal{F}_p(G, \omega)\| \leq 1$ such that*

$$(11) \quad \left(\int_{\hat{G}^\omega} \|\pi_\xi(f)\|_{p'}^{p'} d\mu(\xi) \right)^{1/p'} \leq \|\mathcal{F}_p(G, \omega)\| \left(\int_G |f(x)|^p dx \right)^{1/p}$$

for $f \in L^p(G)$, where $1/p + 1/p' = 1$.

REMARK 11. It is natural as a generalization of [15] to try to improve the estimate $\|\mathcal{F}_p(G, \omega)\| \leq 1$ as G and ω vary. This question is not pursued here. For the example $G = \mathbf{R}^2$ and $\omega = \omega_\gamma$ presented in [10 I, p. 490] one can easily show, using Theorem 1, that $\|\mathcal{F}_p(\mathbf{R}^2, \omega_\gamma)\| \leq A_p (< 1)$. It is also clear that the exact value of $\|\mathcal{F}_p(\mathbf{R}^2, \omega_\gamma)\|$ can be obtained once $\|\mathcal{O}_p(\mathbf{R})\|$ is known (see §5).

PROPOSITION 12. *Let G be unimodular and of type I and separable and let N be the center of G . Suppose that for all $\gamma \in \hat{N}$, the group (G/N) $(\bar{\omega}_\gamma)$ is of type I. Then*

$$\|\mathcal{F}_p(G)\| \leq \|\mathcal{F}_p(N)\| \quad \text{for} \quad 1 \leq p \leq 2.$$

Proof. For groups G satisfying the hypothesis the Plancherel formula is described in [10 I: Th. 8.1]. That and an interpolation argument as in Prop. 10 imply

$$(12) \quad \|\hat{\varphi}\|_{p'} = \left(\int_{\hat{N}} \int_{\widehat{(G/N)^{\bar{\omega}_\gamma}}} \|\pi_{\gamma, \sigma}(\varphi)\|_{p'}^{p'} d\mu_{G/N, \bar{\omega}_\gamma}(\sigma) d\gamma \right)^{1/p'} \leq \|\varphi\|_p$$

for $\varphi \in L^p(G)$, $1 < p < 2$, $1/p + 1/p' = 1$, $N = \text{Cent } G$. Recall from [10 I] that $\pi_{\gamma, \sigma} = \gamma' \otimes \sigma''$ where γ' is an ω_γ -representation of G which extends $\gamma \in \hat{N}$ and σ'' is the lift of $\sigma \in \widehat{(G/N)^{\bar{\omega}_\gamma}}$ to G and ω_γ is the multiplier on G/N corresponding to γ . It follows that

$$\begin{aligned}
 \pi_{\gamma, \sigma}(\varphi) &= \int_G \varphi(g) \gamma'(g) \sigma''(g) dg = \int_{G/N} \int_N \varphi(ng) \gamma'(ng) \sigma''(ng) dnd\bar{g} \\
 &= \int_{G/N} \int_N \varphi(ng) \overline{\omega_\gamma(n, g)} \gamma(n) \gamma'(g) \sigma(\bar{g}) dn d\bar{g} \\
 &= \sigma(\Psi_\gamma) \quad \text{where} \quad \Psi_\gamma(\bar{g}) = [\varphi(\cdot g) \overline{\omega_\gamma(\cdot, g)} \gamma'(g)]^\wedge(-\gamma).
 \end{aligned}$$

Therefore

$$\begin{aligned}
 \|\hat{\varphi}\|_{p'}^{p'} &= \int_{\hat{N}} \int_{(\widehat{G/N})^{\sigma_\gamma}} \|\sigma(\Psi_\gamma)\|_{p'}^{p'} d\mu_{G/N, \bar{\omega}_\gamma}(\sigma) d\gamma \\
 &\cong \int_{\hat{N}} \|\Psi_\gamma\|_{p'}^{p'} d\gamma = \int_{\hat{N}} \left(\int_{G/N} |\Psi_\gamma(\bar{g})|^p d\bar{g} \right)^{p'/p} d\gamma \\
 &\cong \left(\int_{G/N} \left(\int_{\hat{N}} |\Psi_\gamma(\bar{g})|^{p'} d\gamma \right)^{p/p'} d\bar{g} \right)^{p'/p} \\
 &\cong \left(\int_{G/N} \|\mathcal{F}_p(N)\|_p^p \int_N |\varphi(ng) \overline{\omega_\gamma(n, g)} \gamma'(g)|^p dn d\bar{g} \right)^{p'/p} \\
 &= \|\mathcal{F}_p(N)\|_{p'}^{p'} \left(\int_{G/N} \int_N |\varphi(ng)|^p dn d\bar{g} \right)^{p'/p} = \|\mathcal{F}_p(N)\|_{p'}^{p'} \|\varphi\|_{p'}^{p'}.
 \end{aligned}$$

This result is meaningful only if $\|\mathcal{F}_p(N)\| < 1$, i.e., $N = \text{Cent } G$ has no compact open subgroups, e.g., if G is a connected simply connected real nilpotent group. It can also be applied to certain solvable groups e.g. the oscillator group, and to an example of a semi-direct product given in [10 II: §3 (c)] which gives rise to the Weil representation. The estimates given by Prop. 12 for these examples are still crude. For example, for the Heisenberg group Γ_3 with one dimensional center it is already known that $\|\mathcal{F}_p(\Gamma_3)\| \cong A_p^2 < A_p = \|\mathcal{F}_p(\mathbf{R})\|$. Note that Theorem 1 was not used in the proof of Proposition 12. The next example satisfies the hypothesis of Proposition 12 but has a compact center. However a meaningful estimate can still be obtained by using Theorem 1.

EXAMPLE 13. Let $G_1 = G/D$ where G is the three dimensional Heisenberg group and D is the subgroup

$$D = \left\{ \begin{pmatrix} 1 & 0 & 2\pi n \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}; n \in \mathbf{Z} \right\}.$$

The Plancherel formula is given in [10 I: p. 514]. It reads

$$(13) \quad \int_{G_1} |\varphi(g_1)|^2 dg_1 = \int_{\mathbf{R}^2} |\pi_{0,\sigma}(\varphi)|^2 d\sigma + \sum_{n \neq 0} |n| \|\pi_n(\varphi)\|_2^2$$

for $\varphi \in L^1(G_1) \cap L^2(G_1)$, where for $n \neq 0$ the kernel of the integral operator $\pi_n(\varphi)$ is $k_n(x, u) = \varphi(x - u, \cdot, \cdot) \wedge (-nu, -n)$ (Fourier transform on $\mathbf{R} \times T$) and $\pi_{0,\sigma}$ is just the character on \mathbf{R}^2 given by $\sigma \in \mathbf{R}^2$. Interpolating (13) yields

$$\|\hat{\varphi}\|_{p'}^{p'} = \int_{\mathbf{R}^2} |\pi_{0,\sigma}(\varphi)|^{p'} d\sigma + \sum_{n \neq 0} |n| \|\pi_n(\varphi)\|_{p'}^{p'}.$$

Now

$$\begin{aligned} \sum_{n \neq 0} |n| \|k_n\|_{p',p'}^{p'} &= \sum_{n \neq 0} |n| \int \left(\int |\varphi(x - u, \cdot, \cdot) \wedge (-nu, -n)|^p dx \right)^{p'/p} du \\ &= \sum_{n \neq 0} \int_{\mathbf{R}} \left(\int_{\mathbf{R}} |\varphi(x, \cdot, \cdot) \wedge (-v, -n)|^p dx \right)^{p'/p} dv \\ &\cong \left(\int_{\mathbf{R}} \left(\sum_{n \neq 0} \int_{\mathbf{R}} |\varphi(x, \cdot, \cdot) \wedge (-v, -n)|^p dv \right)^{p'/p} dx \right)^{p'/p} \\ &= \left(\int_{\mathbf{R}} (F(x) - G(x))^{p'/p} dx \right)^{p'/p} = \|F - G\|_{p'/p}. \end{aligned}$$

where $F(x) = \|\varphi(x, \cdot, \cdot) \wedge \cdot\|_{p'}^{p'}$ and $G(x) = \int_{\mathbf{R}} |\varphi(x, \cdot, \cdot) \wedge (-v, 0)|^{p'} dv$. But $F - G \geq 0$ so by [7: (13.9)], $\|F\|_{p'/p'} \geq \|F - G\|_{p'/p'} + \|G\|_{p'/p'}$. Thus

$$\begin{aligned} \sum_{n \neq 0} |n| \|k_n\|_{p',p'}^{p'} &\leq \|F\|_{p'/p'} - \|G\|_{p'/p'} = \left(\int \|\varphi(x, \cdot, \cdot) \wedge \cdot\|_{p'}^p dx \right)^{p'/p} \\ &\quad - \left(\int_{\mathbf{R}} \left(\int_{\mathbf{R}} |\varphi(x, \cdot, \cdot) \wedge (-v, 0)|^p dv \right)^{p'/p} dx \right)^{p'/p}. \end{aligned}$$

Similarly

$$\sum_{n \neq 0} |n| \|k_n^*\|_{p',p'}^{p'} \leq \|F\|_{p'/p'} - \|G\|_{p'/p'}.$$

and therefore

$$\begin{aligned}
 \|\hat{\varphi}\|_p^{p'} &\leq \int_{\mathbf{R}^2} |\pi_{0,\sigma}(\varphi)|^{p'} d\sigma + \sum_{n \neq 0} |n| \left(\|k_n\|_{p,p'}^{p'} \|k_n^*\|_{p,p'}^{p'} \right)^{1/2} \\
 &\leq \int_{\mathbf{R}^2} |\pi_{0,\sigma}(\varphi)|^{p'} d\sigma + \left(\sum |n| \|k_n\|_{p,p'}^{p'} \right)^{1/2} \left(\sum |n| \|k_n^*\|_{p,p'}^{p'} \right)^{1/2} \\
 &\leq \int_{\mathbf{R}^2} |\pi_{0,\sigma}(\varphi)|^{p'} d\sigma + \|F\|_{p/p'} - \|G\|_{p/p'} \\
 &\leq \int_{\mathbf{R}^2} |\pi_{0,\sigma}(\varphi)|^{p'} d\sigma - \left(\int \left(\int |\varphi(x, \cdot, \cdot)^\wedge(-v, 0)|^{p'} dv \right)^{p'/p} dx \right)^{p'/p} \\
 &\quad + \|\mathcal{F}_p(\mathbf{R} \times T)\|^{p'} \|\varphi\|_p^{p'} \\
 &\leq \|\mathcal{F}_p(\mathbf{R} \times T)\|^{p'} \|\varphi\|_p^{p'}
 \end{aligned}$$

since

$$\begin{aligned}
 \int_{\mathbf{R}^2} |\pi_{0,\sigma}(\varphi)|^{p'} d\sigma &= \int \int |\pi_{0,(\omega,v)}(\varphi)|^{p'} d\omega dv \\
 &= \int \int |\hat{\varphi}(\omega, v, 0)|^{p'} d\omega dv \\
 &\leq \int A_p^{p'} \left(\int |\varphi(x, \cdot, \cdot)^\wedge(v, 0)|^p dx \right)^{p'/p} dv \\
 &\leq \left(\int \left(\int |\varphi(x, \cdot, \cdot)^\wedge(v, 0)|^{p'} dv \right)^{p'/p} dx \right)^{p'/p}.
 \end{aligned}$$

We conclude:

PROPOSITION 14. *For the group G_1 of the preceding example*

$$\|\mathcal{F}_p(G_1)\| \leq \|\mathcal{F}_p(\mathbf{R})\|.$$

4. Nonunimodular groups. The result of this section is worked out for the two-dimensional solvable group G known as the “ $ax + b$ ” group. It is clear that a calculation virtually identical to that given below will establish similar results for all of the solvable groups considered in [8: Ch. 4]. Before describing which realization of G will be used it is noted that in this section the Fourier transform on \mathbf{R}^n is defined by

$$\hat{f}(y) = \int_{\mathbf{R}^n} e^{2\pi ixy} f(x) dx$$

and therefore

$$(14) \quad ((-\Delta)^{\beta/2} f)^\wedge(x) = (2\pi |x|)^\beta \hat{f}(x) \quad [17: p. 117]$$

where Δ is the Laplacean.

Let $G = \mathbf{R} \times \mathbf{R}_+^*$ with group operation $(\lambda, s)(\mu, t) = (\lambda + s\mu, st)$ right Haar measure $d_R x = d\lambda ds/s$, and modular function $\delta_G(\lambda, s) = 1/s$. The Plancherel formula for G may be stated as follows [13: Th. 4] (cf. [10 I, 8, 9]): there is an operator M in $L^2(G)$ such that

$$(15) \quad \|\hat{\Psi}\|_2^2 \equiv \|\pi_+(\Psi)\|_2^2 + \|\pi_-(\Psi)\|_2^2 = \int_G |M\Psi(x)|^2 d_R x \equiv \|M\Psi\|_2^2$$

for $\Psi \in C_c^\infty(G)$. This M is determined in Proposition 15 and $\{\pi_+, \pi_-\}$ are the two infinite dimensional members of the unitary dual \hat{G} of G .

PROPOSITION 15. *Let G be the “ $ax + b$ ” group, and let $1 < p \leq 2$, $1/p + 1/p' = 1$. Then*

$$\|\hat{\Psi}\|_{p'} = (\|\pi_+(\Psi)\|_{p'} + \|\pi_-(\Psi)\|_{p'})^{1/p'} \leq \|\mathcal{F}_p(\mathbf{R})\| \|M_p \Psi\|_{p'}, \quad \Psi \in C_c^\infty(G)$$

where $M_p = (2\pi)^{1/p'} (-\Delta)^{-1/2p'} \otimes I$. Equality holds (for all Ψ) for $p = 2$.

Proof. The representation π_\pm can be realized as acting on $L^2(\mathbf{R}_+^*, ds/s)$ by $(\pi_\pm(\mu, t)g)(s) = e^{\pm 2\pi i s \mu} g(st)$, $g \in L^2(\mathbf{R}_+^*, ds/s)$, $(\mu, t) \in G$, $s \in \mathbf{R}_+^*$. Then $\pi_\pm(\Psi)$ is an integral operator with kernel $k_\pm(t, s) = \Psi(\cdot, t)^\wedge(\pm st^{-1})$. Letting Σ denote summation over $\{+, -\}$,

$$\begin{aligned} \|\hat{\Psi}\|_{p'}^{p'} &= \sum \|\pi_\pm(\Psi)\|_{p'}^{p'} \leq \sum (\|k_\pm\|_{p', p'}^{p'} \|k_\pm^*\|_{p', p'}^{p'})^{1/2} \\ &\leq \left(\sum \|k_\pm\|_{p', p'}^{p'} \right)^{1/2} \left(\sum \|k_\pm^*\|_{p', p'}^{p'} \right)^{1/2}. \end{aligned}$$

Now

$$\begin{aligned} \sum \|k_\pm\|_{p', p'}^{p'} &= \sum \int_0^\infty \left(\int_0^\infty |\Psi(\cdot, t)^\wedge(\pm st^{-1})|^{p'} \frac{dt}{t} \right)^{p'/p} \frac{ds}{s} \\ &\leq \left(\int_0^\infty \left(\sum \int_0^\infty |\Psi(\cdot, t)^\wedge(\pm st^{-1})|^{p'} \frac{ds}{s} \right)^{p'/p} \frac{dt}{t} \right)^{p'/p} \\ &= \left(\int_0^\infty \left(\sum \int_0^\infty |\Psi(\cdot, t)^\wedge(\pm s)|^{p'} s^{-1/p'} |^{p'} ds \right)^{p'/p} \frac{dt}{t} \right)^{p'/p} \\ &= \left(\int_0^\infty \left(\int_{-\infty}^\infty |\Psi(\cdot, t)^\wedge(\lambda)|^{p'} |\lambda|^{-1/p'} |^{p'} d\lambda \right)^{p'/p} \frac{dt}{t} \right)^{p'/p} \\ &= \left(\int_0^\infty \left(\int_{-\infty}^\infty |(2\pi)^{1/p'} ((-\Delta)^{-1/2p'} \Psi(\cdot, t))^\wedge(\lambda)|^{p'} d\lambda \right)^{p'/p} \frac{dt}{t} \right)^{p'/p} \end{aligned}$$

$$\begin{aligned} &\leq \left(\int_0^\infty A_p^p \|(2\pi)^{1/p'} (-\Delta)^{-1/2p'} \Psi(\cdot, t)\|_p^p \frac{dt}{t} \right)^{p'/p} \\ &= A_p^{p'} \|M_p \Psi\|_p^{p'}. \end{aligned}$$

The same inequality holds for $\Sigma \|k_\pm^*\|_{p, p'}^{p'}$ and the result follows.

5. Analog of the Fourier transform. Let G be a locally compact Abelian group with dual group \hat{G} . For a measurable function F on $\hat{G} \times G$ consider the kernel k_F , $k_F(\alpha, \beta) = F(\alpha - \beta, \cdot)^\wedge(\beta)$, $\alpha, \beta \in \hat{G}$ and the corresponding integral operator K_F defined by $K_F\varphi(\beta) = \int_{\hat{G}} k_F(\alpha, \beta)\varphi(\alpha)d\alpha$. Using the estimates $\|K_F\|_\infty \leq \|F\|_1$ and $\|K_F\|_2 = \|F\|_2$ and the general theory of interpolation one obtains a mapping $\mathcal{O}_p(G): L^p(\hat{G} \times G) \rightarrow c_p(L^2(\hat{G}))$, where $1 < p < 2$, $1/p + 1/p' = 1$, of norm ≤ 1 : $\|K_F\|_{p'} \leq \|F\|_p$.

This type of kernel appears in the Fourier decomposition of the Heisenberg group, see §2 (cf. [4: §2], [15 II: §2]).

PROPOSITION 16. *Let G be a locally compact Abelian group and let $1 < p < 2$. Then*

- (1) $\|\mathcal{O}_p(G)\| \leq \|\mathcal{F}_p(G)\|$
- (2) $\|\mathcal{O}_p(G)\| = 1$ if and only if G has a compact open subgroup.
- (3) If $\|K_F\|_{p'} = \|\mathcal{F}_p(\mathbf{R})\| \|F\|_p$ and p' is an even integer then $F = 0$ a.e. ($G = \mathbf{R}$).

Proof. (1) Using Minkowski's integral inequality and changing variables one obtains that $\|k_F\|_{p, p'}$ and $\|k_F^*\|_{p, p'}$ are both $\leq \|\mathcal{F}_p(G)\| \|F\|_p$ so (1) follows from Theorem 1. (2) If $\|\mathcal{O}_p(G)\| = 1$ then by (1), $\|\mathcal{F}_p(G)\| = 1$ so G has a compact open subgroup by [6: §43] or [15 I: Th. 2]. Conversely if G has a compact open subgroup H , then setting $F(\alpha, x) = c\chi_H \perp(\alpha)\chi_H(x)$ where c is the measure of H one finds that $\|K_F\|_{p'} = \|F\|_p$. (3) The assumption implies that $\|k_F\|_{p, p'} = \|k_F^*\|_{p, p'} = \|\mathcal{F}_p(\mathbf{R})\| \|F\|_p$ and hence that $F(\alpha, \cdot)$ is extremal in $L^p(\mathbf{R})$ for a.e. α and that $|F(\alpha, \cdot)^\wedge(\beta)|^p = \Phi(\alpha)\Psi(\beta)$. Thus

$$F(\alpha, \beta) = c(\alpha)\exp\{-a(\alpha)\beta^2 + ib(\alpha)\beta\}$$

where $c(\alpha) \in \mathbf{C}$, $a(\alpha) > 0$ and $b(\alpha) \in \mathbf{R}$. One finds that

$$F(\alpha, \cdot)^\wedge(\beta) = \text{const } c(\alpha)/(a(\alpha))^{1/2} \times \exp\{-(b(\alpha) - \beta)^2/4a(\alpha)\}$$

and it follows from the above conditions that $a(\alpha)$ and $b(\alpha)$ do not depend on α . Thus $F(\alpha, \beta)$ is of the form $f(\alpha)g(\beta)$, g Gaussian, so that $K_F = L_f M_{\hat{g}}$ and $\|\mathcal{F}_p(\mathbf{R})\| \|F\|_p = \|\mathcal{F}_p(\mathbf{R})\| \|f\|_p \|g\|_p = \|f\|_p \|\hat{g}\|_{p'} \equiv \|L_f M_{\hat{g}}\|_{p'} = \|K_F\|_{p'}$. By Proposition 8 $f = 0$ or $\hat{g} = 0$ a.e. so $F = 0$ a.e.

REMARK 17. The property (3) in Proposition 16 indicates but does not prove that the numbers $\mathcal{F}_p(G)$ and $\mathcal{O}_p(G)$ are either distinct or both equal to 1.

REMARK 18. The behavior of \mathcal{O}_p on products is not as simple as that of \mathcal{F}_p . In particular the computation of $\|\mathcal{O}_p(\mathbf{R}^n)\|$ is not automatically reducible to that of $\|\mathcal{O}_p(\mathbf{R})\|$ (cf. [2: Lemma 2]).

REFERENCES

1. K. I. Babenko, *An inequality in the theory of Fourier integrals*, Izv. Akad. Nauk. SSSR Ser. Mat., **25** (1961), 531–542. English Transl., Amer. Math. Soc. Transl. (2) **44**, 115–128.
2. William Beckner, *Inequalities in Fourier analysis*, Ann. of Math., **102** (1975), 159–182.
3. A. Benedek and R. Panzone, *The spaces L^p with mixed norm*, Duke Math. J., **28** (1961), 301–324.
4. Jacques Dixmier, *Sur les représentations unitaires des groupes de Lie nilpotents. VI*, Canad. J. Math., **12** (1960), 324–352.
5. G. H. Hardy, J. E. Littlewood, and G. Polya, *Inequalities*, Cambridge Univ. Press, 1964.
6. E. Hewitt and K. A. Ross, *Abstract harmonic analysis. Vol. II*, Springer-Verlag, Berlin and New York, 1970.
7. E. Hewitt and K. Stromberg, *Real and abstract analysis*, Springer-Verlag, Berlin and New York, 1969. MR 43 # 428.
8. Frederick W. Keene, *Square integrable representations of Iwasawa subgroups of a semi-simple Lie group*, Doctoral dissertation, Univ. of Calif., Berkeley, 1974.
9. Idriss Khalil, *Sur l'analyse harmonique du groupe affine de la droite*, Stud. Math., **51** (1974), 139–167.
10. Adam Kleppner and Ronald L. Lipsman, *The Plancherel formula for group extensions*, I. Ann. Sci. École Norm. Sup., (4) **5** (1972), 459–516.; II, same journal **6** (1973) 103–132.
11. Ray A. Kunze, *L_p -Fourier transforms on locally compact unimodular groups*, Trans. Amer. Math. Soc., **89** (1958), 519–540.
12. Ray A. Kunze and Elias M. Stein, *Uniformly bounded representations and harmonic analysis of the 2×2 real unimodular group*, Amer. J. Math., **82** (1960), 1–62.
13. Lajos Pukanzsky, *Unitary representations of solvable Lie groups*, Ann. Sci. École Norm. Sup., (4) **4** (1971), 457–608.
14. Bernard Russo, *Algebra di operatori e gruppi localmente compatti*, mimeographed notes, University of Genoa, 1972.
15. ———, *The norm of the L^p -Fourier transform on unimodular groups*, Trans. Amer. Math. Soc., **192** (1974), 293–305. II, Canad. J. Math., **28** (1976), 1121–1131.

16. Robert Schatten, *Norm ideals of completely continuous operators*, Springer-Verlag, 1960.
17. Elias M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Univ. Press, Princeton, N. J., 1970.

Received January 13, 1976 and in revised form May 5, 1976.

UNIVERSITY OF CALIFORNIA
IRVINE, CA 92717

Richard Julian Bagby, <i>On L^p, L^q multipliers of Fourier transforms</i>	1
Robert Beauwens and Jean-Jacques Van Binnebeek, <i>Convergence theorems in Banach algebras</i>	13
James Cyril Becker, <i>Skew linear vector fields on spheres in the stable range</i>	25
Michael James Beeson, <i>Continuity and comprehension in intuitionistic formal systems</i>	29
James K. Deveney, <i>Generalized primitive elements for transcendental field extensions</i>	41
Samuel S. Feder, Samuel Carlos Gitler and K. Y. Lam, <i>Composition properties of projective homotopy classes</i>	47
Nathan Jacob Fine, <i>Tensor products of function rings under composition</i>	63
Benno Fuchssteiner, <i>Iterations and fixpoints</i>	73
Wolfgang H. Heil, <i>On punctured balls in manifolds</i>	81
Shigeru Itoh, <i>A random fixed point theorem for a multivalued contraction mapping</i>	85
Nicolas P. Jewell, <i>Continuity of module and higher derivations</i>	91
Roger Dale Konyndyk, <i>Residually central wreath products</i>	99
Linda M. Lesniak and John A. Roberts, <i>On Ramsey theory and graphical parameters</i>	105
Vo Thanh Liem, <i>Some cellular subsets of the spheres</i>	115
Dieter Lutz, <i>A perturbation theorem for spectral operators</i>	127
P. H. Maserick, <i>Moments of measures on convex bodies</i>	135
Stephen Joseph McAdam, <i>Unmixed 2-dimensional local domains</i>	153
D. B. McAlister and Norman R. Reilly, <i>E-unitary covers for inverse semigroups</i>	161
William H. Meeks, III and Julie Patrusky, <i>Representing codimension-one homology classes by embedded submanifolds</i>	175
Premalata Mohapatro, <i>Generalised quasi-Nörlund summability</i>	177
Takahiko Nakazi, <i>Superalgebras of weak-*Dirichlet algebras</i>	197
Catherine Louise Olsen, <i>Norms of compact perturbations of operators</i>	209
William Henry Ruckle, <i>Absolutely divergent series and isomorphism of subspaces. II</i>	229
Bernard Russo, <i>On the Hausdorff-Young theorem for integral operators</i>	241
Arthur Argyle Sagle and J. R. Schumi, <i>Anti-commutative algebras and homogeneous spaces with multiplications</i>	255
Robert Evert Stong, <i>Stiefel-Whitney classes of manifolds</i>	271
D. Suryanarayana, <i>On a theorem of Apostol concerning Möbius functions of order k</i>	277
Yoshio Tanaka, <i>On closedness of C- and C*-embeddings</i>	283
Chi Song Wong, <i>Characterizations of certain maps of contractive type</i>	293