

Pacific Journal of Mathematics

**AN ANALOGUE OF OKA'S THEOREM FOR WEAKLY
NORMAL COMPLEX SPACES**

WILLIAM ALLEN ADKINS, ALDO ANDREOTTI AND JOHN VINCENT LEAHY

AN ANALOGUE OF OKA'S THEOREM FOR WEAKLY NORMAL COMPLEX SPACES

WILLIAM A. ADKINS, ALDO ANDREOTTI, J. V. LEAHY

Two well known results concerning normal complex spaces are the following. First, the singular set of a normal complex space has codimension at least two. Second, this property characterizes normality for complex spaces which are local complete intersections. This second result is a theorem of Abhyankar [1] which generalizes Oka's theorem. The purpose of this paper is to prove analogues of these facts for the class of weakly normal complex spaces, which were introduced by Andreotti-Norguet [3] in a study of the space of cycles on an algebraic variety. A weakly normal complex space can have singularities in codimension one, but it will be shown that an obvious class of such singularities is generic.

1. Preliminaries. All complex spaces are assumed to be reduced. If X is a complex space, there is the sheaf \mathcal{O}_X of holomorphic functions on X , and the sheaf \mathcal{O}_X^c of c -holomorphic functions on X . A section of \mathcal{O}_X^c on an open subset U of X is a continuous function $f: U \rightarrow \mathbb{C}$ such that f is holomorphic on the regular points of U . The complex space X is said to be weakly normal if $\mathcal{O}_X = \mathcal{O}_X^c$. Examples of weakly normal spaces are normal spaces and unions of submanifolds of \mathbb{C}^m in general position.

Let $V_j = \{(x_1, \dots, x_m) \in \mathbb{C}^m : x_k = 0 \text{ for } n \leq k < j \text{ and } j < k \leq m\}$ where $n \leq j \leq m$. Then V_j is an n -dimensional linear subspace of \mathbb{C}^m . Let

$$V_{(n,m)} = \bigcup_{j=n}^m V_j = \{(x_1, \dots, x_m) \in \mathbb{C}^m : x_i x_j = 0 \text{ for } n \leq i < j \leq m\}$$

and let $S(V_{(n,m)})$ be the singular set of $V_{(n,m)}$.

LEMMA. $V_{(n,m)}$ is a weakly normal complex space and $\dim S(V_{(n,m)}) = n - 1$.

Proof. Since $S(V_{(n,m)}) = \{(x_1, \dots, x_m) \in \mathbb{C}^m : x_n = \dots = x_m = 0\}$, $\dim S(V_{(n,m)}) = n - 1$. Let $f: V_{(n,m)} \rightarrow \mathbb{C}$ be a continuous function which is holomorphic on the regular points of $V_{(n,m)}$. To prove weak normality of $V_{(n,m)}$, we need to show that f is holomorphic. Let $f_j = f|_{V_j}$. By the Riemann extension theorem, f_j is holomorphic on the n -plane V_j and

thus $f_j = f_j(x', x_j)$ is a convergent power series, where $x' = (x_1, \dots, x_{n-1})$ and x_j are coordinates on V_j . Since $f_j|_{x_j=0} = f_k|_{x_k=0}$ for $n \leq j, k \leq m$, we let $f_0(x') = f_j(x', 0)$ and set $g_j(x', x_j) = f_j(x', x_j) - f_0(x')$ for $n \leq j \leq m$. Then $f(x_1, \dots, x_m) = f_0(x') + \sum_{j=n}^m g_j(x', x_j)$ and hence f is holomorphic on $V_{(n,m)}$.

If X is a complex space with $\dim X = n$, let $Sg(X) = S(X) \cup (\bigcup_{0 \leq k < n} X^{(k)})$ where $S(X)$ is the singular set of X and $X^{(k)}$ is the analytic subset of X defined by $X^{(k)} = \{x \in X: X \text{ has a branch of dimension } k \text{ at } x\}$. If $C_4(X, x)$ denotes the fourth Whitney tangent cone of X at x , then Stutz [6] has shown that $W_4 = Sg(X) \cap \{x \in X: \dim C_4(X, x) > n\}$ is an analytic subset of X of codimension at least two.

2. Codimension one singularities of weakly normal spaces. Let X be a complex space. A point $x \in X$ is said to be an elementary point of type (n, m) , for $n \leq m$, if the germ (X, x) is isomorphic to the germ $(V_{(n,m)}, 0)$. Note that if $x \in X$ is an elementary point of type (n, m) , then the germ (X, x) is of pure dimension n and the imbedding dimension of (X, x) is m . The set of elementary points of X contains the set of regular points of X , i.e. the elementary points of type (n, n) for some n . In addition, it contains a particularly simple class of singular points of X . If x is an elementary point of type (n, m) with $n < m$, then x is singular and $\dim(S(X), x) = n - 1 = \dim(X, x) - 1$.

If $\dim X = n$, let $Y = \bigcup_{0 \leq k < n} X^{(k)}$ and let $X_1 = \overline{X \setminus Y}$. By a theorem of Remmert, X_1 is an analytic set of pure dimension n . Let X_s denote the set of all elementary points of X of type (n, m) for some m with $m \geq n = \dim X$. Hence $X_s \subseteq X_1$ and X_s contains the regular points of X of maximal dimension.

THEOREM 1. *Let X be a weakly normal complex space. Then $A = X_1 \setminus X_s$ is an analytic subset of X_1 of codimension at least 2.*

Proof. Let $n = \dim X$. If $\dim S(X) \leq n - 2$ then $A = X_1 \cap S(X)$. Hence A is analytic and codimension $A \geq 2$. Now suppose that $\dim S(X) = n - 1$. We will show that $A = X_1 \cap (Sg(Sg(X)) \cup W_4)$. Since $Sg(Sg(X)) \cup W_4$ is an analytic set of codimension at least 2 in X and since $\dim X = \dim X_1$, this will prove the theorem.

Let $x \in X_s$. If x is a regular point of X , then $x \notin Sg(Sg(X)) \cup W_4$. If x is an elementary point of type (n, m) where $m > n$, then $\dim C_4(X, x) = n$. Hence $x \notin W_4$. Moreover, $S(X)$ is a manifold of dimension $n - 1$ in a neighborhood of x . Thus $x \notin Sg(Sg(X))$. Hence $X_s \subseteq X_1 \setminus (Sg(Sg(X)) \cup W_4)$ and $X_1 \cap (Sg(Sg(X)) \cup W_4) \subseteq A$.

Now suppose that $x_0 \in X_1 \cap S(X) \cap (X_1 \setminus (Sg(Sg(X)) \cup W_4))$. Thus

$x_0 \in \text{Sg}(X) \setminus \text{Sg}(\text{Sg}(X))$ and $\dim C_4(X, x_0) = n$. Note also that the germ (X, x_0) is of pure dimension n . Since the result to be proved is local, we may assume that $X \subseteq \mathbb{C}^t$. By Proposition 4.2 of Stutz [6], there is a neighborhood N of x_0 in X , a polydisc $D \subseteq \mathbb{C}^n$, and a choice of coordinates x_1, \dots, x_n in \mathbb{C}^n and y_1, \dots, y_t in \mathbb{C}^t centered at x_0 with the following properties.

If B_0, \dots, B_r are the global branches of $X \cap N$, then for each j ($0 \leq j \leq r$) there is a holomorphic map $f_j: D \rightarrow B_j$ such that

- (a) f_j is a homeomorphism;
- (b) with respect to the coordinates $x_1, \dots, x_n, y_1, \dots, y_t, f_j(0) = 0$ and

$$f_j(x) = (x_1, \dots, x_{n-1}, x_n^{p_j}, f_{n+1,j}(x), \dots, f_t(x))$$

where p_j is a positive integer for $0 \leq j \leq r$;

- (c) $f_j(x_1, \dots, x_n) = \sum_{v=p_j}^{\infty} f_{ij}^{(v)}(x_1, \dots, x_{n-1}) \cdot x_n^v$ for $n+1 \leq i \leq t$ and $0 \leq j \leq r$.

Let $g_j: B_j \rightarrow D$ be the continuous inverse of f_j and define a map $h: X \cap N \rightarrow \mathbb{C}^{n+r}$ by $\pi_j \circ h|_{B_j} = g_j$ where $\pi_j: \mathbb{C}^{n+r} \rightarrow \mathbb{C}_{x_1, \dots, x_{n-1}, x_{n+j}}$ is the natural linear projection onto the n -plane with coordinates $x_1, \dots, x_{n-1}, x_{n+j}$, for $0 \leq j \leq r$. To see that the map h is well defined, note first that $S(X)$ is an $n-1$ dimensional manifold in a neighborhood of x_0 . Furthermore, $B_j \cap B_k \subseteq S(X) \cap N$ for all j, k . But $f_j(x', 0) = (x', 0, \dots, 0) = f_k(x', 0)$ where $x' = (x_1, \dots, x_{n-1})$. Therefore, if N is chosen small enough, then $B_j \cap B_k = S(X) \cap N = \{y_n = \dots = y_t = 0\}$ for $0 \leq j, k \leq r$. For each $(y_1, \dots, y_t) \in S(X) \cap N$, it follows that $g_j(y) = (y_1, \dots, y_{n-1}, 0)$ for $0 \leq j \leq r$. Thus h is a well defined continuous map.

Since the jacobian matrix $\partial f_j / \partial x$ is given by

$$\frac{\partial f_j}{\partial x} = \begin{bmatrix} I_{n-1} & 0 \\ 0 & p_j x_n^{p_j-1} \\ * & * \end{bmatrix}$$

h is holomorphic on the regular points of $X \cap N$. Since X is weakly normal and h is a homeomorphism onto its image, it follows that h is biholomorphic. Therefore x_0 is an elementary singularity of type $(n, n+r)$. Hence $A \subseteq X_1 \cap (\text{Sg}(\text{Sg}(X)) \cup W_4)$ and the theorem is proved.

REMARK. Let X be a weakly normal complex space and suppose that $\text{codim } S(X) = 1$. Theorem 1 shows that there is an elementary singularity of type (n, m) where $m > n$. Since such a singular point is not normal, Theorem 1 implies the well-known theorem that $\text{codim } S(X) \geq 2$ for a normal complex space X .

THEOREM 2. *Let X be a pure dimensional local complete intersection. Then X is weakly normal if and only if $\text{codim } X \setminus X_s \geq 2$.*

Proof. Let $A = X \setminus X_s$. If X is weakly normal then $\text{codim } A \geq 2$ by Theorem 1.

Conversely, suppose $\text{codim } A \geq 2$. Since $X \setminus A = X_s$, the germ (X, x) is weakly normal for each $x \in X \setminus A$. Since X is a pure dimensional local complete intersection, $pf(\mathcal{O}_{X,x}) = \dim X$ for each $x \in X$, where pf = profondeur. From the Hartog theorem for weak normality [2], we conclude that X is weakly normal.

REMARKS. (1) For the case of curves, the assumption of local complete intersection is not needed. A curve X is weakly normal if and only if $X \setminus X_s = \emptyset$. An algebraic proof of this fact was given by Bombieri [5].

(2) If X is a pure dimensional hypersurface in \mathbf{C}^{n+1} , then Theorem 2 can be proved without the use of the Hartog theorem for weak normality. This case follows from the result of Becker in [4].

(3) Let $X \subseteq \mathbf{C}^{n+1}$ be a pure dimensional hypersurface. If X is weakly normal, there is another characterization of $X \setminus X_s$, than that which is given by the proof of Theorem 1. This description is as follows. There is a holomorphic function $f \in \mathcal{O}(\mathbf{C}^{n+1})$ such that $X = V(f) = \{x \in \mathbf{C}^{n+1} : f(x) = 0\}$ and such that there is a sheaf equality $(f) \cdot \mathcal{O} = \mathcal{I}_X$ where \mathcal{I}_X is the sheaf of ideals of X . Then

$$S(X) = \left\{ x \in X : \frac{\partial f}{\partial z_i}(x) = 0 \text{ for } 1 \leq i \leq n+1 \right\}.$$

At a point $x_0 \in S(X)$ the Hessian form is defined by

$$H(f)_{x_0}(u) = \sum_{i,j=1}^{n+1} \frac{\partial^2 f}{\partial z_i \partial z_j}(x_0) \cdot u_i u_j.$$

Let $\mu(x_0) = \text{rank } H(f)_{x_0}$ and set $S_2(X) = \{x \in S(X) : \mu(x) \leq 1\}$.

Claim. *If X is weakly normal and $\dim S(X) = n - 1$, then*

$$W_4 \cap (S(X) \setminus \text{Sg}(S(X))) = S_2 \cap (S(X) \setminus \text{Sg}(S(X))).$$

Proof. From the proof of Theorem 1, $X \setminus X_s = \text{Sg}(S(X)) \cup W_4$. Suppose $x \in S(X) \setminus \text{Sg}(S(X))$ but $x \notin W_4$. Then the proof of Theorem 1 shows that x is an elementary singular point of type $(n, n+1)$. A proper choice of local coordinates about x shows that (X, x) is isomorphic to $(V(z_1 z_2), 0)$. Hence $\mu(x) = 2$ and $x \notin S_2(X)$.

Now suppose that $x \in S(X) \setminus \text{Sg}(S(X))$ but $x \notin S_2(X)$. Thus $\mu(x) \geq 2$. If $\mu(x) > 2$ then the implicit function theorem shows that $\dim(S(X), x) \leq n - 2$. Therefore $\mu(x) = 2$ and choosing convenient local coordinates centered at x gives $f(z) = az_1z_2 + 0(3)$ where $a \neq 0$. Hence x is an elementary singular point of type $(n, n + 1)$. Therefore, $x \notin W_4$ and the claim is proved.

For weakly normal hypersurfaces this claim gives an easy differential criterion for computing the portion of the set W_4 which is contained in $S(X) \setminus \text{Sg}(S(X))$. This claim is false for hypersurfaces which are not weakly normal.

EXAMPLE. Let $X = \{(x, y, z) \in \mathbb{C}^3: x^2 - zy^2 = 0\}$ be the Cayley umbrella in \mathbb{C}^3 . Then $X \setminus X_s = \{(0, 0, 0)\}$ so that X is weakly normal by Theorem 2. Remark (3) then shows that $W_4 = \{0, 0, 0\}$.

REFERENCES

1. Shreeram Abhyankar, *Concepts of order and rank on a complex space and a condition for normality*, Math. Annalen, **141** (1960), 171–192.
2. Aldo Andreotti and Per Holm, *Parametric spaces*, preprint.
3. Aldo Andreotti and Francois Norguet, *La convexite holomorphe dans l'espace analytique des cycles d'une variete algebrique*, Ann. Scuola Norm. Sup. Pisa, s. 3, **21** (1967), 31–82.
4. Joseph Becker, *Normal Hypersurfaces*, Pacific J. Math. **61** (1975), 17–19.
5. Enrico Bombieri, *Seminormalite e singolarite ordinarie*, Symposia Mathematica XI, Academic Press, (1973), 205–210.
6. John Stutz, *Analytic sets as branched coverings*, Trans. Amer. Math. Soc., **166** (1972), 241–259.

Received December 20, 1976.

INSTITUTE FOR ADVANCED STUDY
PRINCETON, NJ 08540

OREGON STATE UNIVERSITY
CORVALLIS, OR 97331

AND

UNIVERSITY OF OREGON
EUGENE, OR 97403

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, CA 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. A. BEAUMONT
University of Washington
Seattle, WA 98105

R. FINN AND J. MILGRAM
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

• • •
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate, may be sent to any one of the four editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$ 72.00 a year (6 Vols., 12 issues). Special rate: \$ 36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to *Pacific Journal of Mathematics*, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Printed at Jerusalem Academic Press, POB 2390, Jerusalem, Israel.

Copyright © 1977 Pacific Journal of Mathematics
All Rights Reserved

Pacific Journal of Mathematics

Vol. 68, No. 2

April, 1977

William Allen Adkins, Aldo Andreotti and John Vincent Leahy, <i>An analogue of Oka's theorem for weakly normal complex spaces</i>	297
Ann K. Boyle, M. G. Deshpande and Edmund H. Feller, <i>On nonsingularly k-primitive rings</i>	303
Rolando Basim Chuaqui, <i>Measures invariant under a group of transformations</i>	313
Wendell Dan Curtis and Forrest Miller, <i>Gauge groups and classification of bundles with simple structural group</i>	331
Garret J. Etgen and Willie Taylor, <i>The essential uniqueness of bounded nonoscillatory solutions of certain even order differential equations</i>	339
Paul Ezust, <i>On a representation theory for ideal systems</i>	347
Richard Carl Gilbert, <i>The deficiency index of a third order operator</i>	369
John Norman Ginsburg, <i>S-spaces in countably compact spaces using Ostaszewski's method</i>	393
Basil Gordon and S. P. Mohanty, <i>On a theorem of Delaunay and some related results</i>	399
Douglas Lloyd Grant, <i>Topological groups which satisfy an open mapping theorem</i>	411
Charles Lemuel Hagopian, <i>A characterization of solenoids</i>	425
Kyong Taik Hahn, <i>On completeness of the Bergman metric and its subordinate metrics. II</i>	437
G. Hochschild and David Wheeler Wigner, <i>Abstractly split group extensions</i>	447
Gary S. Itzkowitz, <i>Inner invariant subspaces</i>	455
Jiang Luh and Mohan S. Putcha, <i>A commutativity theorem for non-associative algebras over a principal ideal domain</i>	485
Donald J. Newman and A. R. Reddy, <i>Addendum to: "Rational approximation of e^{-x} on the positive real axis"</i>	489
Akio Osada, <i>On the distribution of a-points of a strongly annular function</i>	491
Jeffrey Lynn Spielman, <i>A characterization of the Gaussian distribution in a Hilbert space</i>	497
Robert Moffatt Stephenson Jr., <i>Symmetrizable-closed spaces</i>	507
Peter George Trotter and Takayuki Tamura, <i>Completely semisimple inverse Δ-semigroups admitting principal series</i>	515
Charles Irvin Vinsonhaler and William Jennings Wickless, <i>Torsion free abelian groups quasi-projective over their endomorphism rings</i>	527
Frank Arvey Wattenberg, <i>Topologies on the set of closed subsets</i>	537
Richard A. Zalik, <i>Integral representation of Tchebycheff systems</i>	553