

Pacific Journal of Mathematics

THE RANGE OF ANALYTIC EXTENSIONS

JOSIP GLOBEVNIK

THE RANGE OF ANALYTIC EXTENSIONS

J. GLOBEVNIK

Denote by Δ , $\bar{\Delta}$, $\partial\Delta$ the open unit disc in C , its closure and its boundary, respectively. Let X be a complex Banach space and denote by $\mathcal{A}(X)$ the class of all non-empty sets $P \subset X$ having the following property: given any closed set $F \subset \partial\Delta$ of measure 0 and any continuous function $f: F \rightarrow P$ there exists a continuous extension $\tilde{f}: \bar{\Delta} \rightarrow X$ of f , analytic on Δ and satisfying $\tilde{f}(\bar{\Delta} - F) \subset \text{Int } P$.

THEOREM. $P \in \mathcal{A}(X)$ if and only if $\text{Int } P$ is connected, locally connected at every point of P and satisfies $P \subset \text{closure}(\text{Int } P)$.

THEOREM. If $P \subset C$ consists of more than one point then $P \in \mathcal{A}(C)$ if and only if given any F and f as above there exists a continuous extension $\tilde{f}: \bar{\Delta} \rightarrow C$ of f , analytic on Δ and satisfying $\tilde{f}(\bar{\Delta}) \subset P$.

This generalizes a theorem of Rudin which asserts that such \tilde{f} exists if $P \subset C$ is homeomorphic to $\bar{\Delta}$.

THEOREM. If $P \in \mathcal{A}(X)$ then given any relatively open set $B \subset \partial\Delta$, any relatively closed set $F \subset B$ of measure 0 and any continuous function $f: F \rightarrow P$ there exists a continuous extension $\tilde{f}: \Delta \cup B \rightarrow X$ of f , analytic on Δ and satisfying $\tilde{f}((\Delta \cup B) - F) \subset \text{Int } P$.

0. Introduction. Throughout, we denote by Δ , $\bar{\Delta}$ and $\partial\Delta$ the open unit disc in C , its closure and its boundary, respectively. If X is a complex Banach space and $r > 0$ we write $B_r(X) = \{x \in X: \|x\| < r\}$. Let $x \in X$ and $S, T \subset X$. We write $x + S = \{x + u: u \in S\}$ and $S + T = \{u + v: u \in S, v \in T\}$. We denote by $\text{Int } S$, \bar{S} the interior of S and the closure of S , respectively. If F is a compact Hausdorff space we denote by $C(F, X)$ the set of all continuous functions from F to X and write $C(F)$ for $C(F, C)$. If $B \subset \partial\Delta$ is a relatively open set we denote by $H_B(\Delta, X)$ the set of all continuous functions from $\Delta \cup B$ to X which are analytic on Δ . For $H_{\partial\Delta}(\Delta, X)$ we write $A(\Delta, X)$ and for $A(\Delta, C)$, the disc algebra, we write $A(\Delta)$. We denote the set of all positive integers by N . If $a, b \in R$, $a < b$ we write $[a, b] = \{t \in R: a \leq t \leq b\}$ and we denote $[0, 1]$ by I .

The well known Rudin-Carleson theorem [3, 19, 22] states that given a closed set $F \subset \partial\Delta$ of measure 0 and $f \in C(F)$ there exists an extension $\tilde{f} \in A(\Delta)$ of f satisfying

$$\max_{z \in \bar{\Delta}} |\tilde{f}(z)| = \max_{s \in F} |f(s)|.$$

Actually the following theorem was proved by Rudin: Given F as above, let $P \subset C$ be homeomorphic to \bar{A} and let $f \in C(F)$ satisfy $f(F) \subset P$. There exists an extension $\tilde{f} \in A(\Delta)$ of f satisfying $\tilde{f}(\bar{A}) \subset P$. An interesting consequence is that given any $P \subset C$ homeomorphic to \bar{A} there exists $\tilde{f} \in A(\Delta)$ satisfying $\tilde{f}(\bar{A}) = \tilde{f}(\partial\Delta) = P$ [2, 4].

The Rudin-Carleson theorem was generalized into several directions. A consequence of its generalization to the functions into a Banach space [21, 18, 7] is that for any finite-dimensional complex normed space X there exists $\tilde{f} \in A(\Delta, X)$ such that $\tilde{f}(\bar{A}) = \overline{B_1(X)}$ [7]. Heard and Wells [12] generalized the Rudin-Carleson theorem as follows: Let $B \subset \partial\Delta$ be a relatively open set and $F \subset B$ a relatively closed set of measure 0. Given any bounded continuous function $f: F \rightarrow C$ there exists an extension $\tilde{f} \in H_B(\Delta, C)$ of f satisfying

$$\sup_{z \in \Delta} |\tilde{f}(z)| = \sup_{s \in F} |f(s)|.$$

The generalization of this result to the functions into a Banach space X [8] makes possible, in the case when X is separable, to prove the existence of a continuous function $\tilde{f}: \bar{A} - \{1\} \rightarrow X$, analytic on Δ , whose range is contained and dense in $\overline{B_1(X)}$ and whose cluster set at 1 is $\overline{B_1(X)}$ [8].

The applications above seem interesting enough to consider the following general problem.

PROBLEM. Let X be a complex Banach space, $B \subset \partial\Delta$ a relatively open set, $F \subset B$ a relatively closed set of measure 0 and let $f: F \rightarrow X$ be a continuous function. Assume that a subset P of X contains $f(F)$. Under what conditions on P does there exist an extension $\tilde{f} \in H_B(\Delta, X)$ of f satisfying $\tilde{f}(\Delta \cup B) \subset P$?

By the results mentioned above such an \tilde{f} exists if $P \subset X$ is a closed ball. To prove this one needs the fact that the subspace of all bounded functions in $H_B(\Delta, X)$ is a left $A(\Delta)$ -module and the fact that P is absolutely convex in order to make the necessary norm estimations on the interpolating function \tilde{f} assuring that $\tilde{f}(\Delta \cup B) \subset P$ [8]. Nothing similar is true in general when we consider the functions in $H_B(\Delta, X)$ whose ranges are contained in other sets than balls. Consequently one has to apply different techniques in the general case. In [11] this was done in the special case when the set P was open and it was proved that \tilde{f} above exists if P is (open and) connected. Of course this was not a generalization of the Rudin-Carleson theorem although it was enough to reprove the main result of [9]: Given any open connected set P in a separable complex Banach space X there exists a continuous function $\tilde{f}: \bar{A} - \{1\} \rightarrow X$, analytic on Δ , whose range is contained and dense in P .

In the present paper we study the general case when the set $P \subset X$ is not necessarily open. In the special case when $X = C$ we obtain a simple complete topological description of the sets $P \subset C$ having the following property: given any closed set $F \subset \partial \Delta$ of measure 0 and any $f \in C(F)$ satisfying $f(F) \subset P$ there exists an extension $\tilde{f} \in A(\Delta)$ of f satisfying $\tilde{f}(\bar{\Delta}) \subset P$. If $P \subset C$ has such a property and if P consists of more than one point we show that for every relatively open set $B \subset \partial \Delta$, every relatively closed set $F \subset B$ of measure 0 and every continuous function $f: F \rightarrow P$ there exists a "peak" extension of f , i.e. an extension $\tilde{f} \in H_b(\Delta, C)$ of f satisfying $\tilde{f}((\Delta \cup B) - F) \subset \text{Int } P$. In the general case we study only the sets $P \subset X$ with the property that given any closed set $F \subset \partial \Delta$ of measure 0, every function $f \in C(F, X)$ satisfying $f(F) \subset P$ admits a peak extension $\tilde{f} \in A(\Delta, X)$, and obtain their topological description.

In §1 we state the main results. In §2 we give the complete proofs; this section contains some lemmas and theorems which might be of independent interest. In §3 we present some simple applications to the ranges and cluster sets of analytic functions.

1. Main results.

DEFINITION 1. [10, 11] Let $B \subset \partial \Delta$ be a relatively open set. A subset P of a complex Banach space X is said to have the *analytic extension property* (AEP) with respect to $H_b(\Delta, X)$ if, given any relatively closed set $F \subset B$ of measure 0 and any continuous function $f: F \rightarrow P$ there exists an extension $\tilde{f} \in H_b(\Delta, X)$ of f which satisfies $\tilde{f}(\Delta \cup B) \subset P$. We say that $P \subset X$ has AEP if it has AEP with respect to $H_b(\Delta, X)$ for every relatively open set $B \subset \partial \Delta$.

DEFINITION 2. Let $B \subset \partial \Delta$ be a relatively open set. A subset P of a complex Banach space X is said to have the *peak analytic extension property* (PAEP) with respect to $H_b(\Delta, X)$ if, given any F and f as above there exists an extension $\tilde{f} \in H_b(\Delta, X)$ of f satisfying $\tilde{f}((\Delta \cup B) - F) \subset \text{Int } P$. We call every such extension a *peak extension with respect to P* (whether P has PAEP or not). We say that $P \subset X$ has PAEP if it has PAEP with respect to $H_b(\Delta, X)$ for every relatively open set $B \subset \partial \Delta$.

Let O be an open subset of a complex Banach space X . O is called *locally connected* (LC) at a point $x \in X$ if given any $\varepsilon > 0$ there exists $\delta > 0$ such that if $(x + B_\delta(X)) \cap O$ is not empty it is contained in a connected component of $(x + B_\varepsilon(X)) \cap O$ [17]. Note that O is LC at every point of O .

Now we are able to state our main results.

THEOREM 1. *Let P be a nonempty subset of a complex Banach space X . Then the following are equivalent*

- (A) *there exists a closed set $F \subset \partial\Delta$ of measure 0 with infinitely many points such that every continuous function $f: F \rightarrow P$ admits a peak extension $\tilde{f} \in A(\Delta, X)$ with respect to P*
- (B) *P has PAEP with respect to $A(\Delta, X)$*
- (C) *P has PAEP*
- (D) *P has the following properties:*
 - (i) *$P \subset \overline{\text{Int } P}$.*
 - (ii) *$\text{Int } P$ is connected and locally connected at every point of P .*

THEOREM 2. *Let P be a subset of C containing more than one point. Then the following are equivalent*

- (A) *there exists a closed set $F \subset \partial\Delta$ of measure 0 with infinitely many points such that every continuous function $f: F \rightarrow P$ admits an extension $\tilde{f} \in A(\Delta)$ satisfying $\tilde{f}(\bar{\Delta}) \subset P$*
- (B) *P has AEP*
- (C) *P has PAEP.*

COROLLARY. *Let P be a subset of C containing more than one point. Then the following are equivalent*

- (A) *given any closed set $F \subset \partial\Delta$ of measure 0 and any $f \in C(F)$ satisfying $f(F) \subset P$ there exists an extension $\tilde{f} \in A(\Delta)$ of f satisfying $\tilde{f}(\bar{\Delta}) \subset P$*
- (B) *given any closed set $F \subset \partial\Delta$ of measure 0 and any $f \in C(F)$ satisfying $f(F) \subset P$ there exists an extension $\tilde{f} \in A(\Delta)$ of f satisfying $\tilde{f}(\bar{\Delta} - F) \subset \text{Int } P$*
- (C) *P has the following properties*
 - (i) *$P \subset \overline{\text{Int } P}$*
 - (ii) *$\text{Int } P$ is connected and locally connected at every point of P .*

REMARK. The above Corollary gives a complete topological description of the sets $P \subset C$ for which the Rudin theorem holds. If (C) in the Corollary above is satisfied by a (nonempty) compact set $P \subset C$ whose interior is simply connected note that then P is homeomorphic to $\bar{\Delta}$ [17]. Finally, note that the main results of [7, 8, 9, 11] follow from Theorem 1 and Theorem 2 above.

2. Proofs.

LEMMA 1. Let P be a nonempty subset of a complex Banach space X and let $F \subset \partial\Delta$ be a closed set of measure 0 containing infinitely many points. Suppose that given any continuous function $f: F \rightarrow P$ there exists an extension $\tilde{f} \in A(\Delta, X)$ of f such that $\tilde{f}(\bar{\Delta} - F) \subset \text{Int } P$. Then

- (i) $P \subset \overline{\text{Int } P}$
- (ii) $\text{Int } P$ is connected
- (iii) $\text{Int } P$ is locally connected at every point of P .

Proof. Let $x \in P$. By the assumption there exists $\tilde{f} \in A(\Delta, X)$ satisfying $\tilde{f}(F) = \{x\}$ and $\tilde{f}(\bar{\Delta} - F) \subset \text{Int } P$. Let $\{z_n\} \subset \Delta$ converge to a point of F . By the continuity of \tilde{f} $\{\tilde{f}(z_n)\}$ converges to x . Since $\tilde{f}(z_n) \in \text{Int } P$ ($n \in N$) (i) is proved.

By the assumption F is nowhere dense on $\partial\Delta$ and contains more than one point. Consequently $F = F_1 \cup F_2$ where F_1, F_2 are nonempty disjoint compact sets. Let $x, y \in \text{Int } P$ and define $f(s) = x$ ($s \in F_1$), $f(s) = y$ ($s \in F_2$). Clearly $f \in C(F, X)$ and by the assumption there exists $\tilde{f} \in A(\Delta, X)$ satisfying $\tilde{f}(F_1) = \{x\}$, $\tilde{f}(F_2) = \{y\}$ and $\tilde{f}(\bar{\Delta} - F) \subset \text{Int } P$. Let $z_1 \in F_1, z_2 \in F_2$. Now $t \mapsto \varphi(t) = \tilde{f}(z_1 + t(z_2 - z_1))$ is a path joining x and y . By the properties of f we have $\varphi(I) \subset \text{Int } P$ which proves (ii).

To prove (iii) assume that $\text{Int } P$ is not LC at a point $x \in P$. This means that there exists $\varepsilon > 0$ such that for every $\delta > 0$ the set $(x + B_\delta(X)) \cap \text{Int } P$ meets at least two connected components of the set $(x + B_\varepsilon(X)) \cap \text{Int } P$. It follows that there exist two sequences $\{x_n\} \subset \text{Int } P, \{y_n\} \subset \text{Int } P$ converging to x such that for every $n \in N$ x_n and y_n lie in different components of $(x + B_\varepsilon(X)) \cap \text{Int } P$. Assume for a moment that there exist a sequence $\{t_n\} \subset \partial\Delta$ converging to $t \in \partial\Delta$ and $\tilde{f} \in A(\Delta, X)$ satisfying $\tilde{f}(\Delta) \subset \text{Int } P$ and $\tilde{f}(t_{2n-1}) = x_n, \tilde{f}(t_{2n}) = y_n$ ($n \in N$). By the continuity of \tilde{f} there exists a neighbourhood $U \subset \bar{\Delta}$ of t such that $\tilde{f}(U) \subset x + B_\varepsilon(X)$. Consequently there is some $n \in N$ such that the closed segment J joining t_{2n-1} and t_{2n} is contained in U . Then by the properties of \tilde{f} , $\tilde{f}(J)$ is a path joining x_n and y_n in $(x + B_\varepsilon(X)) \cap \text{Int } P$, a contradiction.

It remains to prove the existence of f and $\{t_n\}$ above. By the assumption F contains infinitely many points which implies that there is some $t \in F$ which is a cluster point of $F - \{t\}$. Since F is nowhere dense on $\partial\Delta$ there exists a decreasing sequence $\{T_n\}$ of open arcs in $\partial\Delta$ all of whose endpoints lie in $\partial\Delta - F$ and such that $F \subset T_1, \bigcap_{n=1}^\infty T_n = \{t\}$. Define $F_n = F \cap (T_n - T_{n+1})$ ($n \in N$). Then F_n ($n \in N$) are disjoint compact sets infinitely many of which are not empty. Passing to a subsequence if necessary we may assume that all sets F_n ($n \in N$) are nonempty. Define $f: F \rightarrow X$ by

$$\begin{aligned} f(s) &= x_n \quad (s \in F_{2n-1}; n \in N), \\ f(s) &= y_n \quad (s \in F_{2n}; n \in N), \\ f(t) &= x. \end{aligned}$$

It is easy to see that f is continuous. By the assumption there exists an extension $\tilde{f} \in A(\Delta, X)$ of f satisfying $\tilde{f}(\bar{\Delta} - F) \subset \text{Int } P$. Choose $t_n \in F_n (n \in N)$.

An open subset O of a complex Banach space X is called *uniformly locally connected* (ULC) on a subset K of X if given any $\varepsilon > 0$ there exists $\delta > 0$ such that for every $x \in K$ the set $(x + B_\delta(X)) \cap O$, if not empty, is contained in a connected component of the set $(x + B_\varepsilon(X)) \cap O$ [17]. We call any such $\varepsilon \mapsto \delta(\varepsilon)$ a *modulus of ULC* of O on K .

LEMMA 2. *Let O be an open subset of a complex Banach space X and let $K \subset X$ be a compact set. Suppose that O is locally connected at each point $x \in K$. Then O is uniformly locally connected on K .*

Proof. Simple. For the idea see [17, p. 160].

LEMMA 3. [11] *Let $B \subset \partial\Delta$ be a relatively open set, $G \subset B$ a relatively closed set of measure 0, and $H \subset B$ a compact set of measure 0, disjoint from G . Let $U \subset \Delta \cup B$ be a neighbourhood of H and let $\varepsilon > 0$. Assume that P is an open connected set in a complex Banach space X which contains the point 0 and let $f: H \rightarrow P$ be a continuous function. There exists $\tilde{f} \in H_B(\Delta, X)$ which satisfies*

- (i) $\tilde{f}|_H = f$
- (ii) $\tilde{f}(G) = \{0\}$
- (iii) $\|\tilde{f}(z)\| < \varepsilon \quad (z \in (\Delta \cup B) - U)$
- (iv) $\tilde{f}(\Delta \cup B) \subset P$.

Lemma 3, which gives an approximate solution to our Problem, is the most important tool in the present paper.

LEMMA 4. *Let O be an open connected set in a complex Banach space X . Let $F \subset \partial\Delta$ be a closed set of measure 0 and f_1, f_2 two continuous functions from F to O . There exists a homotopy $p: F \times I \rightarrow O$ satisfying*

$$\begin{aligned} p(s, 0) &= f_1(s) \\ p(s, 1) &= f_2(s) \end{aligned} \Big\} (s \in F).$$

Proof. Choose $s_0 \in \partial A - F$ and $x \in O$. By Lemma 3 there exist $g_i \in A(A, X)$ ($i = 1, 2$) satisfying

$$\left. \begin{aligned} g_i|F &= f_i \\ g_i(\bar{A}) &\subset O \\ g_i(s_0) &= x \end{aligned} \right\} (i = 1, 2).$$

Put

$$p(s, t) = \begin{cases} g_1(s + 2t(s_0 - s)) & (s \in F, 0 \leq t \leq 1/2) \\ g_2(s_0 - (1 - 2t)(s - s_0)) & (s \in F, 1/2 \leq t \leq 1). \end{cases}$$

It is easy to check that p has the required properties.

LEMMA 5. Let O be an open subset of a complex Banach space and let $F \subset \partial A$ be a closed set of measure 0. Suppose that $f: F \rightarrow \bar{O}$ is a continuous function and assume that O is ULC on $f(F)$. Let $\varepsilon \mapsto \delta(\varepsilon)$ be a modulus of ULC of O on $f(F)$. Let $R > 0$ and assume that $g: F \rightarrow O$ is a continuous function satisfying

$$\|f(s) - g(s)\| < \delta(R)/2 \quad (s \in F).$$

Let $\varepsilon > 0$. There exists a homotopy $\pi: F \times I \rightarrow O$ satisfying

- (i) $\pi(s, 0) = g(s) \quad (s \in F)$,
- (ii) $\|\pi(s, 1) - f(s)\| < \varepsilon \quad (s \in F)$,
- (iii) $\|\pi(s, t) - f(s)\| < 2R \quad (s \in F, 0 \leq t \leq 1)$.

Proof. By the properties of F

$$F = \bigcup_{i=1}^m F_i$$

where F_i ($i = 1, 2, \dots, m$) are disjoint compact sets such that

$$(1.1) \quad \|f(\xi) - f(\eta)\| < \min\{R, \delta(R)/2, \varepsilon/2\} \quad (\xi, \eta \in F_i; i = 1, 2, \dots, m).$$

Choose $\zeta_i \in F_i$ ($i = 1, 2, \dots, m$). Since $f(F) \subset \bar{O}$ there exist $x_i \in O$ ($i = 1, 2, \dots, m$) such that

$$(1.2) \quad \|f(\zeta_i) - x_i\| < \min\{\delta(R)/2, \varepsilon/2\} \quad (i = 1, 2, \dots, m).$$

Define $h(s) = x_i$ ($s \in F_i; i = 1, 2, \dots, m$). By the assumption and by (1.1) we have

$$(1.3) \quad \begin{aligned} \|f(\zeta_i) - g(s)\| &\leq \|f(\zeta_i) - f(s)\| + \|f(s) - g(s)\| \\ &< \delta(R) \end{aligned} \quad (s \in F_i; i = 1, 2, \dots, m)$$

and similarly by (1.1) and (1.2)

$$(1.4) \quad \begin{aligned} \|h(s) - f(s)\| &\leq \|h(s) - f(\zeta_i)\| + \|f(\zeta_i) - f(s)\| \\ &< \varepsilon \quad (s \in F_i; i = 1, 2, \dots, m). \end{aligned}$$

By (1.2) and (1.3) we have

$$x_i = h(s) \in (f(\zeta_i) + B_{\delta(R)}(X)) \cap O \quad (s \in F_i; i = 1, 2, \dots, m)$$

and

$$g(s) \in (f(\zeta_i) + B_{\delta(R)}(X)) \cap O \quad (s \in F_i; i = 1, 2, \dots, m)$$

and it follows that for each $i = 1, 2, \dots, m$ both $h(F_i)$ and $g(F_i)$ are contained in the same connected component of $(f(\zeta_i) + B_R(X)) \cap O$. Now Lemma 4 applies to show that for each $i = 1, 2, \dots, m$ there exists a homotopy $p_i: F_i \times I \rightarrow O$ satisfying

$$\left. \begin{aligned} p_i(s, 0) &= g(s) \\ p_i(s, 1) &= h(s) \end{aligned} \right\} (s \in F_i; i = 1, 2, \dots, m)^*$$

and

$$(1.5) \quad \begin{aligned} p_i(s, t) &\in (f(\zeta_i) + B_R(X)) \cap O \\ &(s \in F_i; 0 \leq t \leq 1; i = 1, 2, \dots, m). \end{aligned}$$

Define $\pi: F \times I \rightarrow O$ by $\pi|_{F_i \times I} = p_i$ ($i = 1, 2, \dots, m$). Since F_i ($i = 1, 2, \dots, m$) are disjoint compact sets it follows that $F_i \times I$ ($i = 1, 2, \dots, m$) are disjoint compact subsets of $F \times I$ and consequently π is continuous. Clearly $\pi(s, 0) = g(s)$ ($s \in F$). By (1.1) and (1.5) we have

$$\begin{aligned} \|p_i(s, t) - f(s)\| &\leq \|p_i(s, t) - f(\zeta_i)\| + \|f(\zeta_i) - f(s)\| \\ &< 2R \quad (s \in F_i; 0 \leq t \leq 1; i = 1, 2, \dots, m) \end{aligned}$$

which implies (iii). Finally, by (1.4) we have $\|\pi(s, 1) - f(s)\| < \varepsilon$ ($s \in F$).

THEOREM 3. *Let O be an open subset of a complex Banach space X and let $F \subset \partial\Delta$ be a closed set of measure 0. Let $f: F \rightarrow \bar{O}$ be a continuous function and assume that O is locally connected at each point of $f(F)$. Then there exists a homotopy $p: F \times I \rightarrow \bar{O}$ satisfying*

- (i) $p(s, t) \in O$ ($s \in F; 0 \leq t < 1$)
- (ii) $p(s, 1) = f(s)$ ($s \in F$).

Proof. Observe first that $f(F)$ being compact O is ULC on $f(F)$ by Lemma 2. Let $\varepsilon \mapsto \delta(\varepsilon)$ be a modulus of ULC of O on $f(F)$. Choose a decreasing sequence $\{R_n\}$ of positive numbers converging to 0 and a strictly increasing sequence $\{t_n\}$ of nonnegative numbers converging to 1 where $t_1 = 0$.

Assume for a moment that there exists a continuous function $p: F \times (I - \{1\}) \rightarrow O$ satisfying

$$(2.1) \quad \|p(s, t) - f(s)\| < 2R_n \quad (s \in F; t_n \leq t < 1; n \in N).$$

Since $\{R_n\}$ converges to 0 we have $\lim_{t \rightarrow 1} p(s, t) = f(s)$ uniformly for $s \in F$. Since f is continuous on F it follows that the extension $p: F \times I \rightarrow \bar{O}$ of p defined on $F \times \{1\}$ by $p(s, 1) = f(s)$ ($s \in F$) is continuous on $F \times I$ and satisfies (i), (ii) above.

To prove the existence of p satisfying (2.1) we first prove that there exists a sequence of homotopies $\pi_n: F \times I \rightarrow O$ ($n \in N$) satisfying

$$(2.2) \quad \pi_{n+1}(s, 0) = \pi_n(s, 1) \quad (s \in F, n \in N)$$

$$(2.3) \quad \|\pi_n(s, 1) - f(s)\| < \delta(R_{n+1})/2 \quad (s \in F, n \in N)$$

and

$$(2.4) \quad \|\pi_n(s, t) - f(s)\| < 2R_n \quad (s \in F, 0 \leq t \leq 1, n \in N).$$

To define π_1 observe that $F = \bigcup_{i=1}^m F_i$ where F_i ($i = 1, 2, \dots, m$) are disjoint compact sets such that

$$(2.5) \quad \|f(\xi) - f(\eta)\| < \delta(R_1)/4 \quad (\xi, \eta \in F_i, i = 1, 2, \dots, m)$$

Choose $\zeta_i \in F_i$ ($i = 1, 2, \dots, m$). Since $f(F) \subset \bar{O}$ there exist $x_i \in \bar{O}$ ($i = 1, 2, \dots, m$) satisfying

$$(2.6) \quad \|x_i - f(\zeta_i)\| < \delta(R_1)/4 \quad (i = 1, 2, \dots, m).$$

Define $g(s) = x_i$ ($s \in F_i, i = 1, 2, \dots, m$). Then $g: F \rightarrow O$ is a continuous function which satisfies $\|f(s) - g(s)\| < \delta(R_1)/2$ ($s \in F$) by (2.5) and (2.6). By Lemma 5 applied to $f, g, R = R_1$ and $\varepsilon = \delta(R_2)/2$ there exists a homotopy $\pi_1: F \times I \rightarrow O$ satisfying $\|\pi_1(s, 1) - f(s)\| < \delta(R_2)/2$ ($s \in F$) and $\|\pi_1(s, t) - f(s)\| < 2R_1$ ($s \in F, 0 \leq t \leq 1$).

Assume that there exist π_n ($n = 1, 2, \dots, l$) satisfying (2.2) for $1 \leq n \leq l-1$ and (2.3), (2.4) for $1 \leq n \leq l$. By Lemma 5 applied to f , to $s \mapsto g(s) = \pi_l(s, 1)$ ($s \in F$), to $R = R_{l+1}$ and $\varepsilon = \delta(R_{l+2})/2$ there exists a homotopy $\pi_{l+1}: F \times I \rightarrow O$ satisfying (2.2) for $n = l$ and (2.3), (2.4) for $n = l+1$. Now the existence of $\{\pi_n: n \in N\}$ satisfying (2.2), (2.3) and (2.4) follows by induction.

Now define $p(s, t_n + t(t_{n+1} - t_n)) = \pi_n(s, t)$ ($s \in F, 0 \leq t \leq 1, n \in N$). It is easy to check that p has all the required properties.

LEMMA 6. *Let O be an open subset of a complex Banach space X . Let $B \subset \partial A$ be a relatively open set, $H \subset B$ a compact set of measure 0 and $G \subset B$ a relatively closed set of measure 0, disjoint from H . Suppose that $\pi: H \times I \rightarrow O$ is a continuous function*

satisfying

$$\text{diam } \{\pi(s, t): 0 \leq t \leq 1\} \leq M \quad (s \in H)$$

for some M . Let $U \subset \Delta \cup B$ be a neighbourhood of H and let $\varepsilon > 0$ be arbitrary. Assume that $h: \Delta \cup B \rightarrow O$ is a continuous function such that

- (a) $h(s) = \pi(s, 0)$ ($s \in H$)
- (b) there exists $\tau > 0$ such that

$$h(\Delta \cup B) + B_\tau(X) \subset O.$$

Then there exists $g \in H_b(\Delta, X)$ satisfying

- (i) $(h + g)(s) = \pi(s, 1)$ ($s \in H$)
- (ii) $g(s) = 0$ ($s \in G$)
- (iii) $\|g(z)\| < \varepsilon$ ($z \in (\Delta \cup B) - U$)
- (iv) $\|g(z)\| < M + \varepsilon$ ($z \in \Delta \cup B$)
- (v) there exists $\delta > 0$ such that

$$(h + g)(\Delta \cup B) + B_\delta(X) \subset O.$$

Proof. By the assumption $\pi: H \times I \rightarrow O$ is continuous hence $\pi(H \times I)$ is a compact set contained in O . Consequently by (b) there exists $\delta > 0$ which satisfies

$$(3.0) \quad \delta < \varepsilon/3$$

$$(3.1) \quad h(\Delta \cup B) + B_{2\delta}(X) \subset O$$

and

$$(3.2) \quad \pi(H \times I) + B_{5\delta}(X) \subset O.$$

By the properties of H

$$H = \bigcup_{j=1}^m H_j$$

where H_j ($j = 1, 2, \dots, m$) are disjoint compact sets such that

$$(3.3) \quad \begin{aligned} \|\pi(\xi, 0) - \pi(\eta, 0)\| &< \delta \\ \|\pi(\xi, 1) - \pi(\eta, 1)\| &< \delta \end{aligned} \quad (\xi, \eta \in F_j: j = 1, 2, \dots, m).$$

Since h is continuous on $\Delta \cup B$ there exist disjoint neighbourhoods $V_j \subset U$ of the sets H_j ($j = 1, 2, \dots, m$), respectively, such that

$$(3.4) \quad \|h(\xi) - h(\eta)\| < \delta \quad (\xi, \eta \in V_j: j = 1, 2, \dots, m).$$

Let $1 \leq j \leq m$. Choose $\xi_j \in H_j$ and consider the set

$$P_j = -\pi(\xi_j, 0) + \{\pi(\xi_j, t): 0 \leq t \leq 1\} + B_{2\delta}(X).$$

By the continuity of π , p_j is an open connected set in X containing the point 0. By (3.3) we have

$$\left. \begin{aligned} \pi(s, 0) &\in \pi(\xi_j, 0) + B_\delta(X) \\ \pi(s, 1) &\in \pi(\xi_j, 1) + B_\delta(X) \end{aligned} \right\} (s \in H_j)$$

and it follows that $\pi(s, 1) - \pi(s, 0) \in P_j$ ($s \in H_j$). Hence $s \mapsto \pi(s, 1) - \pi(s, 0)$ is a continuous function from H_j into P_j . By Lemma 3 there exists $g_j \in H_B(\Delta, X)$ satisfying

$$(3.5) \quad g_j(s) = \pi(s, 1) - \pi(s, 0) \quad (s \in H_j)$$

$$(3.6) \quad g_j(s) = 0 \quad (s \in G \cup (H - H_j))$$

$$(3.7) \quad \|g_j(z)\| < \delta/m \quad (z \in (\Delta \cup B) - V_j)$$

$$(3.8) \quad g_j(\Delta \cup B) \subset P_j.$$

Define

$$g = \sum_{j=1}^m g_j.$$

By (3.7) we have

$$(3.9) \quad \|g(z)\| < \delta \quad (z \in (\Delta \cup B) - \bigcup_{j=1}^m V_j).$$

Since $V_j \subset U$ ($j = 1, 2, \dots, m$) it follows that $\|g(z)\| < \delta$ ($z \in (\Delta \cup B) - U$) and by (3.0) (iii) is satisfied.

If $z \in V_j$ for some j , $1 \leq j \leq m$ then $z \notin V_k$ ($k \neq j$, $1 \leq k \leq m$). Consequently we have by (3.7) and (3.8)

$$(3.10) \quad \begin{aligned} g(z) &= g_j(z) + \sum_{k \neq j} g_k(z) \\ &\in P_j + B_\delta(X) \quad (z \in V_j, j = 1, 2, \dots, m) \end{aligned}$$

By the assumption $\text{diam } \{\pi(s, t): 0 \leq t \leq 1\} \leq M$ ($s \in H$) which implies that

$$\sup_{y \in P_j} \|y\| \leq M + 2\delta \quad (j = 1, 2, \dots, m).$$

By (3.0), (3.9) and (3.10) (iv) follows.

Now, (3.9) implies that

$$h(z) + g(z) \in h(\Delta \cup B) + B_\delta(X) \quad (z \in (\Delta \cup B) - \bigcup_{j=1}^m V_j)$$

and by (3.1) it follows that

$$(3.11) \quad h(z) + g(z) + B_\delta(X) \subset O \quad (z \in (\Delta \cup B) - \bigcup_{j=1}^m V_j).$$

If $z \in V_j$ for some j , $1 \leq j \leq m$ then $z \notin V_k$ ($k \neq j, 1 \leq k \leq m$) so by (a), by (3.4) and by (3.10) it follows that

$$\begin{aligned} h(z) + g(z) &\in h(\xi_j) + B_\delta(X) + P_j + B_\delta(X) \\ &= h(\xi_j) + B_\delta(X) - \pi(\xi_j, 0) + \{\pi(\xi_j, t): 0 \leq t \leq 1\} + B_{2\delta}(X) + B_\delta(X) \\ &\subset \pi(F \times I) + B_{4\delta}(X). \end{aligned}$$

Now (3.2) implies that $h(z) + g(z) + B_\delta(X) \subset O$ ($z \in \bigcup_{j=1}^m V_j$), which, together with (3.11) gives (v).

By the properties of g_j ($1 \leq j \leq m$) it is easy to see that (i) and (ii) are also satisfied.

THEOREM 4. *Let O be an open connected subset of a complex Banach space X . Let $B \subset \partial \Delta$ be a relatively open set and $F \subset B$ a relatively closed set of measure 0. Assume that $p: F \times I \rightarrow \bar{O}$ is a continuous function satisfying $p(s, t) \in O$ ($s \in F, 0 \leq t < 1$).*

There exists $f \in H_B(\Delta, X)$ such that

- (a) $f(s) = p(s, 1)$ ($s \in F$)
- (b) $f(z) \in O$ ($z \in (\Delta \cup B) - F$).

REMARK. In particular, Theorem 4 implies the following: Let O, B and F be as above and let $u: F \rightarrow \bar{O}$ be a continuous function. If there exists a peak continuous extension of u , i.e., a continuous extension $v: \Delta \cup B \rightarrow \bar{O}$ of u satisfying $v((\Delta \cup B) - F) \subset O$ then there exists a peak analytic extension of u , i.e. a continuous extension $w: \Delta \cup B \rightarrow \bar{O}$ of u , analytic on Δ and satisfying $w((\Delta \cup B) - F) \subset O$. To see this, put $p(s, t) = v(st)$ ($s \in F, 0 \leq t \leq 1$) and apply Theorem 4.

Proof of Theorem 4. We consider only the case when F is not compact. It is easy to see how to simplify the proof below in the case of compact F ; note that in the latter case the proof is considerably simpler.

Part 1. With no loss of generality we may assume that $0 \in O$. As in [8] write

$$F = \bigcup_{j=1}^{\infty} F_j$$

where $F_j \subset \Delta \cup B$ ($j \in N$) are compact sets infinitely many of which are not empty, such that there exist disjoint open sets $0_j \subset \Delta \cup B$ ($j \in N$) with the property that $F_j \subset 0_j$ ($j \in N$) and such that every compact subset of $\Delta \cup B$ misses all but a finite number of the sets 0_j . Passing to a subsequence if necessary we may assume that all

sets F_j ($j \in N$) are nonempty. Passing to a smaller 0_i if necessary we may assume that 0_i is contained in a compact subset of $\mathcal{A} \cup B$.

By Lemma 4 there exists for each $j \in N$ a homotopy $p_j: F \times I \rightarrow O$ satisfying $p_j(s, 0) = 0$, $p_j(s, 1) = p(s, 0)$ ($s \in F_j$). With no loss of generality we may assume that $p(s, 0) = 0$ ($s \in F$) (otherwise we replace p by π defined as follows

$$\begin{aligned}\pi(s, t) &= p_j(s, 2t) & (0 \leq t \leq 1/2, s \in F_j, j \in N) \\ \pi(s, t) &= p(s, 2t - 1) & (1/2 \leq t \leq 1, s \in F);\end{aligned}$$

by the properties of the sets F_j ($j \in N$) it is easy to see that $\pi: F \times I \rightarrow \bar{O}$ is continuous).

By the compactness of the sets F_j ($j \in N$) and by the continuity of p there exists for each $j \in N$ an increasing sequence $\{t_{ij}: i \in N\}$, $0 < t_{ij} < 1$ ($i \in N$), converging to 1 and satisfying

$$(4.1) \quad \text{diam} \{p(s, t): t_{ij} \leq t \leq 1\} < 1/2^{i+2} \quad (s \in F_j, j \in N, i \in N).$$

Part 2. In the sequel we will prove the following:

(A) for each $j \in N$ there exists a decreasing sequence $U_{ij} \subset \mathcal{A} \cup B$ ($i \in N$) of neighbourhoods of F_j contained in 0_j and satisfying

$$\bigcap_{i=1}^{\infty} U_{ij} = F_j \quad (j \in N)$$

(B) there exists a decreasing sequence $\{\varepsilon_i\}$ of positive numbers where

$$(4.2) \quad B_{\varepsilon_1}(X) \subset O$$

such that

(C) there exists a sequence $\{g_i\} \subset H_B(\mathcal{A}, X)$ satisfying

- (i) $(\sum_{m=1}^i g_m)(s) = p(s, t_{ij})$ ($s \in F_j, 1 \leq j \leq i, i \in N$)
 $g_i(s) = 0$ ($s \in F_j, j > i, j \in N, i \in N$)
- (ii) $\|g_i(z)\| < 1/2^i$ ($z \in (\mathcal{A} \cup B) - U_{ii}, i \in N$)
- (iii) $\|g_i(z)\| < \varepsilon_i/2^i$ ($z \in (\mathcal{A} \cup B) - \bigcup_{j=1}^i U_{ij}, i \in N$)
- (iv) $(\sum_{m=1}^i g_m)(\mathcal{A} \cup B) + \varepsilon_{i+1}(X) \subset O$ ($i \in N$).

Assume for a moment that we have proved (A), (B), (C) above.

Define

$$f(z) = \sum_{i=1}^{\infty} g_i(z) \quad (z \in \mathcal{A} \cup B).$$

We show that f has all the required properties.

Since each compact of $\mathcal{A} \cup B$ misses all but a finite number of the sets 0_j , (C) (ii) implies that the series converges uniformly on

each compact subset of $\mathcal{A} \cup B$. Consequently $f \in H_B(\mathcal{A}, X)$. By (C) (i) and by the continuity of p we have for all $j \in N$ and for all $s \in F_j$

$$f(s) = \lim_{n \rightarrow \infty} \sum_{i=1}^n g_i(s) = \lim_{\substack{n \rightarrow \infty \\ n > j}} \sum_{i=1}^n g_i(s) = \lim_{n \rightarrow \infty} p(s, t_{nj}) = p(s, 1).$$

Consequently $f(s) = p(s, 1)$ ($s \in F$) so (a) is satisfied.

To check (b), let $z \in \mathcal{A} \cup B - F$. Let first $z \in (\mathcal{A} \cup B) - \bigcup_{j=1}^{\infty} U_{1j}$. Now (C) (iii) implies that $\|f(z)\| < \varepsilon_1$ and by (4.2) it follows that $f(z) \in O$. Now, let $z \in U_{1j}$ for some $j \in N$. Then $z \notin U_{1k}$ ($k \neq j, k \in N$). Further, since $z \notin F_j$ it follows that there exists $i \in N$ such that $z \in U_{ij}$, $z \notin U_{i+1,j}$. Consequently $z \in (\mathcal{A} \cup B) - \bigcup_{j=1}^{\infty} U_{i+1,j}$, so that $z \in (\mathcal{A} \cup B) - \bigcup_{j=1}^{\infty} U_{kj}$ for every $k \in N$, $k \geq i+1$. By (C) (iii) it follows that

$$\left\| \sum_{k=i+1}^{\infty} g_k(z) \right\| < \varepsilon_{i+1}$$

which, by (C) (iv) implies that $f(z) \in O$.

Part 3. It remains to prove (A), (B), (C) above and we do this by induction.

First, choose $\varepsilon_1 > 0$ such that (4.2) holds and put $U_{11} = 0_1$. Choose a decreasing sequence $\{U_{i1}\}$ of neighbourhoods of F_1 in $\mathcal{A} \cup B$ contained in 0_1 and satisfying $\bigcap_{i=1}^{\infty} U_{i1} = F_1$. By Lemma 3 there exists $g_1 \in H_B(\mathcal{A}, X)$ satisfying $g_1(s) = p(s, t_{11})$ ($s \in F_1$), $g_1(s) = 0$ ($s \in F - F_1$), $g_1(\mathcal{A} \cup B) \subset O$ and

$$(4.3) \quad \|g_1(z)\| < \min\{1/2, \varepsilon_1/2\} \quad (z \in (\mathcal{A} \cup B) - U_{11}).$$

Now 0_1 is contained in a compact subset of $\mathcal{A} \cup B$. Consequently $g_1(U_{11})$ is contained in a compact set contained in O by the continuity of g_1 . It follows by (4.2) and (4.3) that there exists ε_2 , $0 < \varepsilon_2 < \varepsilon_1$ satisfying $g_1(\mathcal{A} \cup B) + B_{\varepsilon_2}(X) \subset O$. So we have proved the existence of a sequence $\{U_{ki}, k \in N\}$ $\varepsilon_1, \varepsilon_2 < \varepsilon_1$ and $g_1 \in H_B(\mathcal{A}, X)$ satisfying (A) and (C) (i)-(iv) for $i = 1$.

Assume that we have proved the existence of the sequences $\{U_{kj}, k \in N\}$ ($1 \leq j \leq n$), of a decreasing sequence $\{\varepsilon_k, 1 \leq k \leq n+1\}$ of positive numbers and of a sequence $\{g_i, 1 \leq i \leq n\} \subset H_B(\mathcal{A}, X)$ such that (A) is satisfied for $1 \leq j \leq n$ and (C) (i)-(iv) is satisfied for $1 \leq i \leq n$. We show below how to choose a sequence $\{U_{k, n+1}, k \in N\}$ to satisfy (A) for $j = n+1$ and then how to choose ε_{n+2} : $0 < \varepsilon_{n+2} < \varepsilon_{n+1}$ and $g_{n+1} \in H_B(\mathcal{A}, X)$ to satisfy (C) (i)-(iv) for $i = n+1$.

By the compactness of the set $p(F_{n+1} \times [0, t_{n+1, n+1}])$ there exist $\delta > 0$ and $\varepsilon > 0$ satisfying

$$(4.4) \quad p(F_{n+1} \times [0, t_{n+1, n+1}]) + B_{\varepsilon}(X) \subset O$$

$$(4.5) \quad 4\delta < \varepsilon_{n+1}$$

$$(4.6) \quad \delta + \varepsilon < 1/2^{n+1} - 1/2^{n+2}$$

$$(4.7) \quad \delta + \varepsilon < \varepsilon_{n+1}/2^{n+1}.$$

We choose $\{U_{k,n+1}, k \in N\}$ as follows: $\sum_{m=1}^n g_m$ is continuous on $\Delta \cup B$ and by (C) (i) $(\sum_{m=1}^n g_m)(s) = 0$ ($s \in F_{n+1}$). Consequently there exists a neighbourhood $U_{n+1,n+1} \subset O_{n+1}$ of F_{n+1} in $\Delta \cup B$ such that

$$(4.8) \quad \|(\sum_{m=1}^n g_m)(z)\| < \delta \quad (z \in U_{n+1,n+1}).$$

Now choose a decreasing sequence $U_{k,n+1} \subset O_{n+1}$ ($k \in N$) of neighbourhoods of F_{n+1} in $\Delta \cup B$ satisfying $U_{k,n+1} = U_{n+1,n+1}$ ($k \in N, k \leq n$) and $\bigcap_{k=1}^{\infty} U_{k,n+1} = F_{n+1}$, so that (A) is satisfied for $j = n+1$.

By Lemma 3 there exists $e \in H_B(\Delta, X)$ satisfying

$$(4.9) \quad e(s) = p(s, t_{n+1,n+1}) \quad (s \in F_{n+1})$$

$$(4.10) \quad e(s) = 0 \quad (s \in F_k, k \in N, k \neq n+1)$$

$$(4.11) \quad \|e(z)\| < \delta \quad (z \in (\Delta \cup B) - U_{n+1,n+1})$$

and

$$(4.12) \quad e(\Delta \cup B) \subset p(F_{n+1} \times [0, t_{n+1,n+1}]) + B_\delta(X)$$

since $p(F_{n+1} \times [0, t_{n+1,n+1}]) + B_\delta(X)$ is an open connected set which contains 0 and the set $\{p(s, t_{n+1,n+1}), s \in F_{n+1}\}$.

Consider the function $h = \sum_{m=1}^n g_m + e$. Since (C) (iv) holds for $i = n$ it follows by (4.5) that $(\sum_{m=1}^n g_m)(\Delta \cup B) + B_{4\delta}(X) \subset O$ which, by (4.11) implies that

$$(4.13) \quad h(z) + B_{2\delta}(X) \subset O \quad (z \in (\Delta \cup B) - U_{n+1,n+1}).$$

Now let $z \in U_{n+1,n+1}$. By (4.8) and (4.12) we have $h(z) \in p(F_{n+1} \times [0, t_{n+1,n+1}]) + B_{2\delta}(X)$ which, by (4.4) implies that $h(z) + B_{2\delta}(X) \subset O$ ($z \in U_{n+1,n+1}$) and by (4.13) it follows that

$$(4.14) \quad h(\Delta \cup B) + B_{2\delta}(X) \subset O.$$

Now, put

$$\begin{aligned} H &= \bigcup_{k=1}^{n+1} F_k, \quad G = F - H, \\ \pi(s, t) &= \begin{cases} p(s, t_{nj} + t(t_{n+1,j} - t_{nj})) & (s \in F_j, 1 \leq j \leq n) \\ p(s, t_{n+1,n+1}) & (s \in F_{n+1}), \end{cases} \\ U &= \bigcup_{k=1}^{n+1} U_{n+1,k}, \quad M = 1/2^{n+2}. \end{aligned}$$

Observe that by (C) (i), (4.9) and (4.10) $\pi(s, 0) = h(s)$ ($s \in H$). Note also that by (4.1) $\text{diam} \{\pi(s, t), 0 \leq t \leq 1\} < M$ ($s \in H$). Now Lemma 6 applies to show that there exists $g \in H_B(\Delta, X)$ with the following properties:

$$(4.15) \quad (h + g)(s) = \pi(s, 1) = p(s, t_{n+1, j}) \quad (s \in F_j, 1 \leq j \leq n + 1)$$

$$(4.16) \quad g(s) = 0 \quad (s \in F_j, j \in N, j \geq n + 2)$$

$$(4.17) \quad \|g(z)\| < \varepsilon \quad (z \in (\Delta \cup B) - \bigcup_{k=1}^{n+1} U_{n+1, k})$$

$$(4.18) \quad \|g(z)\| < 1/2^{n+2} + \varepsilon \quad (z \in \Delta \cup B),$$

and that there exists ε_{n+2} , $0 < \varepsilon_{n+2} < \varepsilon_{n+1}$ such that

$$(4.19) \quad (h + g)(\Delta \cup B) + B_{\varepsilon_{n+2}}(X) \subset O.$$

Define $g_{n+1} = e + g$. Clearly $g_{n+1} \in H_B(\Delta, X)$. By (4.15) and (4.16) it follows that (C) (i) is satisfied for $i = n + 1$. Further, by (4.19) (C) (iv) is satisfied for $i = n + 1$.

Let $z \in (\Delta \cup B) - U_{n+1, n+1}$. By (4.11), (4.18) and (4.6) we have $\|g_{n+1}(z)\| \leq \|e(z)\| + \|g(z)\| < \delta + \varepsilon + 1/2^{n+2} < 1/2^{n+1}$ which implies that (C) (ii) is satisfied for $i = n + 1$. Finally, let $z \in (\Delta \cup B) - \bigcup_{j=1}^{n+1} U_{n+1, j}$. By (4.11), (4.17) and (4.7) it follows that

$$\|g_{n+1}(z)\| < \delta + \varepsilon < \varepsilon_{n+1}/2^{n+1}$$

which implies that (C) (iii) is also satisfied for $i = n + 1$.

Proof of Theorem 1. Clearly (C) \Rightarrow (B) \Rightarrow (A). By Lemma 1, (A) \Rightarrow (D). To show that (D) implies (C) assume that $P \subset \overline{\text{Int } P}$ and that $\text{Int } P$ is connected and locally connected at every point of P . Let $B \subset \partial \Delta$ be a relatively open set, $F \subset B$ a relatively closed set of measure 0 and $f: F \rightarrow P$ a continuous function. In the case of non-compact F write $F = \bigcup_{j=1}^{\infty} F_j$ where F_j ($j \in N$) are nonempty compact subsets of $\Delta \cup B$ such that there exist disjoint open sets $0_j \subset \Delta \cup B$ ($j \in N$) satisfying $F_j \subset 0_j$ ($j \in N$) ([8], see also Part 1 of the proof of Theorem 4). For each $j \in N$, $f_j = f|F_j$ is a continuous function from F_j to P . Now Theorem 3 applies to show that by the properties of P there exists for each $j \in N$ a homotopy $p_j: F_j \times I \rightarrow P$ satisfying $p_j(s, t) \in \text{Int } P$ ($s \in F_j, 0 \leq t < 1$) and $p_j(s, 1) = f_j(s)$ ($s \in F_j$). Define $p: F \times I \rightarrow P$ by $p|F_j \times I = p_j$ ($j \in N$). By the properties of p_j and F_j ($j \in N$) p is continuous and satisfies $p(s, t) \in \text{Int } P$ ($s \in F, 0 \leq t < 1$) and $p(s, 1) = f(s)$ ($s \in F$). In the case when F is compact the existence of such a p is immediate by Theorem 3. Now by Theorem 4 there exists $\tilde{f} \in H_B(\Delta, X)$ satisfying $\tilde{f}|F = f$ and $\tilde{f}((\Delta \cup B) - F) \subset \text{Int } P$.

Proof of Theorem 2. Clearly $(C) \Rightarrow (B) \Rightarrow (A)$. It remains to prove that (A) implies (C) . Assume that there exists a closed set $F \subset \partial\Delta$ of measure 0 with infinitely many points such that every continuous function $f: F \rightarrow P$ admits an extension $\tilde{f} \in A(\Delta)$ satisfying $\tilde{f}(\bar{\Delta}) \subset P$.

Let $f: F \rightarrow P$ be a nonconstant continuous function. By the above assumption there exists a necessarily nonconstant extension $\tilde{f} \in A(\Delta)$ of f satisfying $\tilde{f}(\bar{\Delta}) \subset P$. Since every nonconstant complex-valued analytic function is an open map [20] we have $\tilde{f}(\Delta) \subset \text{Int } P$. Consequently $(s, t) \mapsto p(s, t) = f(st)$ is a continuous function from $F \times I$ to P satisfying

$$(5.1) \quad \begin{aligned} p(s, t) &\in \text{Int } P & (s \in F, 0 \leq t < 1) \\ p(s, 1) &= f(s) & (s \in F) . \end{aligned}$$

In the case when $f: F \rightarrow P$ is a constant, say $f(s) = x$ ($s \in F$) write $F = F_1 \cup F_2$ where F_1, F_2 are disjoint compact sets. By the assumption P consists of more than one point so that there is a $y \in P$, $y \neq x$. Further, by the above assumptions there exist necessarily nonconstant functions $f_1, f_2 \in A(\Delta)$ satisfying $f_1(F_1) = f_2(F_2) = \{x\}$, $f_1(F_2) = f_2(F_1) = \{y\}$ and $f_1(\bar{\Delta}) \subset P$, $f_2(\bar{\Delta}) \subset P$. Now define $p: F \times I \rightarrow P$ as follows

$$p(s, t) = \begin{cases} f_1(st) & (s \in F_1, 0 \leq t \leq 1) \\ f_2(st) & (s \in F_2, 0 \leq t \leq 1) . \end{cases}$$

Since f_1 and f_2 are open maps p is a continuous function satisfying (5.1).

Having proved the existence of a continuous function $p: F \times I \rightarrow P$ satisfying (5.1) Theorem 4 applies to show that there exists $g \in A(\Delta, X)$ satisfying $g|F = f$ and $g(\bar{\Delta} - F) \subset \text{Int } P$. Since f was arbitrary it follows that (A) in Theorem 1 is satisfied for our F and P . Consequently (C) follows by Theorem 1.

3. Application and remarks. Given any set $P \subset C$ homeomorphic to $\bar{\Delta}$ there exists $f \in A(\Delta)$ satisfying $f(\bar{\Delta}) = f(\partial\Delta) = P$. This is a consequence of Rudin's theorem. Below we present some generalizations of this result. The existence of $f \in A(\Delta)$ such that $f(\partial\Delta)$ fills some square had been proved before Rudin's theorem and had raised certain interest [16]. Church [5] gave a complete topological description of the sets $f(\partial\Delta)$, $f \in A(\Delta)$.

APPLICATION 1. Given any nonempty compact set $P \subset C$ satisfying $P = \overline{\text{Int } P}$ and such that $\text{Int } P$ is connected and locally connected at every point of P , there exists $\tilde{f} \in A(\Delta)$ satisfying $\tilde{f}(\bar{\Delta}) = \tilde{f}(\partial\Delta) = P$.

Proof. Let $F \subset \partial\Delta$ be a Cantor set of measure 0. By the compactness of P there exists $f \in C(F)$ satisfying $f(F) = P$ [13]. Now by Corollary 1 there exists an extension $\tilde{f} \in A(\Delta)$ satisfying $\tilde{f}(\bar{\Delta}) \subset P$.

APPLICATION 2. Given any nonempty set $P \subset C$ satisfying $P = \overline{\text{Int } P}$ and such that $\text{Int } P$ is connected and locally connected at every point of P , there exists a continuous function $\tilde{f}: \bar{\Delta} - \{1\} \rightarrow C$, analytic on Δ and satisfying

$$\tilde{f}((\bar{\Delta} - \{1\}) \cap U) = \tilde{f}((\partial\Delta - \{1\}) \cap U) = P$$

for every neighbourhood $U \subset C$ of the point 1.

Proof. Let $\{V_n: n \in N\}$ be a sequence of disjoint open arcs in $\partial\Delta$ such that for every neighbourhood U of the point 1 there exists $n_U \in N$ such that $V_n \subset U$ ($n \in N, n > n_U$). For each $n \in N$ let $F_n \subset V_n$ be a Cantor set of measure 0. It is easy to construct a function $\alpha: N \rightarrow N$ satisfying

$$N \subset \alpha(\{n, n+1, \dots\}) \quad (n \in N).$$

Write $P = \bigcup_{n=1}^{\infty} P_n$ where P_n are compact sets. For each $n \in N$ there exists $f_n \in C(F_n)$ satisfying $f_n(F_n) = P_{\alpha(n)}$ [13]. Put $F = \bigcup_{n=1}^{\infty} F_n$ and define $f: F \rightarrow P$ by $f|F_n = f_n$ ($n \in N$). Clearly f is a continuous function. By Theorem 1 there exists a continuous extension $\tilde{f}: \bar{\Delta} - \{1\} \rightarrow X$, analytic on Δ and satisfying $\tilde{f}(\bar{\Delta} - \{1\}) \subset P$. It is easy to see that \tilde{f} has all the required properties.

The following application to vector-valued functions is perhaps more interesting. Its proof is the same as the proof of Application 1.

APPLICATION 3. Given any nonempty compact set P in a finite dimensional complex normed space X satisfying $P = \overline{\text{Int } P}$ and such that $\text{Int } P$ is connected and locally connected at every point of P there exists $\tilde{f} \in A(\Delta, X)$ such that $\tilde{f}(\bar{\Delta}) = \tilde{f}(\partial\Delta) = P$.

REMARK. Note that in Application 2 we can replace C by any finite-dimensional complex normed space.

Note that in the case when X is infinite-dimensional one can not fill a subset of X having nonempty interior with $f(\Delta \cup B)$ for some $f \in H_B(\Delta, X)$ since $f(\Delta \cup B)$ is always a countable union of compact subsets of X and consequently its interior is empty. For the results about cluster sets in this case see [8, 9].

REMARK. If $X = C$ then by Theorem 2 a set $P \subset X$ containing more than one point has AEP if and only if it has PAEP. If $\dim X \geq 2$ this is no longer true since there exist subsets P of X having AEP whose interior is empty. An example is $P = \overline{I}x$ where $x \in X$, $x \neq 0$.

PROBLEM. Give an example of a nonempty set $P \subset C^2$ satisfying $P = \overline{\text{Int } P}$ which has AEP but not PAEP.

REMARK. The results of the present paper give a partial solution to Problem 3 in [10].

ACKNOWLEDGMENT. The author wishes to express his thanks to Professor Ivan Vidav and to professor Jože Vrabec for several very stimulating discussions while this paper was being prepared. A part of the results was obtained in November 1975 while the author was visiting Trinity College, Dublin. He wishes to express his thanks to Professor Trevor West and to Professor Richard Aron for organizing the visit and specially to Professor Aron for his hospitality.

REFERENCES

1. E. Bishop, *A general Rudin-Carleson theorem*, Proc. Amer. Math. Soc., **13** (1962), 140-143.
2. L. Brown, P. M. Gauthier and W. Hengartner, *Continuous boundary behaviour for functions defined in the open unit disc*, Nagoya Math. J., **57** (1975), 49-58.
3. L. Carleson, *Representations of continuous functions*, Math. Z., **66** (1957), 447-451.
4. P. T. Church, *Global boundary behavior of meromorphic functions*, Acta Math., **105** (1961), 49-62.
5. ———, *Boundary images of meromorphic functions*, Trans. Amer. Math. Soc., **110** (1964), 52-78.
6. E. F. Collingwood and A. J. Lohwater, *The theory of cluster sets*, Cambridge Tracts Math. Math. Phys. No. **56**. Cambridge University Press, (1966).
7. J. Globevnik, *The Rudin-Carleson theorem for vector-valued functions*, Proc. Amer. Math. Soc., **53** (1975), 250-252.
8. ———, *Analytic functions whose range is dense in a ball*, J. Functional Analysis, **22** (1976), 32-38.
9. ———, *The range of vector-valued analytic functions*, Ark. Mat., **14** (1976), 113-118. *The range of vector-valued analytic functions II*, Ark. Mat.; **14** (1976), 297-298.
10. ———, *On the range of analytic functions into a Banach space*, To appear in Proc. Symp. Inf. Dim. Holomorphy, Campinas, (1975), Notas de Mat., North Holland.
11. ———, *Analytic extensions of vector-valued functions*, Pacific J. Math., **63** (1976), 389-395.
12. E. A. Heard and J. H. Wells, *An interpolation problem for subalgebras of H^∞* , Pacific J. Math., **28** (1969), 543-553.
13. J. G. Hocking and G. S. Young, *Topology*, Addison-Wessley, 1961.
14. K. Hoffman, *Banach Spaces of Analytic Functions*, Prentice-Hall, 1962.
15. G. M. Leibowitz, *Lectures on Complex Function Algebras*, Scott, Foresman, 1970.

16. G. R. MacLane, *On the Peano curve associated with some conformal maps*, Proc. Amer. Math. Soc., **6** (1955), 625-630.
17. M. H. A. Newman, *Elements of the Topology of Plane Sets of Points*, Cambridge Univ. Press, 1964.
18. D. M. Oberlin, *Interpolation and vector-valued functions*, J. Functional Analysis **15** (1974), 428-439.
19. W. Rudin, *Boundary values of continuous analytic functions*, Proc. Amer. Math. Soc., **7** (1956), 808-811.
20. ———, *Real and Complex Analysis*, McGraw-Hill, 1966.
21. E. L. Stout, *On Some Restriction Algebras*, Function Algebras, Ed. F. T. Birtel, Scott, Foresman, 1966, pp. 6-11.
22. ———, *The Theory of Uniform Algebras*, Bogden and Quigley, 1971.
23. G. T. Whyburn, *Topological Analysis*, Princeton University Press, 1964.

Received July 1, 1976 and in revised form October 27, 1976. This work was supported in part by the Boris Kidrič Fund, Ljubljana, Yugoslavia

UNIVERSITY OF LJUBLJANA
LJUBLJANA, YUGOSLAVIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, CA 90024

R. A. BEAUMONT

University of Washington
Seattle, WA 98105

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 69, No. 2

June, 1977

Carol Alf and Thomas Alfonso O'Connor, <i>Unimodality of the Lévy spectral function</i>	285
S. J. Bernau and Howard E. Lacey, <i>Bicontractive projections and reordering of L_p-spaces</i>	291
Andrew J. Berner, <i>Products of compact spaces with bi-k and related spaces</i>	303
Stephen Richard Bernfeld, <i>The extendability and uniqueness of solutions of ordinary differential equations</i>	307
Marilyn Breen, <i>Decompositions for nonclosed planar m-convex sets</i>	317
Robert F. Brown, <i>Cohomology of homomorphisms of Lie algebras and Lie groups</i>	325
Jack Douglas Bryant and Thomas Francis McCabe, <i>A note on Edelstein's iterative test and spaces of continuous functions</i>	333
Victor P. Camillo, <i>Modules whose quotients have finite Goldie dimension</i>	337
David Downing and William A. Kirk, <i>A generalization of Caristi's theorem with applications to nonlinear mapping theory</i>	339
Daniel Reuven Farkas and Robert L. Snider, <i>Noetherian fixed rings</i>	347
Alessandro Figà-Talamanca, <i>Positive definite functions which vanish at infinity</i>	355
Josip Globevnik, <i>The range of analytic extensions</i>	365
André Goldman, <i>Mesures cylindriques, mesures vectorielles et questions de concentration cylindrique</i>	385
Richard Grassl, <i>Multisectioned partitions of integers</i>	415
Haruo Kitahara and Shinsuke Yorozu, <i>A formula for the normal part of the Laplace-Beltrami operator on the foliated manifold</i>	425
Marvin J. Kohn, <i>Summability R_r for double series</i>	433
Charles Philip Lanski, <i>Lie ideals and derivations in rings with involution</i>	449
Solomon Leader, <i>A topological characterization of Banach contractions</i>	461
Daniel Francis Xavier O'Reilly, <i>Cobordism classes of fiber bundles</i>	467
James William Pendergrass, <i>The Schur subgroup of the Brauer group</i>	477
Howard Lewis Penn, <i>Inner-outer factorization of functions whose Fourier series vanish off a semigroup</i>	501
William T. Reid, <i>Some results on the Floquet theory for disconjugate definite Hamiltonian systems</i>	505
Caroll Vernon Riecke, <i>Complementation in the lattice of convergence structures</i>	517
Louis Halle Rowen, <i>Classes of rings torsion-free over their centers</i>	527
Manda Butchi Suryanarayana, <i>A Sobolev space and a Darboux problem</i>	535
Charles Thomas Tucker, II, <i>Riesz homomorphisms and positive linear maps</i>	551
William W. Williams, <i>Semigroups with identity on Peano continua</i>	557
Yukinobu Yajima, <i>On spaces which have a closure-preserving cover by finite sets</i>	571