

Pacific Journal of Mathematics

**INNER-OUTER FACTORIZATION OF FUNCTIONS WHOSE
FOURIER SERIES VANISH OFF A SEMIGROUP**

HOWARD LEWIS PENN

INNER-OUTER FACTORIZATION OF FUNCTIONS WHOSE FOURIER SERIES VANISH OFF A SEMIGROUP

HOWARD LEWIS PENN

Let G be a compact, connected, Abelian group. Its dual, Γ , is discrete and can be ordered. Let Γ_1 be a semigroup which is a subset of the positive elements for some ordering, but which contains the origin of Γ . Let $H^p(\Gamma_1)$ be the subspace of $L^p(G)$ consisting of functions which have vanishing off Γ_1 . The question that this paper is concerned with is what conditions on a function in $H^p(\Gamma_1)$ assure an inner-outer factorization.

An inner function is a function $f \in H^\infty(\Gamma_1)$ such that $|f|=1$ a.e. (dx) on G . A function $f \in H^p(\Gamma_1)$ is said to be outer if

$$\int_G \log |f(x)| = \log \left| \int_G f(x) dx \right| > -\infty.$$

A function $f \in H^1(\Gamma_1)$ is said to be in the class $LRP(\Gamma_1)$ if $\log |f| \in L^1(G)$ and $\log |f|$ has Fourier coefficients equal to zero off $\Gamma_1 \cup -\Gamma_1$. The main result of the paper is that if Γ_1 is the intersection of half planes and $f \in H^1(\Gamma_1)$ with $\int_G \log |f(x)| dx > -\infty$ then f has an inner-outer factorization if and only if $\log |f|$ is in $LRP(\Gamma_1)$.

A semigroup, P , in Γ_1 is called a half plane if $P \cup -P = \Gamma$ and $P \cap -P = \{0\}$. Helson and Lowdenslager [2] proved that if Γ_1 is a half plane then every function $f \in H^p(\Gamma_1)$ with $\int \log |f| dx > -\infty$ has a factorization as a product of an outer function, $h \in H^p(\Gamma_1)$ and an inner function, g , and this factorization is unique up to multiplication by constants of magnitude 1. From now on we shall assume $\int \log |f| dx > -\infty$.

Helson and Lowdenslager also showed [3] that if u is a real function such that u and e^u are summable, and v is the conjugate function of u with respect to the half plane, Γ_1 , then e^{u+iv} is an outer function in $H^1(\Gamma_1)$. Conversely, if a summable outer function has the representation e^{u+iv} with u and v real then u is summable and v is equal to its conjugate modulo 2π except for an additive constant.

Let P be a half plane which contains Γ_1 . Then, for $u \in L^1_r(G)$ there exists a conjugate function, v , which is unique if we assume $v(0) = 0$, such that $u + iv$ has its Fourier series supported on P . The function, v , is in L^p , $p < 1$. If u has its Fourier coefficients supported only on $\Gamma_1 \cup -\Gamma_1$ then $u + iv$ has its Fourier coefficients supported only on Γ_1 [4, Chap. 8, §7]. Therefore, $f \in LRP(\Gamma_1)$ if

and only if $\log |f| \in L^1$ and $\log |f|$ is the real part of a function whose Fourier coefficients vanish off Γ_1 .

THEOREM. *Assuming that $f \in H^1(\Gamma_1)$ and $\int \log |f| dx > -\infty$, and that $\Gamma_i = \bigcap_{i \in I} P_i$, then f has an inner-outer factorization if and only if $\log |f|$ has its Fourier coefficients vanish off $\Gamma_1 \cup -\Gamma_1$.*

Proof. Assume $f \in LRP(\Gamma_1)$. Let $u = \log |f| \in L^1(G)$. Take any $i \in I$ and consider P_i . $\Gamma_1 \subset P_i$ and f has an inner-outer factorization with respect to P_i . The outer factor is given by e^{u+iv_i} where v_i is the conjugate function to u with respect to P_i . Since u has its Fourier coefficients supported on $\Gamma_1 \cup -\Gamma_1$, it follows that v_i also has its Fourier coefficients supported there. Therefore, v_i is the same as the conjugate function of u with respect to any of the other half planes $P_j, j \in I$. Therefore, the outer factor of f in $H^1(P_i)$ is given by e^{u+iv_i} . Also, if P_j is any of the other half planes whose intersection gives Γ_1 , then the outer factor f in $H^1(P_j)$ is e^{u+iv_j} . Therefore, $e^{u+iv_i} \in \bigcap_{i \in I} H^1(P_i)$, which is just equal to $H^1(\Gamma_1)$. For each half plane $P_i, i \in I$, we have that the inner factor is given by $f e^{-(u+iv_i)}$. Therefore, the inner factor is also in $H^1(\Gamma_1)$.

Conversely, assume that f has an inner-outer factorization, gh , in $H^1(\Gamma_1)$. Choose $P_j, j \in I$, then the outer factor, h , of f in $H^1(\Gamma_1)$, and hence in $H^1(P_j)$, is given by e^{u+v_j} , where v_j is the conjugate function of $u = \log |f|$ with respect to P_j . Since this is true for all $P_j, j \in I$, it follows that e^{iv_j} is the same regardless of which half plane, P_j is used. Now assume P_k is another of the half planes whose intersection is Γ_1 . Then $e^{iv_k} = e^{iv_j}$ where v_k is the conjugate function of u with respect to P_j . It follows that $v_k(x) = v_j(x) + 2n\pi$ where n might change from point to point. We will now show that $n = 0$. Consider the function $h^{1/2}$ which is outer in $H^2(\Gamma_1) \subset H^1(\Gamma_1)$. It follows that $\log |h^{1/2}| = u/2$. The conjugate function of $u/2$ with respect to P_j is $v_j/2$ and its conjugate function with respect to P_k is $v_k/2$. By the Helson and Lowdenslager theorem $h^{1/2} = e^{(u+iv_j)/2}$ and also $h^{1/2} = e^{(u+iv_k)/2}$. Therefore,

$$h = h^{1/2} h^{1/2} = e^{u+i(v_j+v_k)/2}.$$

Hence

$$v_k(x) = (v_j(x) + v_k(x))/2 + 2n\pi.$$

So,

$$v_k(x) = v_j(x) + 4n\pi.$$

Now consider $h^{1/4} = e^{(u+iv_j)/4} = e^{(u+iv_k)/4}$. Therefore,

$$h = h^{3/4}h^{1/4} = e^{3(u+iv_k)/4}e^{(u+iv_j)/4} = e^{u+i(3v_k+v_j)/4}.$$

Hence,

$$v_k(x) = v_j(x) + 8n\pi.$$

By considering the 2^m th roots of h we can show that the difference between v_k and v_j must be $2^{m+1}n\pi$. This must hold for all values of m . The only integer for which this is true is 0. Therefore u has the same conjugate function with respect to each of the half planes.

We will show that u has its Fourier coefficients supported on $\Gamma_1 \cup -\Gamma_1$. Suppose that $\hat{u}(\gamma) \neq 0$, where $\gamma \notin \Gamma_1 \cup -\Gamma_1$. Then there exists $P_j, j \in I$ such that $\gamma \notin P_j$. There also exists $P_k, k \in I$, such that $\gamma \notin -P_k$. Let v_j be the conjugate functions of u with respect to the half plane, P_j and let v_k be the conjugate function of u with respect to P_k . Since $\gamma \notin P_j$, we have

$$\hat{v}_j(\gamma) = i\hat{u}(\gamma)$$

[4, Chap. 8, §7]. Likewise, since $\gamma \notin -P_k$ it follows that $\gamma \in P_k \setminus 0$ and that

$$\hat{v}_k(\gamma) = -i\hat{u}(\gamma).$$

But since $\hat{u}(\gamma) \neq 0$, we have $\hat{v}_j(\gamma) \neq \hat{v}_k(\gamma)$, and hence v_j and v_k are different functions. But we have just shown that u has the same conjugate function with respect to each half plane. Therefore $\hat{u}(\gamma) = 0$ and u has its Fourier series supported on $\Gamma_1 \cup -\Gamma_1$. Therefore $f \in LRP(\Gamma_1)$.

COROLLARY. *If $f \in H^1(\Gamma_1)$ where Γ_1 is the intersection of half planes and $f \in LRP(\Gamma_1)$, then $f = p_1 p_2$ where $p_1, p_2 \in H^2(\Gamma_1)$ and $|p_1|^2 \equiv |p_2|^2 \equiv |f|$*

EXAMPLE. In [1] Ebenstein discusses the H^p functions on a semigroup which is the intersection of a countable collection of half planes. This semigroup fulfills the hypothesis of the theorem. Let T^ω be the compact group which is the Cartesian product of countably many circles. The dual $\sum_{i=1}^\infty Z$, is the direct sum of countably many copies of the integers. Define $A \subset \sum_{i=1}^\infty Z$ by

$$A = \{x: x_i \geq 0 \text{ for all } i\}.$$

We may define $H^p(T^\omega)$, $p \geq 1$ as the subset of $L^p(T^\omega)$ consisting of these functions whose Fourier coefficients vanish off A . The semigroup, A , is the intersection of half planes P_i defined as follows:

$$P_i = \{x: x_i \geq 0, \text{ if } x_i = 0 \text{ then } x_1 \geq 0, \\ \text{if } x_1, x_2, \dots, x_j = 0 \text{ then } x_{j+1} = 0\}.$$

Therefore the theorem applies to $H^p(T^\omega)$.

REMARK. One might hope that certain theorems which hold for the H^p spaces of the disk would remain true, at least for the class $LRP(\Gamma_1)$. One such theorem is Szego's theorem which states if $w \in L^1(dx)$ and $w \geq 0$, then

$$\inf_{g \in A_0} \int |1 - g|^2 w dx = \exp \int \log(w) dx$$

where A_0 consists of those polynomials supported on Γ_1 , with zero-th coefficient equal to zero. This theorem is true if Γ_1 is a half plane [4, Chap. 8, §3]. Rudin has an example [5, Theorem 4.4.8] of a function, f , which is outer, but does not span. This same function can be used to show that Szego's theorem fails even for the class LRP .

REFERENCES

1. Samuel E. Ebenstein, *Some H^p spaces which are uncomplimented in L^p* , Pacific J. Math., **43** (1972), 327-339.
2. H. Helson and D. Lowdenslager, *Prediction theory and Fourier analysis in several variables I*, Acta. Math., **99** (1958), 105-202.
3. ———, *Prediction theory and Fourier analysis in several variables II*, Acta. Math., **106** (1961), 175-213.
4. Walter Rudin, *Fourier Analysis on Groups*, Interscience, New York, (1962).
5. ———, *Function Theory in Polydiscs*, Benjamin, New York, (1969).

Received November 19, 1974 and in revised form November 8, 1976. The work in this paper was completed as part of the author's doctoral thesis at the University of Michigan.

UNITED STATES NAVAL ACADEMY
ANNAPOLIS, MD 21402

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, CA 90024

R. A. BEAUMONT

University of Washington
Seattle, WA 98105

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).

8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 69, No. 2

June, 1977

Carol Alf and Thomas Alfonso O'Connor, <i>Unimodality of the Lévy spectral function</i>	285
S. J. Bernau and Howard E. Lacey, <i>Bicontractive projections and reordering of L_p-spaces</i>	291
Andrew J. Berner, <i>Products of compact spaces with bi-k and related spaces</i>	303
Stephen Richard Bernfeld, <i>The extendability and uniqueness of solutions of ordinary differential equations</i>	307
Marilyn Breen, <i>Decompositions for nonclosed planar m-convex sets</i>	317
Robert F. Brown, <i>Cohomology of homomorphisms of Lie algebras and Lie groups</i>	325
Jack Douglas Bryant and Thomas Francis McCabe, <i>A note on Edelstein's iterative test and spaces of continuous functions</i>	333
Victor P. Camillo, <i>Modules whose quotients have finite Goldie dimension</i>	337
David Downing and William A. Kirk, <i>A generalization of Caristi's theorem with applications to nonlinear mapping theory</i>	339
Daniel Reuven Farkas and Robert L. Snider, <i>Noetherian fixed rings</i>	347
Alessandro Figà-Talamanca, <i>Positive definite functions which vanish at infinity</i>	355
Josip Globevnik, <i>The range of analytic extensions</i>	365
André Goldman, <i>Mesures cylindriques, mesures vectorielles et questions de concentration cylindrique</i>	385
Richard Grassl, <i>Multisectioned partitions of integers</i>	415
Haruo Kitahara and Shinsuke Yorozu, <i>A formula for the normal part of the Laplace-Beltrami operator on the foliated manifold</i>	425
Marvin J. Kohn, <i>Summability R_r for double series</i>	433
Charles Philip Lanski, <i>Lie ideals and derivations in rings with involution</i>	449
Solomon Leader, <i>A topological characterization of Banach contractions</i>	461
Daniel Francis Xavier O'Reilly, <i>Cobordism classes of fiber bundles</i>	467
James William Pendergrass, <i>The Schur subgroup of the Brauer group</i>	477
Howard Lewis Penn, <i>Inner-outer factorization of functions whose Fourier series vanish off a semigroup</i>	501
William T. Reid, <i>Some results on the Floquet theory for disconjugate definite Hamiltonian systems</i>	505
Caroll Vernon Riecke, <i>Complementation in the lattice of convergence structures</i>	517
Louis Halle Rowen, <i>Classes of rings torsion-free over their centers</i>	527
Manda Butchi Suryanarayana, <i>A Sobolev space and a Darboux problem</i>	535
Charles Thomas Tucker, II, <i>Riesz homomorphisms and positive linear maps</i>	551
William W. Williams, <i>Semigroups with identity on Peano continua</i>	557
Yukinobu Yajima, <i>On spaces which have a closure-preserving cover by finite sets</i>	571