

Pacific Journal of Mathematics

ENUMERATION OF DOUBLY UP-DOWN PERMUTATIONS

L. CARLITZ

ENUMERATION OF DOUBLY UP-DOWN PERMUTATIONS

L. CARLITZ

It is well known that $A(n)$, the number of up-down permutations of $\{1, 2, \dots, n\}$ satisfies

$$\sum_{n=0}^{\infty} A(2n) \frac{z^{2n}}{(2n)!} = \sec z,$$

$$\sum_{n=0}^{\infty} A(2n+1) \frac{z^{2n+1}}{(2n+1)!} = \tan z.$$

In the present paper generating functions are obtained for up-down (down-up) permutations in which the peaks themselves are in an up-down configuration.

In a previous paper the writer obtained generating functions for the number of up-down (and down-up) permutations counting the rises among the "peaks".

1. Let $Z_n = \{1, 2, \dots, n\}$ and let (a_1, a_2, \dots, a_n) be an arbitrary [4, pp. 105–112] up-down permutation of Z_n . Then (b_1, b_2, \dots, b_n) , where

$$b_i = n - a_i + 1 \quad (i = 1, 2, \dots, n)$$

is a down-up permutation and vice versa. Thus, for $n > 1$, there is a one-to-one correspondence between up-down and down-up permutations so that it suffices to consider the former.

In the present paper we are concerned with up-down (and down-up) permutations of Z_n in which it is required that the peaks themselves satisfy the up-down or down-up conditions. Thus let (a_1, a_2, \dots, a_n) denote an up-down permutation of Z_n so that

$$(1.1) \quad a_{2k-1} < a_{2k}, \quad a_{2k} > a_{2k+1} \quad (k = 1, 2, \dots, [n/2]).$$


Then the additional requirement is either

$$(1.2) \quad a_{4k-2} < a_{4k}, \quad a_{4k} > a_{4k+2} \quad (k = 1, 2, \dots, [n/4])$$

or

$$(1.3) \quad a_{4k-2} > a_{4k}, \quad a_{4k} < a_{4k+2} \quad (k = 1, 2, \dots, [n/4]).$$

For example the permutation $(1, 3, 2, 8, 4, 7, 6, 13, 9, 10, 5, 12, 11)$



satisfies both (1.1) and (1.2).

Next let (a_1, a_2, \dots, a_n) be a down-up permutation of Z_n so that

$$(1.4) \quad a_{2k-1} > a_{2k}, \quad a_{2k} < a_{2k+1} \quad (k = 1, 2, \dots, [n/2]).$$

Then the additional requirement is either

$$(1.2)' \quad a_{4k-3} > a_{4k-1}, \quad a_{4k-1} < a_{4k+1} \quad (k = 1, 2, \dots, [n/4])$$

or

$$(1.3)' \quad a_{4k-3} < a_{4k-1}, \quad a_{4k-1} < a_{4k+1} \quad (k = 1, 2, \dots, [n/4]).$$

Thus there are four possibilities, namely

- I. (1.1) and (1.2),
- II. (1.1) and (1.3),
- III. (1.4) and (1.2)',
- IV. (1.4) and (1.3)'.

There are various relations between these varieties of permutations; however they depend upon the residue of $n \pmod 4$.

In order to derive generating functions it will be convenient to define the following enumerants. Let $A_{RF}(n)$ denote the number of up-down permutations of Z_n such that the peaks

$$(a_2, a_4, \dots, a_{2[n/2]})$$

begin with a rise, $a_2 < a_4$, and end with a fall. Thus for this case it is necessary that $n \equiv 2$ or $3 \pmod 4$.

We define $A_{RR}(n)$, $A_{FR}(n)$, $A_{FF}(n)$ in a similar way. Note that for RR , $n \equiv 0$ or $1 \pmod 4$; for FR , $n \equiv 2$ or $3 \pmod 4$; for FF , $n \equiv 0$ or $1 \pmod 4$.

We also define $C_{RF}(n)$, $C_{RR}(n)$, $C_{FR}(n)$, $C_{FF}(n)$ in an analogous manner for down-up permutations. Then for RF , $n \equiv 1$ or $2 \pmod 4$, for RR , $n \equiv 0$ or $3 \pmod 4$, for FR , $n \equiv 1$ or $2 \pmod 4$, for FF , $n \equiv 0$ or $3 \pmod 4$.

We shall accordingly consider the following enumerants:

$$(1.5) \quad \left\{ \begin{array}{l} A_{RF}(4n+3), \quad A_{RF}(4n+2) \\ A_{RR}(4n+1), \quad A_{RR}(4n) \\ A_{FR}(4n+3), \quad A_{FR}(4n+2) \\ A_{FF}(4n+1), \quad A_{FF}(4n) \end{array} \right.$$

$$(1.6) \quad \left\{ \begin{array}{l} C_{RF}(4n+1), \quad C_{RF}(4n+2) \\ C_{RR}(4n+3), \quad C_{RR}(4n) \\ C_{FR}(4n+1), \quad C_{FR}(4n+2) \\ C_{FF}(4n+3), \quad C_{FF}(4n). \end{array} \right.$$

Reading a permutation both from left to right and from right to left, we get the following relations connecting the enumerants.

$$(1.7) \quad \left\{ \begin{array}{l} A_{RR}(4n+1) = A_{FF}(4n+1) \\ A_{RR}(4n) = C_{FF}(4n) \\ A_{FF}(4n) = C_{RR}(4n) \\ A_{RF}(4n+2) = C_{RF}(4n+2) \\ A_{FR}(4n+2) = C_{FR}(4n+2) \\ C_{RR}(4n+3) = C_{FF}(4n+3). \end{array} \right.$$

Put

$$(1.8) \quad \left\{ \begin{array}{l} y_{RF}(x) = \sum_{n=0}^{\infty} A_{RF}(4n+3) \frac{x^{4n+3}}{(4n+3)!} \quad (A_{RF}(3) = 2) \\ y_{RR}(x) = \sum_{n=0}^{\infty} A_{RR}(4n+1) \frac{x^{4n+1}}{(4n+1)!} \quad (A_{RR}(1) = 1) \\ y_{FR}(x) = \sum_{n=0}^{\infty} A_{FR}(4n+3) \frac{x^{4n+3}}{(4n+3)!} \quad (A_{FR}(3) = 2) \\ y_{FF}(x) = \sum_{n=0}^{\infty} A_{FF}(4n+1) \frac{x^{4n+1}}{(4n+1)!} \quad (A_{FF}(1) = 1) \end{array} \right.$$

$$(1.9) \quad \left\{ \begin{array}{l} z_{RF}(x) = \sum_{n=0}^{\infty} A_{RF}(4n+2) \frac{x^{4n+2}}{(4n+2)!} \quad (A_{RF}(2) = 1) \\ z_{RR}(x) = \sum_{n=0}^{\infty} A_{RR}(4n) \frac{x^{4n}}{(4n)!} \quad (A_{RR}(0) = 1) \\ z_{FR}(x) = \sum_{n=0}^{\infty} A_{FR}(4n+2) \frac{x^{4n+2}}{(4n+2)!} \quad (A_{FR}(2) = 1) \\ z_{FF}(x) = \sum_{n=0}^{\infty} A_{FF}(4n) \frac{x^{4n}}{(4n)!} \quad (A_{FF}(0) = 1) \end{array} \right.$$

$$(1.10) \left\{ \begin{aligned} \bar{y}_{RF}(x) &= \sum_{n=0}^{\infty} C_{RF}(4n+1) \frac{x^{4n+1}}{(4n+1)!} & (C_{RF}(1) = 1) \\ \bar{y}_{RR}(x) &= \sum_{n=0}^{\infty} C_{RR}(4n+3) \frac{x^{4n+3}}{(4n+3)!} & (C_{RR}(3) = 2) \\ \bar{y}_{FR}(x) &= \sum_{n=0}^{\infty} C_{FR}(4n+1) \frac{x^{4n+1}}{(4n+1)!} & (C_{FR}(1) = 1) \\ \bar{y}_{FF}(x) &= \sum_{n=0}^{\infty} C_{FF}(4n+3) \frac{x^{4n+3}}{(4n+3)!} & (C_{FF}(3) = 2) \end{aligned} \right.$$

$$(1.11) \left\{ \begin{aligned} \bar{z}_{RF}(x) &= \sum_{n=0}^{\infty} C_{RF}(4n+2) \frac{x^{4n+2}}{(4n+2)!} & (C_{RF}(2) = 1) \\ \bar{z}_{RR}(x) &= \sum_{n=0}^{\infty} C_{RR}(4n) \frac{x^{4n}}{(4n)!} & (C_{RR}(0) = 1) \\ \bar{z}_{FR}(x) &= \sum_{n=0}^{\infty} C_{FR}(4n+2) \frac{x^{4n+2}}{(4n+2)!} & (C_{FR}(2) = 1) \\ \bar{z}_{FF}(x) &= \sum_{n=0}^{\infty} C_{FF}(4n) \frac{x^{4n}}{(4n)!} & (C_{FF}(0) = 1). \end{aligned} \right.$$

In view of (1.7), we get

$$(1.12) \left\{ \begin{aligned} y_{RR}(x) &= y_{FF}(x), & z_{RR}(x) &= \bar{z}_{FF}(x), & z_{FF}(x) &= \bar{z}_{RR}(x) \\ z_{RF}(x) &= \bar{z}_{RF}(x), & z_{FR}(x) &= \bar{z}_{FR}(x), & \bar{y}_{RR}(x) &= \bar{y}_{FF}(x). \end{aligned} \right.$$

Note that, for example, in taking

$$A_{RF}(3) = A_{FR}(3) = 2,$$

we are listing the up-down permutation (1, 3, 2) and (2, 3, 1) both under *RF* and *FR*. This is done so that the recurrences given below will be satisfied. A like remark applies in a number of other instances, as is evident from an examination of (1.8), ..., (1.11).

In the remainder of the paper we evaluate the sixteen enumerants defined in (1.8), ..., (1.11). For a summary of results see §6 below.

2. Evaluation of $y_{RF}(x)$, $y_{RR}(x)$, $y_{FF}(x)$, $y_{FR}(x)$. We consider first $y_{RF}(x)$. The method employed is to take a typical permutation of Z_n and consider the effect of removing the largest element. This is indeed the method used in [1]. Clearly the element removed must be a peak. The given permutation breaks into two pieces one of which may be vacuous. Thus for $A_{RF}(4n+3)$ we get the recurrence

$$(2.1) \quad A_{RF}(4n + 3) = \sum_{k=0}^{n-1} \binom{4n + 2}{4k + 3} A_{RF}(4k + 3)A_{RF}(4n - 4k - 1)$$

$$(n \geq 1, \quad A_{RF}(3) = 2).$$

It follows that the generating function $y_{RF}(x)$ satisfies the differential equation

$$(2.2) \quad y'_{RF}(x) = y^2_{RF}(x) + x^2.$$

Now put

$$(2.3) \quad y_{RF}(x) = -\frac{U'(x)}{U(x)},$$

where

$$(2.4) \quad U(x) = \sum_{n=0}^{\infty} a_n \frac{x^{4n}}{(4n)!} \quad (a_0 = 1).$$

Substituting (2.3) in (2.2) we get

$$(2.5) \quad U''(x) + x^2U(x) = 0.$$

This implies the recurrence

$$a_{n+1} + (4n + 1)(4n + 2)a_n = 0 \quad (n = 0, 1, 2, \dots)$$

and therefore

$$(2.6) \quad a_n = (-1)^n 1 \cdot 5 \cdot 9 \dots (4n - 3) \cdot 2 \cdot 6 \cdot 10 \dots (4n - 2).$$

Thus

$$\frac{a_n}{(4n)!} = \frac{(-1)^n}{3 \cdot 7 \cdot 11 \dots (4n - 1) \cdot 4 \cdot 8 \cdot 12 \dots 4n} = \frac{(-1)^n}{4^{2n}} \frac{1}{n! (3/4)_n},$$

where

$$(a)_n = a(a + 1) \dots (a + n - 1).$$

Hence (2.4) becomes

$$(2.7) \quad U(x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{4^{2n}} \frac{x^{4n}}{n! (3/4)_n} = {}_0F_1 \left(-; \frac{3}{4}; \frac{x^4}{16} \right)$$

in the notation of generalized hypergeometric functions [5, Ch. 5].

Alternatively we may write [5, p. 108]

$$(2.8) \quad U(x) = \left(\frac{1}{2}x\right)^{\frac{1}{2}} \Gamma\left(\frac{3}{4}\right) J_{-1/4}\left(\frac{1}{2}x^2\right),$$

where $J_{-1/4}(z)$ denotes the Bessel function of order $-1/4$.

In the next place, we have for $A_{RR}(4n+1)$ the recurrence

$$(2.9) \quad A_{RR}(4n+1) = \sum_{k=0}^{n-1} \binom{4n}{4k+3} A_{RF}(4k+3) A_{RR}(4n-4k+1) \quad (n > 0, \quad A_{RR}(1) = 1).$$

This gives

$$(2.10) \quad y'_{RR}(x) = y_{RF}(x)y_{RR}(x) + 1.$$

Hence, by (2.3), (2.10) becomes

$$U(x)y'_{RR}(x) + U'(x)y_{RR}(x) = U(x)$$

and therefore

$$(2.11) \quad y_{RR}(x) = \frac{1}{U(x)} \int U(x),$$

where generally

$$(2.12) \quad \int f(x) = \int_0^x f(t) dt.$$

By (1.12) this implies

$$(2.13) \quad y_{FF}(x) = \frac{1}{U(x)} \int U(x).$$

Note that, by (2.7) and (2.12),

$$(2.14) \quad \int U = \sum_{n=0}^{\infty} \frac{(-1)^n}{4^{2n}} \frac{x^{4n+1}}{n!(3/4)_n(4n+1)}.$$

The enumerant $A_{FR}(4n+1)$ satisfies the recurrence

$$(2.15) \quad A_{FR}(4n+3) = \sum_{k=0}^n \binom{4n+2}{4k+1} A_{FF}(4k+1) A_{RR}(4n-4k+1) \quad (n \geq 0, \quad A_{FR}(3) = 2).$$

Thus

$$y'_{FR}(x) = y_{FF}(x)y_{RR}(x).$$

Hence, by (2.11) and (2.13),

$$(2.16) \quad y'_{FR}(x) = \left\{ \frac{1}{U(x)} \int U(x) \right\}^2.$$

It can be verified that

$$(2.17) \quad U^2(x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{4^{2n}} \frac{(n+1)_n x^{4n}}{n!(3/4)_n(3/4)_n}.$$

3. Evaluation of $z_{RF}(x)$, $z_{RR}(x)$, $z_{FF}(x)$, $z_{FR}(x)$. As above we have first

$$(3.1) \quad A_{RF}(4n+2) = \sum_{k=0}^{n-1} \binom{4n+1}{4k+3} A_{RF}(4k+3) A_{RF}(4n-4k-2) \quad (n > 0, \quad A_{RF}(2) = 1).$$

This yields

$$(3.2) \quad z'_{RF}(x) = y_{RF}(x)z_{RF}(x) + x.$$

Hence, by (2.3),

$$U(x)z'_{RF}(x) + U'(x)z_{RF}(x) = xU(x),$$

so that

$$(3.3) \quad z_{RF}(x) = \frac{1}{U(x)} \int xU.$$

Note that

$$(3.4) \quad \int xU = \sum_{n=0}^{\infty} \frac{(-1)^n}{4^{2n}} \frac{x^{4n+2}}{n!(3/4)_n(4n+2)}.$$

Next

$$A_{RR}(4n) = \sum \binom{4n-1}{4k+3} A_{RF}(4k+3) A_{RR}(4n-4k-4) \quad (n \geq 1, \quad A_{RR}(0) = 1).$$

Thus

$$(3.5) \quad z'_{RR}(x) = y_{RF}(x)z_{RR}(x).$$

It follows that

$$(3.6) \quad z_{RR}(x) = \frac{1}{U(x)}.$$

For $A_{FF}(4n)$ we have

$$(3.7) \quad A_{FF}(4n) = \sum_{k=0}^{n-1} \binom{4n-1}{4k+1} A_{FF}(4k+1) A_{RF}(4n-4k-2) \quad (n > 0, \quad A_{FF}(0) = 1).$$

This gives

$$(3.8) \quad z'_{FF}(x) = y_{FF}(x)z_{RF}(x).$$

Therefore, by (2.13) and (3.3),

$$(3.9) \quad z'_{FF}(x) = \frac{1}{U^2(x)} \int U(x) \cdot \int xU(x).$$

As for $A_{FR}(4n+2)$, we have

$$(3.10) \quad A_{FR}(4n+2) = \sum_{k=0}^n \binom{4n+1}{4k+1} A_{FF}(4k+1) A_{RR}(4n-2k) \quad (n \geq 0, \quad A_{FR}(2) = 1).$$

Thus

$$(3.11) \quad z'_{FR}(x) = y_{FF}(x)z_{RR}(x).$$

Then, by (2.13) and (3.6),

$$(3.12) \quad z'_{FR}(x) = \frac{1}{U^2(x)} \int U(x).$$

4. Evaluation of $\bar{y}_{RF}(x)$, $\bar{y}_{RR}(x)$, $\bar{y}_{FF}(x)$, $\bar{y}_{FR}(x)$. To begin with

$$(4.1) \quad C_{RF}(4n + 1) = \sum_{k=0}^{n-1} \binom{4n}{4k + 2} C_{RF}(4k + 2) A_{RF}(4n - 4k - 2) \quad (n > 0, \quad C_{RF}(1) = 1).$$

This gives

$$(4.2) \quad \bar{y}'_{RF}(x) = \bar{z}_{RF}(x) z_{RF}(x).$$

By (1.12), $\bar{z}_{RF}(x) = z_{RF}(x)$, so that (4.2) reduces to

$$(4.3) \quad \bar{y}'_{RF}(x) = \{z_{RF}(x)\}^2.$$

This formula together with (3.3) determines $\bar{y}_{RF}(x)$.

Next

$$(4.4) \quad C_{RR}(4n + 3) = \sum_{k=0}^n \binom{4n + 2}{4k + 2} C_{RF}(4k + 2) A_{RR}(4n - 4k) \quad (n \geq 0, \quad C_{RR}(3) = 1),$$

so that

$$(4.5) \quad \bar{y}'_{RR}(x) = \bar{z}_{RF}(x) z_{RR}(x) = z_{RF}(x) z_{RR}(x).$$

Hence

$$(4.6) \quad \bar{y}'_{RR}(x) = \frac{1}{U^2(x)} \int xU(x).$$

Since by (1.12), $\bar{y}_{FF}(x) = \bar{y}_{RR}(x)$, we have also

$$(4.7) \quad \bar{y}'_{FF}(x) = \frac{1}{U^2(x)} \int xU(x).$$

In the next place

$$(4.8) \quad C_{FR}(4n + 1) = \sum_{k=0}^n \binom{4n}{4k} C_{FF}(4k) A_{RR}(4n - 4k) \quad (n \geq 0, \quad C_{FR}(1) = 1).$$

This gives

$$(4.9) \quad \bar{y}'_{FR}(x) = \bar{z}_{FF}(x) z_{RR}(x) = \{z_{RR}(x)\}^2,$$

since $\bar{z}_{FF}(x) = z_{RR}(x)$. Therefore, by (3.6),

$$(4.10) \quad \bar{y}'_{FR}(x) = \frac{1}{U^2(x)}.$$

5. Second solution of (2.5). One solution of the differential equation

$$(5.1) \quad w'' + x^2w = 0$$

is given by $w_1 = U(x)$. For a second solution we take

$$(5.2) \quad w_2 = U(x)V(x), \quad V(x) = \sum_{n=0}^{\infty} b_n \frac{x^{4n+1}}{(4n+1)!} \quad (b_0 = 1).$$

Clearly w_1 and w_2 are linearly independent. Substituting from (5.2) in (5.1) we get

$$(5.3) \quad 2U'(x)V'(x) + U(x)V''(x) = 0.$$

This gives

$$(5.4) \quad V'(x) = \frac{1}{U^2(x)}.$$

Comparing (5.4) with (4.10), it is clear that

$$(5.5) \quad V(x) = \bar{y}_{FR}(x),$$

so that the second solution of (5.1) becomes

$$(5.6) \quad w_2 = U(x)\bar{y}_{FR}(x).$$

6. Summary of results.

$$(6.1) \quad y_{RF}(x) = -\frac{U'(x)}{U(x)}$$

$$(6.2) \quad y_{RR}(x) = y_{FF}(x) = \frac{1}{U(x)} \int U(x)$$

$$(6.3) \quad y'_{FR}(x) = \left\{ \frac{1}{U(x)} \int U(x) \right\}^2$$

$$(6.4) \quad z_{RF}(x) = \bar{z}_{RF}(x) = \frac{1}{U(x)} \int xU(x)$$

$$(6.5) \quad z_{RR}(x) = \bar{z}_{FF}(x) = \frac{1}{U(x)}$$

$$(6.6) \quad z'_{FF}(x) = \bar{z}'_{RR}(x) = \frac{1}{U^2(x)} \int U(x) \cdot \int xU(x)$$

$$(6.7) \quad z'_{FR}(x) = z'_{FR}(x) = \frac{1}{U^2(x)} \int U(x)$$

$$(6.8) \quad \bar{y}'_{RF}(x) = \{z_{RF}(x)\}^2$$

$$(6.9) \quad \bar{y}_{RR}(x) = \bar{y}_{FF}(x) = \frac{1}{U^2(x)} \int xU(x)$$

$$(6.10) \quad \bar{y}'_{FR}(x) = \frac{1}{U^2(x)}$$

$$(6.11) \quad U(x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{4^{2n}} \frac{x^{4n}}{n!(3/4)_n}$$

$$(6.12) \quad U^2(x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{4^{2n}} \frac{(n + \frac{1}{2})_n x^{4n}}{n!(3/4)_n(3/4)_n}.$$

Linearly independent solutions of

$$w'' + x^2w = 0$$

are furnished by

$$(6.13) \quad w_1 = U(x), \quad w_2 = U(x)\bar{y}_{FR}(x).$$

7. Generalizations. We may define *doubly* up-down permutations as permutations (a_1, a_2, \dots, a_n) of Z_n such that

$$(7.1) \quad a_{2k-1} < a_{2k}, \quad a_{2k} > a_{2k+1} \quad (k = 1, 2, \dots, [n/2])$$

and

$$(7.2) \quad a_{4k-2} < a_{4k}, \quad a_{4k} > a_{4k+2} \quad (k = 1, 2, \dots, [n/4]).$$

Similarly we may define *triply* up-down as permutations that satisfy (7.1) and (7.2) and in addition

$$(7.3) \quad a_{8k-4} < a_{8k}, \quad a_{8k} > a_{8k+4} \quad (k = 1, 2, \dots, [n/8]).$$

It is clear how to extend this definition to r -ply up-down permutations. Thus this suggests the enumeration of permutations of these types.

An extension in a different direction is the following. Let $A_3(n)$ denote the number of permutations of Z_n that satisfy

$$(7.4) \quad a_{3k-2} < a_{3k-1} < a_{3k}, \quad a_{3k} > a_{3k+1} \quad (k = 1, 2, \dots, [n/3]).$$

Then, as a special case of a result proved in [2], [3], we have

$$(7.5) \quad \sum_{n \times 0}^{\infty} A_3(3n) \frac{x^{3n}}{(3n)!} = \left\{ \sum_{n=0}^{\infty} \frac{(-1)^n}{(3n)!} \right\}^{-1}.$$

This suggests the consideration of permutations that satisfy (7.4) and in addition

$$(7.6) \quad a_{9k-6} < a_{9k-3} < a_{9k}, \quad a_{9k} > a_{9k+3} \quad (k = 1, 2, \dots, [n/9]).$$

Moreover further restrictions analogous to (7.3) can also be introduced.

However we shall not treat these extensions in the present paper.

REFERENCES

1. L. Carlitz, *Enumeration of up-down permutations by number of rises*, Pacific J. Math., **45** (1973), 49-58.
2. ———, *Generating functions for a special class of permutations*, Proc. Amer. Math. Soc., **47** (1975), 251-256.
3. ———, *Permutations with prescribed pattern*, Mathematische Nachrichten, **58** (1973), 31-53.
4. E. Netto, *Lehrbuch der Combinatorik*, Teubner, Leipzig and Berlin, 1927.
5. E. D. Rainville, *Special Functions*, Macmillan, New York, 1960.

Received July 6, 1976.

DUKE UNIVERSITY
DURHAM, NC 27706

| | |
|---|-----|
| William H. Barker, <i>Noether's theorem for plane domains with hyperelliptic double</i> | 1 |
| Michael James Beeson, <i>Non-continuous dependence of surfaces of least area on the boundary curve</i> | 11 |
| Horst Behncke, <i>Functions acting in weighted Orlicz algebras</i> | 19 |
| Howard Edwin Bell, <i>A commutativity study for periodic rings</i> | 29 |
| Peter Botta and Stephen J. Pierce, <i>The preservers of any orthogonal group</i> | 37 |
| Douglas S. Bridges, <i>The constructive Radon-Nikodým theorem</i> | 51 |
| James Dennis Brom, <i>The theory of almost periodic functions in constructive mathematics</i> | 67 |
| N. Burgoyne and C. Williamson, <i>Semi-simple classes in Chevalley type groups</i> | 83 |
| Douglas Cameron, <i>A class of maximal topologies</i> | 101 |
| L. Carlitz, <i>Enumeration of doubly up-down permutations</i> | 105 |
| Paul Robert Chernoff, <i>The quantum n-body problem and a theorem of Littlewood</i> | 117 |
| Jo-Ann Deborah Cohen, <i>Locally bounded topologies on $F(X)$</i> | 125 |
| Heinz Otto Cordes and Robert Colman McOwen, <i>Remarks on singular elliptic theory for complete Riemannian manifolds</i> | 133 |
| Micheal Neal Dyer, <i>Correction to: "Rational homology and Whitehead products"</i> | 143 |
| Robert Fernholz, <i>Factorization of Radonifying transformations</i> | 145 |
| Lawrence Arthur Fialkow, <i>A note on quasisimilarity. II</i> | 151 |
| Harvey Charles Greenwald, <i>Lipschitz spaces of distributions on the surface of unit sphere in Euclidean n-space</i> | 163 |
| Albrecht Irle, <i>On the measurability of conditional expectations</i> | 177 |
| Tom (Roy Thomas Jr.) Jacob, <i>Matrix transformations involving simple sequence spaces</i> | 179 |
| A. Katsaras, <i>Continuous linear maps positive on increasing continuous functions</i> | 189 |
| Kenneth Kunen and Judith Roitman, <i>Attaining the spread at cardinals of cofinality ω</i> | 199 |
| Lawrence Louis Larmore and Robert David Rigdon, <i>Enumerating normal bundles of immersions and embeddings of projective spaces</i> | 207 |
| Ch. G. Philos and V. A. Staïkos, <i>Asymptotic properties of nonoscillatory solutions of differential equations with deviating argument</i> | 221 |
| Peter Michael Rosenthal and Ahmed Ramzy Sourour, <i>On operator algebras containing cyclic Boolean algebras</i> | 243 |
| Polychronis Strantzalos, <i>Strikt fast gleichgradig-stetige und eigentliche Aktionen</i> | 253 |
| Glenn Francis Webb, <i>Exponential representation of solutions to an abstract semi-linear differential equation</i> | 269 |
| Scott Andrew Wolpert, <i>The finite Weil-Petersson diameter of Riemann space</i> | 281 |