

Pacific Journal of Mathematics

**AN OPERATOR ALGEBRA WHICH IS NOT CLOSED IN THE
CALKIN ALGEBRA**

KENNETH R. DAVIDSON AND CHE-KAO FONG

AN OPERATOR ALGEBRA WHICH IS NOT CLOSED IN THE CALKIN ALGEBRA

K. R. DAVIDSON AND C. K. FONG

An example is constructed of an operator algebra which is closed in the weak operator topology but has a nonclosed image in the Calkin algebra.

Given a norm closed algebra of operators on Hilbert space, is the sum of this algebra with the ideal of all compact operators a closed set? The answer is well known to be yes for C^* -algebras [2]. It was shown by Arveson [1] to be closed for the discrete triangular algebra, and has recently been shown for all nest algebras by Muhly using his methods of [3]. Also, it is easily established for algebras of subnormal operators and Toeplitz operators. In this note, we will construct an algebra for which the sum is not closed. In fact, this algebra will be closed in the weak operator topology.

Let \mathcal{A} be a norm closed algebra on a Hilbert space \mathcal{H} , let \mathcal{KB} denote the compact operators, and let π denote the quotient map of $\mathcal{L}(\mathcal{H})$ onto the Calkin algebra $\mathcal{L}(\mathcal{H})/\mathcal{KB}$. It is apparent that $\mathcal{A} + \mathcal{KB}$ is closed if and only if the image $\pi(\mathcal{A})$ is closed in the Calkin algebra. We will require the following elementary observation.

PROPOSITION 1. *If \mathcal{A} is a norm closed subalgebra of $\mathcal{L}(\mathcal{H})$, then $\mathcal{A} + \mathcal{KB}$ is closed if and only if $\mathcal{A}/\mathcal{A} \cap \mathcal{KB}$ and $\pi(\mathcal{A}) = \mathcal{A} + \mathcal{KB}/\mathcal{KB}$ are isomorphic as Banach algebras.*

COROLLARY 1. *If \mathcal{A} is a norm closed algebra such that \mathcal{A} contains no compact operators except 0, then $\mathcal{A} + \mathcal{KB}$ is closed if and only if the quotient map π is bounded below on \mathcal{A} .*

The proof is a simple application of the Closed Graph Theorem.

PROPOSITION 2. *There is a (commutative) operator algebra \mathcal{A} , closed in the weak operator topology, such that $\mathcal{A} + \mathcal{KB}$ is not closed.*

Proof. Let $\{e_n\}$ indexed on the integers be a basis for \mathcal{H} . Let U be the bilateral shift defined by $Ue_n = e_{n+1}$, and let P be the one dimensional projection onto the span of e_0 . Let $A = U + P$, and let \mathcal{A} be the weakly closed algebra generated by A .

We will show that \mathcal{A} does not contain any nonzero compact

operators. Suppose \mathcal{A} contains a compact operator K . Then we can find polynomials p_n so that $p_n(A)$ tends to K in the weak operator topology. The restriction of $p_n(A)$ to \mathcal{H} , the span of $\{e_n; n \geq 1\}$ is equal to the restriction of $p_n(U)$ to \mathcal{H} which is unitarily equivalent to the Toeplitz operator of multiplication by p_n on H^2 of the circle. Since the algebra of analytic Toeplitz operators is weakly closed, and no nonzero Toeplitz operator is compact, we conclude that $p_n(A)|_{\mathcal{H}}$ tends to zero in the weak operator topology. In particular we get that $(Ke_i, e_j) = 0$ for all $i \geq 1$ and all j . Since the same is true for coanalytic Toeplitz operators and $p_n(A^*)|_{\mathcal{H}^\perp}$, we note also that $(Ke_i, e_j) = 0$ for all i and all $j \leq 0$.

Furthermore, if we write $p_n(z) = \sum_k \alpha_k^{(n)} z^k$, the fact that the Toeplitz operators for p_n tend weakly to zero implies that for each k , $\lim_{n \rightarrow \infty} \alpha_k^{(n)} = 0$. So, without loss of generality, we can assume that $\alpha_k^{(n)} = 0$ for $1 \leq k \leq N$ and all n . We notice that for (i, j) belonging to the triangle T_N :

$$0 \geq i \geq -N, \quad 0 \leq j \leq N, \quad j - i \leq N$$

and for $m \geq N$, we have $(A^m e_i, e_j) = 1$. Hence $(p_n(A)e_i, e_j)$ is constant for (i, j) in T_N . This must also hold in the limit, so (Ke_i, e_j) is constant for (i, j) in T_N . But this N was arbitrary, so (Ke_i, e_j) is constant on $i \leq 0, j \geq 0$. Since K is a bounded operator, this must be zero. Together with the previous paragraph, we see that K is zero.

However, $\|A^n\| \geq \sqrt{n}$ and $\|\pi(A^n)\| = \|\pi(U^n)\| = 1$, so π is not bounded below on \mathcal{A} . Thus by Corollary 1, $\mathcal{A} + \mathcal{L}\mathcal{E}$ is not closed.

COROLLARY 2. *There is a separable (commutative) norm closed algebra \mathcal{B} such that $\mathcal{B} + \mathcal{L}\mathcal{E}$ is not closed.*

Proof. Take \mathcal{B} to be the norm closed algebra generated by A .

We would like to thank Eric Nordgren for some helpful discussions.

REFERENCES

1. W. B. Arveson, *Interpolation problems in nest algebras*, J. Functional Anal., **20** (1975), 208-233.
2. J. Dixmier, *Les C*-Algebres et Leur Representations*, Gauthier-Villars, Paris, 1964.
3. R. I. Loeb and P. S. Muhly, *Analyticity and flows in von Neumann algebras*, to appear.

Received March 16, 1977. Research partially supported by U.S. National Science Foundation Grant MCS76-21044.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
CAMBRIDGE, MA 02139

AND

UNIVERSITY OF TORONTO
TORONTO, CANADA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

C. W. CURTIS

University of Oregon
Eugene, OR 97403

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

| | |
|---|-----|
| Kazuo Anzai and Shiro Ishikawa, <i>On common fixed points for several continuous affine mappings</i> | 1 |
| Bruce Alan Barnes, <i>When is a representation of a Banach *-algebra Naimark-related to a *-representation</i> | 5 |
| Richard Dowell Byrd, Justin Thomas Lloyd, Franklin D. Pedersen and James Wilson Stepp, <i>Automorphisms of the semigroup of finite complexes of a periodic locally cyclic group</i> | 27 |
| Donald S. Coram and Paul Frazier Duvall, Jr., <i>Approximate fibrations and a movability condition for maps</i> | 41 |
| Kenneth R. Davidson and Che-Kao Fong, <i>An operator algebra which is not closed in the Calkin algebra</i> | 57 |
| Garret J. Etgen and James Pawlowski, <i>A comparison theorem and oscillation criteria for second order differential systems</i> | 59 |
| Philip Palmer Green, <i>C*-algebras of transformation groups with smooth orbit space</i> | 71 |
| Charles Allen Jones and Charles Dwight Lahr, <i>Weak and norm approximate identities are different</i> | 99 |
| G. K. Kalisch, <i>On integral representations of piecewise holomorphic functions</i> | 105 |
| Y. Kodama, <i>On product of shape and a question of Sher</i> | 115 |
| Heinz K. Langer and B. Textorius, <i>On generalized resolvents and Q-functions of symmetric linear relations (subspaces) in Hilbert space</i> | 135 |
| Albert Edward Livingston, <i>On the integral means of univalent, meromorphic functions</i> | 167 |
| Wallace Smith Martindale, III and Susan Montgomery, <i>Fixed elements of Jordan automorphisms of associative rings</i> | 181 |
| R. Kent Nagle, <i>Monotonicity and alternative methods for nonlinear boundary value problems</i> | 197 |
| Richard John O'Malley, <i>Approximately differentiable functions: the r topology</i> | 207 |
| Mangesh Bhalchandra Rege and Kalathoor Varadarajan, <i>Chain conditions and pure-exactness</i> | 223 |
| Christine Ann Shannon, <i>The second dual of C(X)</i> | 237 |
| Sin-ei Takahasi, <i>A characterization for compact central double centralizers of C*-algebras</i> | 255 |
| Theresa Phillips Vaughan, <i>A note on the Jacobi-Perron algorithm</i> | 261 |
| Arthur Anthony Yanushka, <i>A characterization of PSp(2m, q) and PΩ(2m + 1, q) as rank 3 permutation groups</i> | 273 |