

Pacific Journal of Mathematics

FIXED POINT SETS OF PEANO CONTINUA

JOHN ROWLAY MARTIN

FIXED POINT SETS OF PEANO CONTINUA

JOHN R. MARTIN

For each positive integer $n = 1, 2, \dots$, it is shown that there is an $(n + 1)$ -dimensional acyclic LC^{n-1} continuum X_n containing an n -dimensional sphere which is not the fixed point set of any self-map of X_n .

1. **Introduction.** A subset A of a topological space X is called a *fixed point set* of X if there is a (continuous) map $f: X \rightarrow X$ such that $f(x) = x$ iff $x \in A$. If X is Hausdorff, then A is closed, and, clearly, every retract of X is a fixed point set of X . It is possible that a space X may have the property that each of its nonempty closed subsets is a fixed point set of X . The problem of determining which spaces have this property, called the *complete invariance property* by L. E. Ward, Jr. in [5], has been investigated by H. Robbins, Helga Schirmer, and L. E. Ward, Jr. Some spaces known to have the complete invariance property include n -cells [1], dendrites [2], convex subsets of Banach spaces [5], compact manifolds without boundary [3], and all compact triangulable manifolds with or without boundary [4].

The general question as to what properties a space must satisfy to insure that it has the complete invariance property has not been resolved. In fact, in [5, p. 553] L. E. Ward, Jr. asks the following question.

Does every Peano continuum have the complete invariance property?

The purpose of this note is to show that even acyclic Peano continua which possess higher order local connectedness need not have the complete invariance property. Indeed, for each positive integer $n = 1, 2, \dots$, we give an example of an $(n + 1)$ -dimensional acyclic LC^{n-1} continuum X_n which fails to have the complete invariance property. Moreover, X_n contains an n -dimensional sphere which is not a fixed point set of X_n .

2. **Notation and the construction of X_n .** We shall let E^n denote Euclidean n -dimensional space, and we shall consider E^m to be canonically imbedded in E^n if $m < n$. The closed unit ball in E^{n+1} shall be denoted by B^{n+1} and the boundary $\text{Bd } B^{n+1}$ of B^{n+1} shall be denoted by S^n .

Consider the rectangle in $E^2 \subset E^{n+2}$ with vertices $(1, -1, 0, \dots, 0)$, $(0, -1, 0, \dots, 0)$, $(0, 1, 0, \dots, 0)$, $(1, 1, 0, \dots, 0)$. Let D denote the closed disk consisting of this rectangle together with its interior in

E^2 , and let A denote the segment in $\text{Bd } D$ with endpoints $(0, -1, 0, \dots, 0)$ and $(0, 1, 0, \dots, 0)$. We shall let C denote the closure of the curve in E^2 whose equation is given by $y = \sin \pi/x$ for $0 < x \leq 1$.

Let $\{B_t^{n+1} | 1 \leq t < \infty\}$ be a disjoint collection of $(n + 1)$ -dimensional balls in E^{n+2} satisfying the following properties.

(1) $\bigcup_{1 \leq t < \infty} B_t^{n+1}$ is homeomorphic to $B^{n+1} \times [1, \infty)$ under a homeomorphism which sends B_t^{n+1} onto $B^{n+1} \times \{t\}$ for each t in the half-open interval $[1, \infty)$.

(2) $\lim_{t \rightarrow \infty} \delta(B_t^{n+1}) = 0$, where $\delta(B_t^{n+1})$ denotes the diameter of B_t^{n+1} .

(3) For each t in $[1, \infty)$,

$$B_t^{n+1} \cap E^2 = \left\{ \left(\frac{1}{t}, \sin \pi t, 0, \dots, 0 \right) \right\}.$$

Let $S_t^n = \text{Bd } B_t^{n+1}$. If J is an interval in the real line, we define

$$B_J^{n+1} = \bigcup_{t \in J} B_t^{n+1} \quad \text{and} \quad S_J^n = \bigcup_{t \in J} S_t^n.$$

Definition. For $n = 1, 2, \dots$, we define

$$X_n = D \cup S_{[1, \infty)}^n.$$

Essentially, $S_{[1, \infty)}^n$ is obtained by taking a cone over an n -sphere, removing the vertex, and then winding the resulting tube of n -spheres in a “ $\sin 1/x$ ” fashion in E^{n+2} so as to converge to the limit interval A . This procedure is carried out so that the intersection of E^2 with the closure of $S_{[1, \infty)}^n$ is precisely C . The disk D is then attached to the closure of $S_{[1, \infty)}^n$ along C to obtain X_n .

3. The properties of X_n .

Property 1. Each X_n is an $(n + 1)$ -dimensional acyclic continuum.

Proof. By acyclic we mean that X_n has the Čech cohomology of a point. Since X_n is clearly an $(n + 1)$ -dimensional continuum, we need only show that it is acyclic. But

$$X_n = \bigcap_{t \geq 1} (D \cup S_{[1, \infty)}^n \cup B_{[t, \infty)}^{n+1})$$

and, for each t , $D \cup B_{[t, \infty)}^{n+1}$ is a deformation retract of $D \cup S_{[1, \infty)}^n \cup B_{[t, \infty)}^{n+1}$. Since $D \cup B_{[t, \infty)}^{n+1}$ is contractible, each of the spaces $D \cup S_{[1, \infty)}^n \cup B_{[t, \infty)}^{n+1}$ is contractible. Therefore, by the continuity axiom for Čech cohomology, X_n is acyclic.

Property 2. Each X_n is LC^{n-1} at points of A , and locally contractible elsewhere.

Proof. Clearly, every point p in $X_n - A$ has a neighborhood which is finitely triangulable, and therefore X_n is locally contractible at p . If $p \in A$, then p has a compact neighborhood consisting of a closed disk K and infinitely many disjoint cylinders of the form $S_{[t, t+\lambda]}^n$, each of which intersects K in an arc. It then follows that X_n is LC^{n-1} at p .

Property 3. S_1^n is not a fixed point set of X_n .

Proof. Since S_1^n is not contractible in $S_{[1, \infty)}^n \cup A$ and $(S_{[1, \infty)}^n \cup A) \cap D = C$, it follows that S_1^n is not contractible in X_n .

Now suppose that $f: X_n \rightarrow X_n$ is a map whose fixed point set is precisely S_1^n . If $f(A) \subset A$, then f has a fixed point in A . Since this is not possible, there is a point p in A such that $f(p) \in X_n - A$. Let V be a contractible neighborhood of $f(p)$ in X_n , and let $\beta: V \times I \rightarrow V$ denote a homotopy which deforms V to a point. Since f is continuous at p , there is a neighborhood U of p in X_n such that $f(U) \subset V$. Then, for some r in $[1, \infty)$, we have $S_r^n \subset U$. Let $\alpha: S_1^n \times I \rightarrow S_{[1, r]}^n$ be a homotopy which deforms S_1^n onto S_r^n .

Define a homotopy $H: S_1^n \times I \rightarrow X_n$ by

$$H(x, t) = \begin{cases} f(\alpha(x, 2t)), & (\text{if } 0 \leq t \leq \frac{1}{2}), \\ \beta(f(\alpha(x, 1)), 2t - 1), & (\text{if } \frac{1}{2} \leq t \leq 1). \end{cases}$$

It is easy to check that H is a homotopy which deforms S_1^n to a point in X_n . This contradiction shows that S_1^n is not a fixed point set of X_n .

4. Problem. Let us restrict our discussion to the class of compacta. Then all the examples of spaces having the complete invariance property which are mentioned in the introduction of this paper are ANR-spaces. In fact, the first three examples are AR-spaces. In view of this, it seems appropriate to suggest the following problem.

Does every AR-space (ANR-space) have the complete invariance property?

REFERENCES

1. H. Robbins, *Some complements to Brouwer's fixed point theorem*, Israel J. Math., **5** (1967), 225-226.

2. H. Schirmer, *Properties of fixed point sets on dendrites*, Pacific J. Math., **36** (1971), 795-810.
3. ———, *Fixed point sets of homeomorphisms of compact surfaces*, Israel J. Math., **10** (1971), 373-378.
4. ———, *Fixed point sets of polyhedra*, Pacific J. Math., **52** (1974), 221-226.
5. L. E. Ward, Jr., *Fixed point sets*, Pacific J. Math., **47** (1973), 553-565.

Received June 16, 1977. The research for this article was supported in part by the National Research Council of Canada (Grant A8205).

UNIVERSITY OF SASKATCHEWAN
SASKATOON, CANADA S7N 0W0

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

C. W. CURTIS

University of Oregon
Eugene, OR 97403

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
OSAKA UNIVERSITY

Gerald Arthur Anderson, <i>Computation of the surgery obstruction groups</i> $L_{4k}(1; Z_p)$	1
R. K. Beatson, <i>The degree of monotone approximation</i>	5
Sterling K. Berberian, <i>The character space of the algebra of regulated functions</i>	15
Douglas Michael Campbell and Jack Wayne Lamoreaux, <i>Continua in the plane with limit directions</i>	37
R. J. Duffin, <i>Algorithms for localizing roots of a polynomial and the Pisot Vijayaraghavan numbers</i>	47
Alessandro Figà-Talamanca and Massimo A. Picardello, <i>Functions that operate on the algebra $B_0(G)$</i>	57
John Erik Fornæss, <i>Biholomorphic mappings between weakly pseudoconvex domains</i>	63
Andrzej Granas, Ronald Bernard Guenther and John Walter Lee, <i>On a theorem of S. Bernstein</i>	67
Jerry Grossman, <i>On groups with specified lower central series quotients</i>	83
William H. Julian, Ray Mines, III and Fred Richman, <i>Algebraic numbers, a constructive development</i>	91
Surjit Singh Khurana, <i>A note on Radon-Nikodým theorem for finitely additive measures</i>	103
Garo K. Kiremidjian, <i>A Nash-Moser-type implicit function theorem and nonlinear boundary value problems</i>	105
Ronald Jacob Leach, <i>Coefficient estimates for certain multivalent functions</i>	133
John Alan MacBain, <i>Local and global bifurcation from normal eigenvalues. II</i>	143
James A. MacDougall and Lowell G. Sweet, <i>Three dimensional homogeneous algebras</i>	153
John Rowlay Martin, <i>Fixed point sets of Peano continua</i>	163
R. Daniel Mauldin, <i>The boundedness of the Cantor-Bendixson order of some analytic sets</i>	167
Richard C. Metzler, <i>Uniqueness of extensions of positive linear functions</i>	179
Rodney V. Nillsen, <i>Moment sequences obtained from restricted powers</i>	183
Keiji Nishioka, <i>Transcendental constants over the coefficient fields in differential elliptic function fields</i>	191
Gabriel Michael Miller Obi, <i>An algebraic closed graph theorem</i>	199
Richard Cranston Randell, <i>Quotients of complete intersections by C^* actions</i>	209
Bruce Reznick, <i>Banach spaces which satisfy linear identities</i>	221
Bennett Setzer, <i>Elliptic curves over complex quadratic fields</i>	235
Arne Stray, <i>A scheme for approximating bounded analytic functions on certain subsets of the unit disc</i>	251
Nicholas Th. Varopoulos, <i>A remark on functions of bounded mean oscillation and bounded harmonic functions. Addendum to: "BMO functions and the $\bar{\partial}$-equation"</i>	257
Charles Irvin Vinsonhaler, <i>Torsion free abelian groups quasi-projective over their endomorphism rings. II</i>	261
Thomas R. Wolf, <i>Characters of p'-degree in solvable groups</i>	267
Toshihiko Yamada, <i>Schur indices over the 2-adic field</i>	273