

Pacific Journal of Mathematics

**A SCHEME FOR APPROXIMATING BOUNDED ANALYTIC
FUNCTIONS ON CERTAIN SUBSETS OF THE UNIT DISC**

ARNE STRAY

A SCHEME FOR APPROXIMATING BOUNDED ANALYTIC FUNCTIONS ON CERTAIN SUBSETS OF THE UNIT DISC

A. STRAY

We denote by H^∞ the space of all bounded analytic functions in the unit disc $D = \{z: |z| < 1\}$. We consider a relatively closed subset S of D with the following property: If $f \in H^\infty$ and its restriction $f|_S$ to S is uniformly continuous, there exists a bounded sequence of polynomials $\{p_n\}$ such that

- (i) $p_n \rightarrow f$ uniformly on compact subsets of D .
- (ii) $p_n \rightarrow f$ uniformly on S .

A set S with this property, is called a Mergelyan set for H^∞ . The object of this paper is to give a new and constructive proof of the following result:

THEOREM. Let S be a relatively closed subset of D . Assume that the clusterpoints of S on the unit circle $T = \{z: |z| = 1\}$ which are not in the nontangential closure of S , has zero linear measure. Then S is a Mergelyan set for H^∞ .

A point $\eta \in T$ is said to be in the nontangential closure of S if η is a limit point of $S \cap D(\lambda, \eta)$ for some $\lambda > 0$ where $D(\lambda, \eta) = \{z \in D: |z - \eta| < \lambda(1 - |z|)\}$.

The theorem above solves a problem raised by J. Detraz [1]. (She also proved the converse of the theorem.)

The first proof of it (due to A. M. Davie) was based on functional analysis, and is rather short and elegant. See [5] for details. But from a constructive point of view, the proof is not very satisfactory. So the main reason for introducing this new and longer proof, is that it gives a rather explicit construction of the approximating polynomials $\{p_n\}_{n=1}^\infty$.

This theorem is part of a more general problem in simultaneous approximation introduced to me by Professor L. Rubel. (See [4], [5], and [6].)

Proof of the theorem. We denote by S^2 the extended complex plane. If $B \subset S^2$, and g is a function defined on B , then we define $\|g\|_B = \sup \{|g(z)|, z \in B\}$. Various absolute constants will be denoted by A_1, A_2, \dots .

Let f and S be as in the above theorem. We assume $\|f\|_D \leq 1$.

If $E \subset T$ is measurable, $|E|$ denotes its linear measure. Let $D(\lambda, \rho)$ be as above. By a well known result of Fatou, ([3], page 34) there is a set $F \subset T$, with $|F| = 0$, such that

$$(1) \quad \lim_{\substack{z \rightarrow \rho \\ z \in D(\lambda, \rho)}} f(z) \stackrel{\text{def}}{=} f^*(\rho)$$

exists for all $\rho \in T \setminus F$ and all $\lambda > 0$. Let E denote the limitpoints of S on T . By (1) and our hypothesis on S , the continuous extension of f to $S \cup E$, coincides with f^* almost everywhere on E . (With respect to linear measure.)

By approximating as in §1 in [7], we can also assume that f^* is continuous on $T \setminus E$. But then f^* (considered as a function on T) is continuous on $T \setminus (E \setminus B)$, where B denotes the interior of E relative to T .

The approximation of f is now done in two steps. If $|E \setminus B| = 0$, the first step can be omitted. So we assume $|E \setminus B| > 0$.

The idea is to modify Vitushkin's scheme for rational approximation (see [2], page 210) to be suitable for our problem. We shall construct functions f_n , $n = 1, 2, \dots$ such that $g = f - \sum_1^\infty f_n$ has the following two properties if $\varepsilon > 0$ is given in advance:

- (2) $\|f - g\|_S < \varepsilon$.
 (3) g is holomorphic in D and continuous in $D \cup T \setminus E_0$, where $E_0 \subset E \setminus B$ is compact and $|E_0| = 0$. Also $\|g\|_D \leq A_1$.

Before we construct $\{f_n\}_{n=1}^\infty$, let us indicate the second and last step in the proof. Denote by $A(D)$, all uniformly continuous analytic functions in D . We choose functions a and b from $A(D)$ such that there is a neighborhood V of E_0 in C with the following property

$$(4) \quad \begin{cases} |a - f| < \varepsilon \cdot 2^{-1} & \text{on } S \cap V. \\ \max \{ \|a\|_D, \|b\|_D \} \leq 2. \\ b = 0 & \text{on } E_0 \text{ and if } z \in S, \text{ then} \\ |a(z) - f(z)| > 2\varepsilon \implies |1 - b(z)| < \varepsilon. \end{cases}$$

The construction of a and b can be done as in [3], page 80. We partition E_0 into finitely many subsets E_1, \dots, E_n with pairwise disjoint neighborhoods W_1, \dots, W_n , such that there are complex numbers λ_j , $1 \leq j \leq n$ with $|f - \lambda_j| < \varepsilon 2^{-2}$ on each E_j . For each j we argue as on page 80 in [3] and find $b_j \in A(D)$ with $b_j = 0$ on E_j and $\operatorname{Re} b_j < 0$ on $D \setminus E_j$. Then we define

$$a = \sum_1^n \lambda_j e^{Nbj}$$

where N is a sufficiently large number. Then $|a - f| < \varepsilon 2^{-1}$ in some neighborhood V of E_0 , and we choose $b \in A(D)$ with $b = 0$ on E_0 , $\|b\|_D \leq 1$ and $|1 - b| < \varepsilon$ on $D \setminus V$.

Consider now $h = b(g - a) + a$. Then $h \in A(D)$ and since $g - h = (g - a)(1 - b)$, we have from (2), (3), and (4) above that $\|g - h\|_S < \max\{(A_1 + 2)\varepsilon, 5\varepsilon\}$. But then $\|f - h\|_S < A_2\varepsilon$, and we also have $\|h\|_D \leq A_3$. Since $h \in A(D)$, this function is easy to approximate uniformly on D by polynomials, and this completes the second step.

We turn to the construction of $\{f_n\}$.

Let $\varepsilon > 0$ be given. We choose pairwise disjoint discs $A_n = A(w_n, \delta_n)$, $n = 1, 2, \dots$ and constants t, M , and λ with the following properties

- (5) $w_n \in E \setminus B$ and there are numbers η_n with $\delta_n < \eta_n < 2\delta_n < t$ such that $[(A(w_n, \eta_n) \setminus A(w_n, \delta_n)) \cap T] \subset T \setminus (E \setminus B)$.
- (6) $|E \setminus (B \cup (\bigcup_n A_n))| = 0$.
- (7) $\sup\{|f(z) - f(\rho)|, z, \rho \in A(w_n, 2\delta_n) \cap D(\lambda, w_n)\} < \varepsilon$.
- (8) $\sup\{|f(z) - f(\rho)|, z, \rho \in S, |z - \rho| \leq Mt\} < \varepsilon$.

More details about how t, λ , and M depend on ε , will be given below. The important thing is that they are given in advance, before $\{A_k\}$ is constructed. The existence of $\{A_k\}$ satisfying (5)–(8) follows from Fatou’s result (1) about nontangential limits mentioned above, and the fact that $E \setminus B$ is totally disconnected. To obtain (8) we use that $f|_S$ is uniformly continuous.

We now choose numbers r_n with $\delta_n < r_n < \eta_n$, and smooth functions φ_n with support in $A(w_n, r_n)$ such that we also have for all n

$$(9) \quad 0 \leq \varphi_n \leq 1, \quad \varphi_n \equiv 1 \text{ near } A_n, \text{ and} \\ \left\| \frac{\partial \varphi_n}{\partial \bar{z}} \right\|_c \leq A_2(\eta_n - \delta_n)^{-1}.$$

We extend f to C by the equation $f(z) = f(1/\bar{z})$. Define functions G_n , $n = 1, 2, \dots$ by

$$(*) \quad G_n(\rho) = T_{\varphi_n} f(\rho) = \frac{1}{\pi} \iint \frac{f(z) - f(\rho)}{z - \rho} \frac{\partial \varphi_n}{\partial \bar{z}} dx dy \\ = \varphi_n(\rho) \cdot f(\rho) + \frac{1}{\pi} \iint \frac{f(z)}{z - \rho} \frac{\partial \varphi_n}{\partial \bar{z}} dx dy.$$

For basic properties of the T_φ -operator we refer to [2], page

30. These properties gives that G_n is analytic in D and continuous wherever f is. Also $f - G_n$ is analytic near $\Delta_n \cap (E \setminus B)$. Since $\|f\|_c \leq 1$, (9) shows that

$$(10) \quad \|G_n\|_c \leq A_3.$$

We also claim that the constants t , M , and λ can be chosen so that

$$(11) \quad \|G_n\|_s < A_4 \varepsilon.$$

To see this, note first that since G_n is analytic in $S^2 \setminus \Delta(w_n, t)$ and $G_n(\infty) = 0$, an easy application of Schwartz lemma shows that $|G_n(\rho)| < A(M - 1)^{-1} < \varepsilon$ if M is sufficiently large, and $|\rho - w_n| > Mt$. This does not contradict (8) if t is sufficiently small. If we use (8), we see that if $p \in S$ and $|\rho - w_n| \leq Mt$, then we can estimate the first integral in (*) so that

$$(12) \quad |G_n(\rho)| \leq A_5 \varepsilon + K(\lambda)$$

where $K(\lambda)$ is a constant tending to zero if $\lambda \rightarrow \infty$. It is easy to see that we can choose λ depending on ε , but not on n , such that $K(\lambda) < \varepsilon$. (The constant $K(\lambda)$ comes from integration over those $z \in \Delta(w_n, r_n)$ where $|f(z) - f(\rho)| > 2\varepsilon$).

We can now write f_n as

$$f_n = G_n - H_n$$

where H_n is a rational function with poles only in $\Delta(w_n, \delta_n) \setminus \bar{D}$ such that

$$\begin{aligned} \|H_n\|_s &\leq \|G_n\|_s \leq \varepsilon A_6 \\ \|H_n\|_D &\leq \|G_n\|_D \leq A_7 \\ \|H_n - G_n\|_{S^2 \setminus \Delta(w_n, \eta_n)} &< \varepsilon 2^{-n}. \end{aligned}$$

The existence of $\{H_n\}$ is not quite trivial. If we map $(S^2 \setminus \Delta(w_n, r_n)) \cup D$ conformally onto D , $S \cup (S^2 \setminus \Delta(w_n, \eta_n))$ is mapped to a Farrel set (see [5] for definition) in D , and the existence of H_n , $n = 1, 2, \dots$, now follows from the main result in [5]. This completes the proof. In fact, the set E_0 in (3) above, is simply $(T \setminus (B \cup (\bigcup_n \Delta_n)))$.

Concluding remark. It is easy to verify that this proof applies to more general planar domains.

REFERENCES

1. J. Detraz, *Algebres de fonctions analytique dans le disque*, Ann. Sci. Ecole Norm. Sup., **3** (1970), 313-352.

2. T. W. Gamelin, *Uniform Algebras*, Prentice Hall, Englewood Cliffs, N.J., 1969.
3. K. Hoffman, *Banach Spaces of Analytic Functions*, Prentice Hall, Englewood Cliffs, N.J., 1962.
4. L. A. Rubel, *Symposium on Complex Analysis*, Canterbury, 1973. Edited by J. Clunie and W. K. Hayman. Cambridge University Press.
5. A. Stray, *Pointwise bounded approximation by functions satisfying a side conditions*, Pacific J. Math., **51** (1974).
6. ———, *Characterization of Mergelyan sets*, Proc. Amer. Math. Soc., **44**, number 2, (1974).
7. ———, *Approximation and interpolation*, Pacific J. Math., **40**, number 2, (1972).

Received January 18, 1977 and in revised form August 24, 1977.

A. D. H.
P. O. BOX 607
4601 KR. SAND
NORWAY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

C. W. CURTIS

University of Oregon
Eugene, OR 97403

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
OSAKA UNIVERSITY

Gerald Arthur Anderson, <i>Computation of the surgery obstruction groups</i> $L_{4k}(1; Z_p)$	1
R. K. Beatson, <i>The degree of monotone approximation</i>	5
Sterling K. Berberian, <i>The character space of the algebra of regulated functions</i>	15
Douglas Michael Campbell and Jack Wayne Lamoreaux, <i>Continua in the plane with limit directions</i>	37
R. J. Duffin, <i>Algorithms for localizing roots of a polynomial and the Pisot Vijayaraghavan numbers</i>	47
Alessandro Figà-Talamanca and Massimo A. Picardello, <i>Functions that operate on the algebra $B_0(G)$</i>	57
John Erik Fornæss, <i>Biholomorphic mappings between weakly pseudoconvex domains</i>	63
Andrzej Granas, Ronald Bernard Guenther and John Walter Lee, <i>On a theorem of S. Bernstein</i>	67
Jerry Grossman, <i>On groups with specified lower central series quotients</i>	83
William H. Julian, Ray Mines, III and Fred Richman, <i>Algebraic numbers, a constructive development</i>	91
Surjit Singh Khurana, <i>A note on Radon-Nikodým theorem for finitely additive measures</i>	103
Garo K. Kiremidjian, <i>A Nash-Moser-type implicit function theorem and nonlinear boundary value problems</i>	105
Ronald Jacob Leach, <i>Coefficient estimates for certain multivalent functions</i>	133
John Alan MacBain, <i>Local and global bifurcation from normal eigenvalues. II</i>	143
James A. MacDougall and Lowell G. Sweet, <i>Three dimensional homogeneous algebras</i>	153
John Rowlay Martin, <i>Fixed point sets of Peano continua</i>	163
R. Daniel Mauldin, <i>The boundedness of the Cantor-Bendixson order of some analytic sets</i>	167
Richard C. Metzler, <i>Uniqueness of extensions of positive linear functions</i>	179
Rodney V. Nillsen, <i>Moment sequences obtained from restricted powers</i>	183
Keiji Nishioka, <i>Transcendental constants over the coefficient fields in differential elliptic function fields</i>	191
Gabriel Michael Miller Obi, <i>An algebraic closed graph theorem</i>	199
Richard Cranston Randell, <i>Quotients of complete intersections by C^* actions</i>	209
Bruce Reznick, <i>Banach spaces which satisfy linear identities</i>	221
Bennett Setzer, <i>Elliptic curves over complex quadratic fields</i>	235
Arne Stray, <i>A scheme for approximating bounded analytic functions on certain subsets of the unit disc</i>	251
Nicholas Th. Varopoulos, <i>A remark on functions of bounded mean oscillation and bounded harmonic functions. Addendum to: "BMO functions and the $\bar{\partial}$-equation"</i>	257
Charles Irvin Vinsonhaler, <i>Torsion free abelian groups quasi-projective over their endomorphism rings. II</i>	261
Thomas R. Wolf, <i>Characters of p'-degree in solvable groups</i>	267
Toshihiko Yamada, <i>Schur indices over the 2-adic field</i>	273