

Pacific Journal of Mathematics

**A REMARK ON FUNCTIONS OF BOUNDED MEAN
OSCILLATION AND BOUNDED HARMONIC FUNCTIONS.
ADDENDUM TO: "BMO FUNCTIONS AND THE $\bar{\partial}$ -EQUATION"**

NICHOLAS TH. VAROPOULOS

A REMARK ON FUNCTIONS OF BOUNDED MEAN OSCILLATION AND BOUNDED HARMONIC FUNCTIONS

N. TH. VAROPOULOS

This note is an Addendum to [7]; it is written, however, in such a way that it can be read independently.

We recall that μ a positive measure on the upper half space $\mathbf{R}_+^{n+1} = \{(x, y); x \in \mathbf{R}^n, y > 0\}$ (the notations we use are those of [6]) is called a Carleson measure if

$$\mu\{x \in I, 0 < y < h\} \leq Ch^n$$

for all I hypercube in \mathbf{R}^n of side length equal to h (cf. [6] VII 44, [1] [7]). In this note we shall prove the following

THEOREM 1. *Let u be a real bounded harmonic function in \mathbf{R}_+^{n+1} such that $\|u\|_\infty \leq 1$ then there exists $v \in C^\infty(\mathbf{R}_+^{n+1})$ a real bounded smooth function such that $|\nabla v| d \text{Vol}$ is a Carleson measure in \mathbf{R}_+^{n+1} and such that $\|u - v\|_\infty \leq 1 - \varepsilon_0$ for some numerical constant ε_0 ($0 < \varepsilon_0 < 1$).*

∇v denotes of course the gradient of v in \mathbf{R}_+^{n+1} .

REMARK. Note that $|\nabla u| d \text{Vol}$ is not in general a Carleson measure (cf. [5]).

THEOREM 2. *Let $f \in \text{BMO}(\mathbf{R}^n)$ then there exists $F \in C^\infty(\mathbf{R}_+^{n+1})$ such that*

- (i) $F(x, y) \xrightarrow{y \rightarrow 0} f(x) \quad x \in \mathbf{R}^n \text{ p.p.}$
- (ii) $\sup_{y > 0} |F(x, y)| \in L^1_{\text{loc}}(\mathbf{R}^n; dx)$
- (iii) $|\nabla F| d \text{Vol}$ is a Carleson measure in \mathbf{R}_+^{n+1} .

Conversely if $f \in L^1_{\text{loc}}(\mathbf{R}^n)$ is such that there exists some $F \in C^1(\mathbf{R}_+^{n+1})$ that satisfies (i), (ii), and (iii) then f is a BMO function.

(For the definition of BMO and background cf. [2].)

In functional terms the above theorem means that the space BMO is the restriction space (i.e., quotient space) of an appropriate space of C^∞ functions in the upper half space (cf. [3] for analogous results).

Both the above theorems can be generalized in other contexts,

e.g., the complex ball or more generally strictly pseudoconvex domains in C^n . Indeed it was in these contexts that theorems of the above type turned out to be useful for the first time cf. [7].

We shall give the proof of both theorems when $n = 1$, the general case is identical. The proof of theorem is based on a construction that was introduced in [1] by L. Carleson and which has proved to be, a very powerful tool in the study of bounded harmonic functions. A number of modifications of the above construction due to D. Sarason and D. Marshall can be used to simplify matters.

Let u be a real bounded harmonic function in R_+^2 and let us suppose that $\|u\|_\infty \leq 1$. Let us denote by $S = \{0 < x < 1, 0 < y < 1\}$ the unit square in R_+^2 .

$$\begin{aligned} A &= \{s \in S; |u(s)| < \alpha\} \\ B_1 &= \{s \in S; u(s) > \beta\} \\ B_2 &= \{s \in S; u(s) < -\beta\} \end{aligned}$$

and $B = B_1 \cup B_2$, where $0 < \alpha < \beta < 1$ are two numbers that will be chosen appropriately. We have then

LEMMA (L. Carleson and D. Marshall). *For an appropriate choice of α and β there exists $\Gamma \subset S$ a closed subset which is the union of countably many horizontal and vertical linear segments in S that satisfies the following two conditions:*

- (i) Γ separates A from B in S i.e., if $\sigma: [0, 1] \rightarrow S$ is a continuous curve s.t. $\sigma(0) \in A$ and $\sigma(1) \in B$ then $\exists 0 \leq t \leq 1$ s.t. $\sigma(t) \in \Gamma$.
- (ii) The measure μ_Γ induced by the arc length of Γ is a Carleson measure. (μ_Γ is defined by $\mu_\Gamma(\Omega) = \text{total length of the intersection of the linear segments of } \Gamma \text{ with } \Omega$, where Ω is an arbitrary open set of S .)

The above lemma is explicitly proved in [4], therefore no proof will be given here. (The proof is far from easy however, it essentially contains the hard part of the Corona theorem!). The best reference (i.e., the most elementary approach) for the above lemma is to be found in D. Marshall's thesis UCLA (1976). In that reference he avoids the use of Hall's lemma.

Proof of Theorem 1. Let $S \setminus \Gamma = \bigcup_{n=1}^\infty S_n$ be the decomposition of $S \setminus \Gamma$ into its connected components, and let us define F a function on $S \setminus \Gamma$ as following

$$\begin{aligned} F(x) &= 1 \quad \forall x \in S_n \quad n \geq 1 \quad \text{if} \quad S_n \cap B_1 \neq \emptyset \\ F(x) &= -1 \quad \forall x \in S_n \quad n \geq 1 \quad \text{if} \quad S_n \cap B_2 \neq \emptyset \\ F(x) &= 0 \quad \forall x \in S_n \quad n \geq 1 \quad \text{if} \quad S_n \cap B = \emptyset. \end{aligned}$$

It is then clear from the lemma that F is well defined and that $|\nabla F|$ taken in the sense of distribution theory is a Carleson measure. It is also evident from the definition that $\|u - F\|_\infty \leq \max[1 - \beta, \beta]$ in S . To obtain the required function in S it suffices to smooth out appropriately the above function. The passage to the whole of \mathbf{R}_+^n is obvious. Observe also that in Theorem 1 the function v can be chosen such that $\lim v(x, y)$ as $y \rightarrow 0$ exists for almost all x .

Proof of Theorem 2. Using a very simple argument that involves Banach spaces and a geometric series (which anyone who has seen a proof of the closed graph theorem can supply) we see that Theorem 1 implies the first part of Theorem 2 if $f \in L^\infty(\mathbf{R}^n)$. The general case follows then from Theorem 1.1.1 of [7]. The inverse is contained in [7] pt. 1.

One question (of mild interest) that remains open is whether ε_0 in Theorem 1 can be chosen to be an arbitrary positive number such that $0 < \varepsilon_0 < 1$. *Added in proof:* J. Garnett has found a way to do that.

REFERENCES

1. L. Carleson, *Interpolation by bounded analytic functions and the Corona problem*, Ann. of Math., **76** (1962), 547-559.
2. C. Fefferman and E. M. Stein, *H^p spaces of several variables*, Acta Math., **129** (1972), 137-193.
3. E. Galiardo, *Caratterizzazioni della trace sulla frontiera relative ad alcune classi di funzioni in n variabili*, Rend. Sem. Mat. Univ. Padova, **27** (1957), 284-305.
4. D. Marshall, *Subalgebras of L^∞ containing H^∞* , Acta Math., **137**, 91-98.
5. W. Rudin, *The radial variation of analytic functions*, Duke Math. J., **22** (1955), 235-242.
6. E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Univ. Press (1970).
7. N. Th. Varopoulos, *BMO functions and the $\bar{\partial}$ equation*, Pacific J. Math., **71** (1) (1977), 221-273.

Received July 15, 1977.

MATHEMATIQUE
UNIVERSITE DE PARIS-SUD
CENTRE D'ORSAY, FRANCE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

C. W. CURTIS

University of Oregon
Eugene, OR 97403

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
OSAKA UNIVERSITY

| | |
|---|-----|
| Gerald Arthur Anderson, <i>Computation of the surgery obstruction groups</i> $L_{4k}(1; Z_p)$ | 1 |
| R. K. Beatson, <i>The degree of monotone approximation</i> | 5 |
| Sterling K. Berberian, <i>The character space of the algebra of regulated functions</i> | 15 |
| Douglas Michael Campbell and Jack Wayne Lamoreaux, <i>Continua in the plane with limit directions</i> | 37 |
| R. J. Duffin, <i>Algorithms for localizing roots of a polynomial and the Pisot Vijayaraghavan numbers</i> | 47 |
| Alessandro Figà-Talamanca and Massimo A. Picardello, <i>Functions that operate on the algebra $B_0(G)$</i> | 57 |
| John Erik Fornæss, <i>Biholomorphic mappings between weakly pseudoconvex domains</i> | 63 |
| Andrzej Granas, Ronald Bernard Guenther and John Walter Lee, <i>On a theorem of S. Bernstein</i> | 67 |
| Jerry Grossman, <i>On groups with specified lower central series quotients</i> | 83 |
| William H. Julian, Ray Mines, III and Fred Richman, <i>Algebraic numbers, a constructive development</i> | 91 |
| Surjit Singh Khurana, <i>A note on Radon-Nikodým theorem for finitely additive measures</i> | 103 |
| Garo K. Kiremidjian, <i>A Nash-Moser-type implicit function theorem and nonlinear boundary value problems</i> | 105 |
| Ronald Jacob Leach, <i>Coefficient estimates for certain multivalent functions</i> | 133 |
| John Alan MacBain, <i>Local and global bifurcation from normal eigenvalues. II</i> | 143 |
| James A. MacDougall and Lowell G. Sweet, <i>Three dimensional homogeneous algebras</i> | 153 |
| John Rowlay Martin, <i>Fixed point sets of Peano continua</i> | 163 |
| R. Daniel Mauldin, <i>The boundedness of the Cantor-Bendixson order of some analytic sets</i> | 167 |
| Richard C. Metzler, <i>Uniqueness of extensions of positive linear functions</i> | 179 |
| Rodney V. Nillsen, <i>Moment sequences obtained from restricted powers</i> | 183 |
| Keiji Nishioka, <i>Transcendental constants over the coefficient fields in differential elliptic function fields</i> | 191 |
| Gabriel Michael Miller Obi, <i>An algebraic closed graph theorem</i> | 199 |
| Richard Cranston Randell, <i>Quotients of complete intersections by C^* actions</i> | 209 |
| Bruce Reznick, <i>Banach spaces which satisfy linear identities</i> | 221 |
| Bennett Setzer, <i>Elliptic curves over complex quadratic fields</i> | 235 |
| Arne Stray, <i>A scheme for approximating bounded analytic functions on certain subsets of the unit disc</i> | 251 |
| Nicholas Th. Varopoulos, <i>A remark on functions of bounded mean oscillation and bounded harmonic functions. Addendum to: "BMO functions and the $\bar{\partial}$-equation"</i> | 257 |
| Charles Irvin Vinsonhaler, <i>Torsion free abelian groups quasi-projective over their endomorphism rings. II</i> | 261 |
| Thomas R. Wolf, <i>Characters of p'-degree in solvable groups</i> | 267 |
| Toshihiko Yamada, <i>Schur indices over the 2-adic field</i> | 273 |