

Pacific Journal of Mathematics

PRINCIPAL IDEAL AND NOETHERIAN GROUPS

S. FEIGELSTOCK AND Z. SCHLUSSEL

PRINCIPAL IDEAL AND NOETHERIAN GROUPS

S. FEIGELSTOCK AND Z. SCHLUSSEL

Let Π be a ring property. An additive group G is said to be an (associative) strongly Π -group if G is not nil, and if every (associative) ring R with additive group G such that R is not a zeroing has property Π . The (associative) strongly principal ideal groups, and the (associative) strongly Noetherian groups are classified for groups which are not torsion free. Some results are also obtained for the torsion free case.

(i) All groups considered here are abelian, with addition the group operation. Rings are not necessarily associative.

Let π be a ring property. A group G is said to be an (associative) π -group, denoted by $(A)\pi$ -group, if there exists an (associative) ring R with additive group G such that R is not a zeroing, and R has property π . G is an (associative) strongly π -group, denoted by $(A)S\pi$ -group, if G is an $(A)\pi$ -group, and every (associative) ring with additive group G which is not the zeroing on G has property π .

If the only (associative) ring with additive group G is the zeroing, then G is said to be an (associative) nil group, denoted by (A) nil group.

The two ring properties π considered in this paper are:

1. every two-sided ideal is principal, denoted by PI ,
2. every two-sided ideal is finitely generated, denoted by N .

In (ii) a complete characterization of the torsion $(A)SPI$ groups will be given. It will be shown that there are no mixed $(A)SPI$ groups. Some results concerning torsion free $(A)SPI$ groups will be obtained. In (iii), the torsion, and mixed SN groups will be completely characterized. Some results concerning torsion free SN groups will be given.

(ii) If X is a nonempty subset of a group or ring, $\langle X \rangle$ denotes the additive subgroup generated by X , and $\langle X \rangle$ denotes the ideal generated by X .

If $G = G_1 \oplus G_2$ is a group, π_{G_i} is the natural projection of G on G_i , for $i = 1, 2$.

LEMMA 1. Let $G = H \oplus K$, $H \neq 0$, $K \neq 0$, be an $ASPI$ -group. Then H and K are either both cyclic or both A nil.

Proof. Suppose that H is not A nil. Let S be an associative ring on H which is not the zeroring on H , and let T be the zeroring on K . The ring direct sum $R = S \oplus T$ is an associative ring on G , which is not the zeroring on G . T is an ideal in R , and hence $T = \langle x \rangle$. Clearly $K = \langle x \rangle$. Therefore K is not A nil. The above argument, interchanging the roles of H and K , yields that H is cyclic.

COROLLARY. *Let $G = H \oplus K$, $H \neq 0$, $K \neq 0$ be an SPI-group. Then H and K are cyclic.*

Proof. It suffices to negate that H and K are both A nil. Suppose this is so. Let R be a ring on G which is not the zeroring on G .

(1) Suppose that $R^2 \subseteq K$. There exist $h_0 \in H$, $k_0 \in K$ such that $R = \langle h_0 + k_0 \rangle$. Let $h \in H$. Since $h \in R$, there exists an integer n , and $x \in R^2$ such that $h = n(h_0 + k_0) + x$. However, $x \in K$. Hence $h = nh_0$, and H is cyclic, contradicting the fact that H is A nil.

(2) Suppose that $R^2 \not\subseteq K$. For all $g_1, g_2 \in G$, define $g_1 \times g_2 = \pi_H(g_1 g_2)$. Then $S = (G, \times)$ is a not necessarily associative ring on G , which is not the zeroring on G , satisfying $S^2 \subseteq H$. The argument employed in (1) yields that K is cyclic, contradicting the fact that K is A nil.

THEOREM 1. *Let G be a nonzero torsion group. The following are equivalent:*

- (1) *Either G is cyclic, or $G \cong Z(p) \oplus Z(p)$ for a prime p .*
- (2) *G is SPI.*
- (3) *G is ASPI.*

Proof. (1) \Rightarrow (2): Nontrivial cyclic groups are clearly SPI. Suppose that $G = \langle x_1 \rangle \oplus \langle x_2 \rangle$ with $|x_i| = p$ a prime, $i = 1, 2$. Let R be a ring on G which is not the zeroring on G , and let l be a proper ideal in R . Then $|l| = 0$ or p , and hence l is generated by a single element. It therefore suffices to show that R is generated by a single element. We may assume that $R \neq \langle x_1 \rangle$, and that $R \neq \langle x_2 \rangle$. Hence $\langle x_i \rangle = \langle x_i \rangle$ for $i = 1, 2$. This implies that

$$x_i x_j = \begin{cases} k_i x_i, 0 < k_i < p, & \text{if } i = j, i = 1, 2 \\ 0 & \text{if } i \neq j, i, j = 1, 2. \end{cases}$$

Put $l = \langle x_1 + x_2 \rangle$. Let r, s be integers such that $rk_1 + sp = 1$. Then $rx_1(x_1 + x_2) = rk_1 x_1 = (1 - sp)x_1 = x_1$. Hence $x_1 \in l$, and so $(x_1 + x_2) - x_1 = x_2 \in l$. Therefore $l = R$.

(2) \Rightarrow (3): Let G be a torsion SPI group. It suffices to show that G

admits an associative, nonzero multiplication. If G is indecomposable then $G \simeq Z(p^n)$, p a prime, and n a positive integer or ∞ [1, Corollary 27.4]. If $n = \infty$ then G is nil. If n is a positive integer then G admits a ring structure isomorphic to Z_p^n the ring of integers modulo p^n . If G is decomposable, then by the Corollary to Lemma 1, $G \simeq Z(n) \oplus Z(m)$. Hence G admits a ring structure isomorphic to the ring direct sum $Z_n \oplus Z_m$.

(3) \Rightarrow (1): Suppose that G is *ASPI*. If G is indecomposable, then $G \simeq Z(p^k)$, p a prime, $1 \leq k \leq \infty$, [1, Corollary 27.4]. However $Z(p^\infty)$ is *A nil* [3, Satz 1, and Zusatz]. Hence G is cyclic. By Lemma 1 and [1, Theorem 120.3] we may assume $G = (x_1) \oplus (x_2)$ with $|x_i| = n_i, i = 1, 2$. If $(n_1, n_2) = 1$, G is cyclic; otherwise, let p be a prime divisor of (n_1, n_2) . Then $G = (y_1) \oplus (y_2) \oplus H$, with $|y_i| = p^{m_i}, i = 1, 2$, and $1 \leq m_1 \leq m_2$. Since $(y_1) \oplus (y_2)$ is neither cyclic nor *A nil*, $H = 0$ by Lemma 1.

Now let R be the ring on G with multiplication defined by $y_i y_j = p^{m_2-1} y_2$ for $i, j = 1, 2$. Then R is an associative ring on G which is not a zeroring, so $R = \langle s_1 y_1 + s_2 y_2 \rangle$ for some $s_1, s_2 \in Z$. Every $x \in R$ has the form $x = k_x s_1 y_1 + (k_x s_2 + m_x p^{m_2-1}) y_2$, for some $k_x, m_x \in Z$. In particular, $y_1 = k_{y_1} s_1 y_1$, and $y_2 = (k_{y_2} s_2 + m_{y_2} p^{m_2-1}) y_2$. Hence if $m_2 > 1, k_{y_1} s_1 \equiv 1 \pmod{p}$, $k_{y_2} s_2 + m_{y_2} p^{m_2-1} \equiv 1 \pmod{p}$, so $p \nmid k_{y_1}$ and $p \nmid s_2$. However $k_{y_1} s_2 + m_{y_1} p^{m_2-1} \equiv 0 \pmod{p}$, so that either $p \mid k_{y_1}$, or $p \mid s_2$. This is a contradiction, so $m_2 = 1 = m_1$.

THEOREM 2. *There are no mixed ASPI-groups.*

Proof. Let G be a mixed *ASPI*-group. G is decomposable, [1, Corollary 27.4], so by Lemma 1, $G = H \oplus K, H \neq 0, K \neq 0$, with H and K either both cyclic or both *A nil*.

(1) Suppose that H and K are both *A nil*. There are no mixed *A nil* groups [4, hilfssatz 9] so we may assume that H is a torsion group, and that K is torsion free. Let R be an associative ring on G , such that R is not a zeroring. Clearly H is an ideal in R , and so $H = \langle h \rangle$. Let $m = |h|$, so $mh = 0$. There are no nontrivial, bounded *A nil*-groups [3, Satz 1 and Zusatz], a contradiction.

(2) Suppose that $H = (x)$, and $K = (e)$ with $|x| = n < \infty$, and $|e| = \infty$. Let R be the ring on G with multiplication induced by $x^2 = xe = ex = 0, e^2 = ne$. Clearly, R is an associative ring on G , and R is not a zeroring. Hence $R = \langle sx + te \rangle, s, t \in Z$. Every $y \in R$ is of the form $y = m_y sx + (m_y + u_y n) te, m_y, u_y \in Z$. In particular, $(m_e + u_e n) t = 1$. Hence $t = \pm 1$. Therefore $m_x + u_x n = 0$, so that $n \mid m_x$. However $x = m_x sx = 0$, a contradiction.

THEOREM 3. *Let G be a torsion free, ASPI-group. Then G is either indecomposable, or the direct sum of two *A nil*-groups.*

Proof. By Lemma 1, it suffices to negate that $G = (x_1) \oplus (x_2)$, $x_i \neq 0$, $i = 1, 2$. Suppose this is so. Let R be the ring on G with multiplication induced by

$$x_i x_j = \begin{cases} 3x_i & \text{if } i = j, i = 1, 2 \\ 0 & \text{if } i \neq j, i, j = 1, 2 \end{cases}$$

Then $R = \langle k_1 x_1 + k_2 x_2 \rangle$, $k_i \in \mathbb{Z}$, $k_i \neq 0$, $i = 1, 2$. Every $x \in R$ is of the form

$$x = (r_x + 3s_x)k_1 x_1 + (r_x + 3t_x)k_2 x_2, r_x, s_x, t_x \in \mathbb{Z}.$$

$r_{x_1} + 3s_{x_1} = \pm 1$, so that $r_{x_1} \equiv \pm 1 \pmod{3}$. However, $r_{x_1} + 3t_{x_1} = 0$, so that $r_{x_1} \equiv 0 \pmod{3}$, a contradiction.

COROLLARY. *Let G be a torsion free SPI-group. Then G is indecomposable.*

Proof. Theorem 3, and the Corollary to Lemma 1.

The above Corollary yields that if G is a torsion free SPI-group then either $G \cong \mathbb{Q}$, the group of rational numbers, or G is reduced, [1, Theorem 21.3].

THEOREM 4. *Let G be a nonzero torsion group. The following are equivalent:*

- (1) G is bounded.
- (2) G is an API-group.
- (3) G is a PI-group.

Proof. (1) \Rightarrow (2): Suppose that $nG = 0$, n a positive integer. Then $G \cong \bigoplus_{p|n} [\bigoplus_{\alpha_k} \mathbb{Z}(p^k)]$, p a prime with $p^k | n$, and α_k a cardinal number, [1, Theorem 17.3, and Theorem 8.4]. For each $p^k | n$, put $H_{p^k} = \bigoplus_{\alpha_k} \mathbb{Z}(p^k)$. Then $G \cong \bigoplus_{p^k | n} H_{p^k}$: There exists an associative unital PI-ring R_{p^k} on H_{p^k} , for all $p^k | n$, [1, Lemma 122.3]. The ring direct sum $R = \bigoplus_{p^k | n} R_{p^k}$ is an API-ring on G which is not the zero ring on G [5, Chapt. 4, Theorem 33].

(2) \Rightarrow (3): Obvious.

(3) \Rightarrow (1): Let R be a PI-ring on G which is not the zero ring on G . Then $R = \langle x \rangle$. Let $n = |x|$. Clearly, $nG = 0$.

COROLLARY. *Let G be a mixed group. If G is an (A)PI-group, then $T(G)$ (the torsion part of G) is bounded, and $T/T(G)$ is an (A)PI-group. Conversely, if $T(G)$ is bounded, and if there exists a unital (A)PI-ring on $G/T(G)$, then G is (A)PI.*

Proof. Let G be an $(A)PI$ -group, and let R be an $(A)PI$ -ring on G which is not a zeroring. Since $T(G)$ is an ideal in R , $T(G) = \langle x \rangle$, and $nT(G) = 0$, $n = |x|$. Clearly $R/T(G)$ is an $(A)PI$ -ring with identity on $G/T(G)$.

Suppose that $T(G)$ is bounded, and that there exists an $(A)PI$ -ring with identity on $G/T(G)$. Then $G \simeq T(G) \oplus G/T(G)$, [1, Theorem 100.1]. There exists an API -ring R_1 with identity on $T(G)$, [1, Lemma 122.3]. Let R_2 be a unital $(A)PI$ -ring on $G/T(G)$. Let R be the ring direct sum $R = R_1 \oplus R_2$, with e_i the identity of R_i , $i = 1, 2$. Let l be an ideal in R . Then $l = (l \cap R_1) \oplus (l \cap R_2)$. Since $l \cap R_1$ is an ideal in R_1 , $l \cap R_1 = \langle x_i \rangle$, $i = 1, 2$. Clearly, $\langle x_1 + x_2 \rangle \subseteq l$. However $x_i = e_i(x_1 + x_2) \in \langle x_1 + x_2 \rangle$ for $i = 1, 2$. Hence $l = \langle x_1 + x_2 \rangle$.

Additional information concerning PI -groups and the classification of π -groups for other ring properties π may be found in [2].

LEMMA 2. *If a group G is finitely generated, then G is SN.*

Proof. Obvious.

LEMMA 3. *Let a group $G = H \oplus K$, $H \neq 0$, $K \neq 0$, be SN. Then either G is finitely generated, or H and K are both nil.*

Proof. Suppose that H is not nil. Let S be a nonzeroring on H , and let T be the zeroring on K . The ring direct sum $R = S \oplus T$ is a ring on G which is not the zeroring, with ideal T . Let t_1, \dots, t_n be a finite set of generators for T . Then $K = \langle t_1, \dots, t_n \rangle$. This implies that K is not nil. The same argument, interchanging the roles of H and K , yields that H is finitely generated. Hence G is finitely generated.

THEOREM 5. *Let G be a non torsion free group. G is SN if and only if G is finitely generated.*

Proof. By Lemma 2, it suffices to show that if G is non torsion free and SN, then G is finitely generated.

(1) Suppose that G is a torsion group. If G is indecomposable then G is cyclic [1, Corollary 27.4 and Theorem 120.3]. We may assume, by Lemma 3, that $G = H \oplus K$, $H \neq 0$, $K \neq 0$, with H and K both nil. This implies that G is nil [1, Theorem 120.3]. A contradiction.

(2) Suppose that G is a mixed group. Then G is decomposable [1, Corollary 27.3]. By Lemma 3, it suffices to negate that $G = H \oplus K$, $H \neq 0$, $K \neq 0$, with H and K both nil. Suppose that this is so. By [1, Theorem 120.3] we may assume that H is a torsion group and that K is torsion free. However, by [1, Proposition 126.2] H is bounded and hence not nil, a contradiction.

COROLLARY. *Let G be an SN group. Then $T(G)$ and $G/T(G)$ are SN.*

Proof. If G is torsion free then the statement is trivial. Otherwise, G is finitely generated by Theorem 5, and so $T(G)$ and $G/T(G)$ are SN by Lemma 2.

A torsion-free SN-group need not be finitely generated; e.g. Q the group of rational numbers. However, we have the following:

THEOREM 6. *Let G be an SN-group. Then G is either indecomposable, or finitely generated.*

Proof. By Lemma 3, it suffices to negate that $G = H \oplus K$, $H \neq 0$, $K \neq 0$, with H and K both nil. Suppose this is so. Let R be a ring on G which is not a zeroing. Then $R = \langle x_1, \dots, x_n \rangle$, n a positive integer. Put $x_i = h_i + k_i$, $h_i \in H$, $k_i \in K$, $1 \leq i \leq n$.

(1) Suppose that $R^2 \subseteq K$. Let $h \in H$. Since $h \in R$, $h = \sum_{i=1}^n r_i(h_i + k_i) + x$, $r_i \in Z$, $1 \leq i \leq n$, $x \in R^2$. However $R^2 \subseteq K$. Hence $h = \sum_{i=1}^n r_i h_i$, and H is finitely generated. This contradicts the fact that H is nil.

(2) Suppose that $R^2 \not\subseteq K$. For $g_1, g_2 \in G$ define $g_1 \times g_2 = \pi_H(g_1 g_2)$. Then $S = (G, \times)$ is a ring on G which is not a zeroing satisfying $S^2 \subseteq H$. The above argument yields that K is finitely generated, contradicting the fact that K is nil.

REFERENCES

1. L. Fuchs, *Infinite Abelian Groups*, Academic Press, New York and London, vol. 1 (1970), vol. 2 (1973).
2. Z. Schlüssel, *On Additive Groups of Rings*, Master's thesis, Bar-Ilan University, 1976 (Hebrew with English summary).
3. T. Szele, *Zur theorie der zeroringe*, Math. Ann., **121** (1949), 242-248.
4. ———, *Gruppentheoretische beziehungen bei gewissen ringkonstruktionen*, Math. Zeitschr., **54** (1951), 168-180.
5. O. Zariski and P. Samuel, *Commutative Algebra*, vol. 1, Van-Nostrand, Princeton (1958).

Received December 28, 1976 and in revised form June 13, 1977.

BAR-ILAN UNIVERSITY
RAMAT-GAN, ISRAEL

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, CA 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. A. BEAUMONT
University of Washington
Seattle, WA 98105

R. FINN AND J. MILGRAM
Stanford University
Stanford, CA 94305

R. C. MOORE
University of California
Berkeley, CA 94720

ASSOCIATE EDITORS

R. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
SAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typewritten or offset-reproduced (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, duplicate, may be sent to any one of the four editors. Please classify according to the scheme of Math. Review Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Bartlett, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Printed at Jerusalem Academic Press, POB 2390, Jerusalem, Israel.

Copyright © 1978 Pacific Journal of Mathematics
All Rights Reserved

Mieczyslaw Altman, <i>General solvability theorems</i>	1
Denise Amar and Eric Amar, <i>Sur les suites d'interpolation en plusieurs variables</i>	15
Herbert Stanley Bear, Jr. and Gerald Norman Hile, <i>Algebras which satisfy a second order linear partial differential equation</i>	21
Marilyn Breen, <i>Sets in R^d having $(d - 2)$-dimensional kernels</i>	37
Gavin Brown and William Moran, <i>Analytic discs in the maximal ideal space of $M(G)$</i>	45
Ronald P. Brown, <i>Quadratic forms with prescribed Stiefel-Whitney invariants</i>	59
Gulbank D. Chakerian and H. Groemer, <i>On coverings of Euclidean space by convex sets</i>	77
S. Feigelstock and Z. Schlussek, <i>Principal ideal and Noetherian groups</i>	87
Ralph S. Freese and James Bryant Nation, <i>Projective lattices</i>	93
Harry Gingold, <i>Uniqueness of linear boundary value problems for differential systems</i>	107
John R. Hedstrom and Evan Green Houston, Jr., <i>Pseudo-valuation domains</i>	137
William Josephson, <i>Coallocation between lattices with applications to measure extensions</i>	149
M. Koskela, <i>A characterization of non-negative matrix operators on l^p to l^q with $\infty > p \geq q > 1$</i>	165
Kurt Kreith and Charles Andrew Swanson, <i>Conjugate points for nonlinear differential equations</i>	171
Shoji Kyuno, <i>On prime gamma rings</i>	185
Alois Andreas Lechicki, <i>On bounded and subcontinuous multifunctions</i>	191
Roberto Longo, <i>A simple proof of the existence of modular automorphisms in approximately finite-dimensional von Neumann algebras</i>	199
Kenneth Millett, <i>Obstructions to pseudoisotopy implying isotopy for embeddings</i>	207
William F. Moss and John Piepenbrink, <i>Positive solutions of elliptic equations</i>	219
Mitsuru Nakai and Leo Sario, <i>Duffin's function and Hadamard's conjecture</i>	227
Mohan S. Putcha, <i>Word equations in some geometric semigroups</i>	243
Walter Rudin, <i>Peak-interpolation sets of class C^1</i>	267
Elias Saab, <i>On the Radon-Nikodým property in a class of locally convex spaces</i>	281
Stuart Sui Sheng Wang, <i>Splitting ring of a monic separable polynomial</i>	293