

Pacific Journal of Mathematics

**METABELIAN GROUPS WITH AN IRREDUCIBLE
PROJECTIVE REPRESENTATION OF LARGE DEGREE**

FRANK RIMI DEMEYER

METABELIAN GROUPS WITH AN IRREDUCIBLE PROJECTIVE REPRESENTATION OF LARGE DEGREE

F. R. DEMEYER

The structure of a metabelian group admitting an irreducible projective complex representation whose degree is the square root of the order of the group studied.

In [7] a class of metabelian groups admitting a faithful irreducible projective representation was characterized. If \bar{G} is a finite group with irreducible projective representation T^* so that $T^*(1)^2 = [\bar{G}:1]$ then T^* is a faithful irreducible projective representation. Conversely, if G is abelian and T^* is a faithful irreducible projective representation then $T^*(1)^2 = [\bar{G}:1]$. We study metabelian groups \bar{G} admitting an irreducible projective representation T^* with $T^*(1)^2 = [\bar{G}:1]$. Given such a group one can construct, using Schur's theory, a central extension G of \bar{G} with center Z and an ordinary faithful irreducible representation T of G with $T(1)^2 = [G:Z]$ and $T^*(gZ) = T(g)$ for all $g \in G$ (see p. 358 of [1]). A group G with an irreducible representation T so that $T(1)^2 = [G:Z]$ is called a group of central type.

Groups of central type have been studied in several places (see for example [3] and [5]). We study metabelian groups \bar{G} admitting an irreducible projective complex representation whose degree is $\sqrt{[\bar{G}:1]}$ by studying the associated central extension G of central type using the methods of ordinary representation theory. In this note all groups are finite and all representations and characters are assumed to be taken over the complex numbers. All unexplained terminology and notation is as in [1].

I. An abelian group G is said to be of symmetric type in case $G = A \times A$ for some abelian group A . It has been shown that an abelian group G is a group of symmetric type if and only if G admits a faithful irreducible projective representation [4] if and only if G admits an irreducible projective representation T^* with $T^*(1)^2 = [G:1](2)$. For any group H let $Z(H)$ denote the center of H .

THEOREM 1. *Let G be a group of central type with center Z . Assume there is a faithful irreducible character ζ on G with $\zeta(1)^2 = [G:Z]$. If there is a normal subgroup H of G with $Z \subseteq H \subseteq G$*

and both G/H and H/Z abelian then

(1) H is a group of central type so $H/Z(H)$ is of symmetric type.

(2) There is a normal subgroup K of G with $H \subseteq K \subseteq G$ so that K/H is of symmetric type and $[G:K] = [Z(H):Z]$.

Proof. First assume G is a p -group, p a prime. By Schur's lemma $\zeta|_Z = \zeta(1)\phi$ for a faithful linear character ϕ on Z . Proceed by induction on $[Z(H):Z]$ to prove: If ψ is an extension of ϕ to $Z(H)$ then $\psi^H = \lambda(1)\lambda$ where $\lambda(1)^2 = [H:Z(H)]$, and thus H is a group of central type with center $Z(H)$. If $[Z(H):Z] = 1$ then H is a group with cyclic center Z such that H/Z is abelian. In this case it is easy to check that the commutator map induces a non-degenerate bilinear function from $H/Z \times H/Z$ to Z . It follows by Proposition 3 of [2] that H is of central type and H/Z is of symmetric type. Moreover $\phi^H = \lambda(e)\lambda$ for an irreducible character λ on H with $\lambda(1)^2 = [H:Z]$. If $Z(H) \neq Z$ let $x \in Z(H) - Z$ with $x^p \in Z$ and $x \cdot Z$ in the center of G/Z . We can find such an x because G is a p -group. Let $H_1 = \{y \in G \mid [x, y] = e\}$. By Theorem 3 of [3], $[G:H_1] = p$ and H_1 is a group of central type with center $Z(H_1) = Z\langle x \rangle$. If ψ is an extension of ϕ to $Z(H_1)$ then $\psi^{H_1} = \rho(e)\rho$ where $\rho(e)^2 = [H_1:Z(H_1)]$. If $X = \ker \psi = \ker \rho$ then X is a subgroup of $Z(H_1)$ so that $X \cap Z = \{e\}$ and H_1/X is a group of central type with a faithful irreducible character ρ . Clearly $Z(H)/X \subseteq Z(H/X)$. If $hX \in Z(H/X)$ then for any $k \in H$, $[h, k] \in X$. But H/Z is abelian so $[h, k] \in Z$, and $Z \cap X = \{e\}$. Thus $h \in Z(H)$ and $Z(H)/X = Z(H/X)$. We apply induction to the group H_1/X , and obtain H/X is a group of central type with center $Z(H)/X$. Thus H is a group of central type with center $Z(H)$ and $H/Z(H)$ is of symmetric type. Moreover, if ψ is an extension of ψ to $Z(H)$ then $\psi^H = \lambda(1)\lambda$ where $\lambda(1)^2 = [H:Z(H)]$.

To prove (2) let $K = \{y \in G \mid \lambda^y = \lambda\}$. Since G/H is abelian, $K \triangleleft G$. We proceed by induction on $[Z(H):Z]$ as in the proof of (1) to show (K, H, λ) is a fully ramified triple [5] or [6], i.e., there is an irreducible character ρ on K so that $\rho|_H = e\lambda$ where $e^2 = [K:H]$. This suffices by Theorem 1.1 of [5] to show K/H is of symmetric type. If $Z(H) = Z$ then $K = G$ and $\rho = \zeta$ works. If $Z(H) \neq Z$ let H_1, X, ψ, λ be as in the proof of (1). If $y \in G - H_1$ then $\lambda^y(x) = \lambda(yxy^{-1}) = \lambda(xz)$ where $z \in Z - \{e\}$. But $\lambda|_{Z(H_1)} = \lambda(e)\psi$ so $\lambda(xz) = [\lambda(e)\psi(x)]\psi(z) = \lambda(x)\phi(z) = \lambda(x)\phi(z)$. Since ϕ is faithful, $y \notin K$ and thus $K \subseteq H_1$. Since $X \cap Z = \{e\}$ we can apply the same argument in G/X to see K/X is the inertia group of λ in G/H . Apply induction to H_1/X . Then $(K/X, H/X, \lambda)$ is a fully ramified triple so (K, H, λ) is fully ramified. To prove $[G:K] = [Z(H):Z]$ we have from

Clifford's theory (p. 342 of [1]) $\rho|_H = e(\lambda_1 + \cdots + \lambda_n)$ where by relabeling we can let $\lambda_1 = \lambda$ where the λ_i are conjugate irreducible characters, and $n = [G:K]$. Thus $\rho(1)^2 = [G:Z] = e^2\lambda(1)^2n^2 = [G:K][K:H][H:Z(H)][Z(H):Z]$. Since $e^2 = [K:H]$, $\lambda(e)^2 = [H:Z(H)]$, and $n = [G:K]$ so we have $[G:K] = [Z(H):Z]$.

Now we show Theorem 1 is true without requiring G to be a p -group. By the hypothesis H is nilpotent so $H = \prod_p H_p$ where p are the primes such that $p|[G:1]$ and the H_p are the p -Sylow subgroups of H . Let G_p be a Sylow p -subgroup of G , then $Z(G_p) = G_p \cap Z$ and $Z(G_p) \subseteq H_p \subseteq G_p$. Moreover, G_p is a group of central type whose irreducible character of large degree is faithful. (Theorem 2 of [3].) Thus H_p is a group of central type with center $Z(H_p)$ and there is a normal subgroup K_p of G_p containing H_p with K_p/H_p of symmetric type and $[G_p:K_p] = [Z(H_p):Z(G_p)]$. By Theorem 2 of [3], H is a group of central type with center $Z(H)$, and if K is the normal subgroup of G which is the inverse image under the natural homomorphism of $\prod_p K_p/H_p$ in G/H then K/H is of symmetric type since each Sylow subgroup is and $[G:K] = \prod_p [G_p:H_p] = \prod_p [Z(H_p):Z(G_p)] = [Z(H):Z]$. This proves the theorem.

There is a central extension G of $A_4 \times X_3$ of order $2^3 \cdot 3^3$ which is of central type, G is not nilpotent, yet G satisfies the hypothesis of Theorem 1.

Theorem 1 does not hold if ζ is not faithful because the center of H does not behave properly. However, if G is a group of central type with center Z and irreducible character ζ with $\zeta(e)^2 = [G:Z]$ then the kernel X of ζ is contained in Z so if one is only interested in the structure of G/Z it is always possible to replace G by G/X where X is the kernel of ζ and apply Theorem 1 to G/X . In Theorem 1 it is not necessarily the case that $G/K \cong Z(H)/Z$ as the following example shows. Let H be the elementary abelian group of order 8 generated by x, y, z . Let J be the cyclic group of order 4 generated by w and let J act as a group of automorphisms of H by $w(x) = y, w(y) = xz, w(z) = z$. Let $G = [H]J$ be the semidirect product. Then the center of G is generated by z and has order 2. Let ϕ be the faithful linear character on Z and extend ϕ to ψ on H by $\psi(y) = \psi(x) = 1$. Then ψ has 4 conjugates under J so $\psi^G = \zeta$ has degree = 4 and is irreducible. Thus G is a group of central type, $H \triangleleft G$, and both H and $G/H = J$ are abelian. Thus H is of central type with center H , the group K of Theorem 1 is H and $G/H \cong J \not\cong H/Z$. Notice, however, that in the group G one can find an abelian normal subgroup N such that $G/N \cong N/Z$, namely $N = \langle xy, w^2z \rangle$. Then $G/N \cong N/Z$ is the Klein 4-group.

Using the results in [7] it is easy to characterize the metacyclic groups \bar{G} admitting an irreducible projective representation of degree

$n = [\bar{G}:1]^{1/2}$. By 1.6 and 4.1 of [7] such a group must be the semi-direct product of two cyclic groups of order n . If $\bar{G} = \langle a, b \mid a^n = b^n = e, ba = a^r b \rangle$ is the semidirect product of a cyclic group of order n with itself then by 4.4 of [7] \bar{G} admits an irreducible projective representation of degree n if and only if $n - 1$ is the least positive integer exponent so that $1 + r + \dots + r^{n-1} = 0 \pmod n$. This was pointed out to me by the referee.

We conclude by showing how to construct a group G of central type with G/Z metabelian. This construction will characterize metabelian groups \bar{G} with an irreducible projective representation of degree $[\bar{G}:1]^{1/2}$.

First it is necessary to recall some terminology and results from § 2 of [6]. Let G be any finite group with normal subgroup N so that G/N is abelian. Let θ be an irreducible character on N with $\theta^g = \theta$ for all $g \in G$. Then (G, N, θ) is called a character triple. Let $x, y \in G$. The character θ can be extended to a character ψ on the subgroup $\langle N, y \rangle$ generated by N and y . The character ψ^x is another extension of ψ to $\langle N, y \rangle$ so $\psi^x = \lambda\psi$ for some linear character λ on $\langle N, y \rangle$. Define a complex valued bilinear function on $G \times G$ by the equation $\langle\langle x, y \rangle\rangle = \lambda(xy x^{-1} y^{-1})$. It is shown in [6] that this inner product is well defined. The bilinear function is nondegenerate if $\{x \in G \mid \langle\langle x, y \rangle\rangle = 1 \text{ for all } y \in G\} = N$. By 2.7 of [6] the triple (G, N, θ) is fully ramified if and only if the bilinear function defined above is nondegenerate.

Begin with an abelian group Z_1 and a cyclic subgroup Z . Let ϕ be a linear character on Z_1 such that $\phi|_Z$ is faithful. Let H be an abelian group of symmetric type and let $\gamma: H \times H \rightarrow Z$ be a nondegenerate bilinear function. Such a function exists if and only if $\exp(H)$ divides $[Z:1]$. Let H be any central extension of Z by H with factor set γ . Then (H, Z_1, ϕ) is a fully ramified triple and thus is a group of central type with center $Z(H) = Z_1$. Also $\phi^H = \lambda(e)\lambda$ for an irreducible character λ on H with $\lambda(e)^2 = [H:Z(H)]$. Next let \bar{K} be an abelian group of symmetric type. Let $\alpha: \bar{K} \rightarrow \text{Aut}(H)$ so that $\phi^{\alpha(x)} = \phi$ for all $x \in \bar{K}$ and $Z \subseteq \{y \in Z_1 \mid \alpha(x)(y) = y \text{ for all } x \in \bar{K}\}$. Note that $\phi^{\alpha(x)} = \phi$ if and only if $\lambda^{\alpha(x)} = \lambda$ since λ vanishes off Z_1 and Z_1 is characteristic in H . Let K be one of the normal extensions of H by \bar{K} determined by α . Then (K, H, λ) is a character triple and (K, H, λ) is fully ramified if and only if the form $\langle\langle \cdot, \cdot \rangle\rangle$ defined above is nondegenerate. In this case $\lambda^K = \rho(e)\rho$ where ρ is irreducible on K and $\rho(e)^2 = [K:Z]$. We choose a K so that (K, H, λ) is fully ramified. Finally let C be an abelian group with $[C:1] = [Z_1:Z]$ and let $\beta: C \rightarrow \text{Aut}(K)$ such that $\phi^{\beta(c)} \neq \phi$ for all $c \neq e$ and $Z \subseteq \{y \in K \mid \beta(c)(y) = y \text{ for all } c \in C\}$. Let G be any normal extension of K by C determined by β with G/H abelian.

Then $\rho^g = \zeta$ is irreducible on G since ρ is distinct from all its conjugates under C and $\zeta(e)^2 = [G:C]^2\rho(e)^2 = [G:Z]$. Thus G is a group of central type satisfying the hypothesis of Theorem 1 and an examination of the proof of Theorem 1 shows every group satisfying the hypothesis of Theorem 1 is constructed in this way.

REFERENCES

1. C. W. Curtis and I. Reiner, *Representation Theory of Finite Groups and Associative Algebras*, Interscience, New York, 1962, 2nd ed., 1966.
2. F. R. DeMeyer, *Galois theory in separable algebras over commutative rings*, III. J. Math., **10** (1966), 287-295.
3. F. R. DeMeyer and G. J. Janusz, *Finite groups with an irreducible representation of large Degree*, Math. Zeit., **108** (1979), 145-153.
4. R. Frucht, *Über die Darstellung endlicher abelscher Gruppen durch Kollineationen*, J. Reine Angew. Math., **166** (1931), 16-29.
5. S. M. Gagola, *Characters fully ramified over a normal subgroup*, Pacific J. Math., **55** (1974), 107-126.
6. I. M. Isaacs, *Characters of solvable and symplectic groups*, Amer. J. Math., **95** (1973), 594-635.
7. H. N. Ng, *Faithful irreducible projective representations of metabelian groups*, J. Algebra, **38** (1976), 8-28.

Received June 9, 1977 and in revised form November 18, 1977.

COLORADO STATE UNIVERSITY
FORT COLLINS, CO 80523

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, CA 90024

CHARLES W. CURTIS

University of Oregon
Eugene, OR 97403

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).

8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1978 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Stephanie Brewster Brewer Taylor Alexander, <i>Local and global convexity in complete Riemannian manifolds</i>	283
Claudi Alsina i Català, <i>On countable products and algebraic convexifications of probabilistic metric spaces</i>	291
Joel David Berman and George Grätzer, <i>Uniform representations of congruence schemes</i>	301
Ajit Kaur Chilana and Kenneth Allen Ross, <i>Spectral synthesis in hypergroups</i>	313
David Mordecai Cohen and Howard Leonard Resnikoff, <i>Hermitian quadratic forms and Hermitian modular forms</i>	329
Frank Rimi DeMeyer, <i>Metabelian groups with an irreducible projective representation of large degree</i>	339
Robert Ellis, <i>The Furstenberg structure theorem</i>	345
Heinz W. Engl, <i>Random fixed point theorems for multivalued mappings</i>	351
William Andrew Ettl, <i>On arc length sharpenings</i>	361
Kent Ralph Fuller and Joel K. Haack, <i>Rings with quivers that are trees</i>	371
Kenneth R. Goodearl, <i>Centers of regular self-injective rings</i>	381
John Gregory, <i>Numerical algorithms for oscillation vectors of second order differential equations including the Euler-Lagrange equation for symmetric tridiagonal matrices</i>	397
Branko Grünbaum and Geoffrey Shephard, <i>Isotoxal tilings</i>	407
Myron Stanley Henry and Kenneth Leroy Wiggins, <i>Applications of approximation theory to differential equations with deviating arguments</i>	431
Mark Jungerman, <i>The non-orientable genus of the n-cube</i>	443
Robert Richard Kallman, <i>Only trivial Borel measures on S_∞ are quasi-invariant under automorphisms</i>	453
Joyce Longman and Michael Rich, <i>Scalar dependent algebras in the alternative sense</i>	463
Richard A. Mollin, <i>The Schur group of a field of characteristic zero</i>	471
David Pokrass, <i>Some radical properties of rings with $(a, b, c) = (c, a, b)$</i>	479
Margaret Shay and Paul Ruel Young, <i>Characterizing the orders changed by program translators</i>	485
Jerrold Norman Siegel, <i>On the structure of $B_\infty(F)$, F a stable space</i>	491
Surjeet Singh, <i>(hnp)-rings over which every module admits a basic submodule</i>	509
A. K. Snyder, <i>Universal interpolating sets and the Nevanlinna-Pick property in Banach spaces of functions</i>	513
Jeffrey D. Vaaler, <i>On the metric theory of Diophantine approximation</i>	527
Roger P. Ware, <i>When are Witt rings group rings? II</i>	541