

# Pacific Journal of Mathematics

**SUBMANIFOLDS WITH  $L$ -FLAT NORMAL CONNECTION OF  
THE COMPLEX PROJECTIVE SPACE**

MASAFUMI OKUMURA

# SUBMANIFOLDS WITH $L$ -FLAT NORMAL CONNECTION OF THE COMPLEX PROJECTIVE SPACE

MASAFUMI OKUMURA

**Real submanifolds with  $L$ -flat normal connection of the complex projective space are studied. As a special case “a complex submanifold with  $L$ -flat normal connection of the complex projective space is necessarily totally geodesic” is proved.**

**Introduction.** As is well known an odd-dimensional sphere  $S^{2n+1}$  is a principal circle bundle over a complex projective space  $P^n(C)$ . The Riemannian structure on  $P^n(C)$  is given by the submersion  $\pi: S^{2n+1} \rightarrow P^n(C)$  which is defined by the Hopf-fibration. If we construct a circle bundle over a real submanifold of  $P^n(C)$  in such a way that it is compatible with the Hopf-fibration, the circle bundle is a submanifold of the odd-dimensional sphere. Thus when we want to study submanifolds of the complex projective space it is useful to study the circle bundle over the submanifold. From this point of view, H. B. Lawson, Jr. [2] and the present author [3, 4, 5] have studied real submanifolds of the complex projective space. In the previous paper [5], the author studied relations between the normal connection of a submanifold of  $P^n(C)$  and that of the circle bundle over the submanifold and established the notion of  $L$ -flatness for the normal connection of a real submanifold of  $P^n(C)$ .

The purpose of the present paper is to study submanifolds with  $L$ -flat normal connection of  $P^n(C)$ . The main result is the following.

**THEOREM 1.** *The totally geodesic complex projective linear subspaces  $P^n(C)$  are the only complex submanifolds with  $L$ -flat normal connection of  $P^{n+p}(C)$ .*

In §1 we state some formulas for real submanifolds of a Kaehlerian manifold and in §2 we discuss the case when the ambient manifold is the complex projective space. There we explain  $L$ -flatness of the normal connection. In §3, we calculate the Laplacian for a function which is defined on the submanifold and prove some theorems including Theorem 1.

1. **Real submanifolds of a Kaehlerian manifold.** Let  $M'$  be a real  $(n + p)$ -dimensional Kaehlerian manifold with Kaehlerian

structure  $(J, G)$ , that is,  $J$  is the endomorphism of the tangent bundle  $T(M')$  satisfying  $J^2 = -$  identity and  $G$  the Riemannian metric of  $M'$  satisfying the Hermitian condition  $G(JX', JY') = G(X', Y')$  for any  $X', Y' \in T(M')$ .

Let  $M$  be an immersed submanifold of  $M'$  and  $i$  be the immersion. Then the tangent bundle  $T(M)$  is identified with a subbundle of  $T(M')$  and a Riemannian metric  $g$  of  $M$  is induced from the Riemannian metric  $G$  of  $M'$  in such a way that  $g(X, Y) = G(iX, iY)$ , where  $X, Y \in T(M)$ . The normal bundle  $N(M)$  is the subbundle of  $T(M')$  consisting of all  $X' \in T(M')$  which are orthogonal to  $T(M)$  with respect to  $G$ . At each point of  $M$ , we choose orthonormal normal vectors  $n_1, n_2, \dots, n_p$  to  $M$  and extend them respectively to  $N_1, N_2, \dots, N_p$  in such a way that they belong to  $N(M)$ .

For any  $X \in T(M)$  and for  $N_A, A = 1, 2, \dots, p$ , the transforms  $JiX$  and  $JN_A$  are respectively written in the following forms:

$$(1.1) \quad JiX = iFX + \sum_{A=1}^p u^A(X)N_A,$$

$$(1.2) \quad JN_A = -iU_A + \sum_{B=1}^p F'^B_A N_B,$$

where  $F, F', U_A$ , and  $u^A$  define respectively an endomorphism of  $T(M)$ , that of  $N(M)$ , local tangent vector fields and local 1-forms on  $M$ . They satisfy the relations  $u^A(X) = g(U_A, X)$ ,  $F'^B_A = -F'^A_B$ . Applying  $J$  to both side members of (1.1) and (1.2), we find

$$(1.3) \quad F^2X = -X + \sum_{A=1}^p u^A(X)U_A,$$

$$(1.4) \quad u^A(FX) = -\sum_{B=1}^p F'^A_B u^B(X),$$

$$(1.5) \quad FU_A = -\sum_{B=1}^p F'^B_A U_B,$$

$$(1.6) \quad \sum_{\sigma=1}^p F'^\sigma_A F'^B_\sigma = -\delta^B_A + u^B(U_A).$$

If at any point of  $M$ ,  $FU_A = 0$  are valid for  $A = 1, 2, \dots, p$ , we know that  $F$  satisfies  $F^3 + F = 0$  and that the rank of  $F \geq n - p$ . In this case the submanifold  $M$  is called an  $F$ -submanifold of rank  $\geq n - p$  [7]. We denote by  $\nabla$  and  $D$  the Riemannian connection of  $M$  and  $M'$  respectively and by  $D^N$  the induced normal connection from  $D$  to  $N(M)$ . Then they are related by the following Gauss and Weingarten equations.

$$(1.7) \quad D_{i_X} i Y = i \nabla_X Y + h(X, Y), \quad h(X, Y) = \sum_{A=1}^p h^A(X, Y) N_A,$$

$$(1.8) \quad D_{i_X} N_A = -i H_A X + D_X^N N_A, \quad D_X^N N_A = \sum_{B=1}^p L_A^B(X) N_B,$$

where  $h(X, Y)$  is the second fundamental form and  $H_A$ 's are symmetric linear transformations of  $T(M)$  which are called the Weingarten maps for the normal  $N_A$ . The last two equations show that  $h^A(X, Y) = g(H_A X, Y)$ . The mean curvature vector field  $\mu$  of  $M$  is defined by

$$(1.9) \quad \mu = \left( \sum_{A=1}^p (\text{trace } H_A) N_A \right) / n$$

and it is well known that  $\mu$  is independent of the choice of  $N_A$ 's. If the mean curvature vector field vanishes identically on  $M$ ,  $M$  is called a minimal submanifold of  $M'$ . By definition  $M$  is minimal if and only if  $\text{trace } H_A = 0$ ,  $A = 1, 2, \dots, p$  at each point of  $M$ . Differentiating (1.1) and (1.2) covariantly and making use of the fact that the Riemannian connection  $D$  of  $M'$  leaves the almost complex structure  $J$  invariant, we have

$$(1.10) \quad (\nabla_Y F) X = \sum_{A=1}^p \{u^A(X) H_A Y - g(H_A X, Y) U_A\},$$

$$(1.11) \quad \nabla_X U_A = F H_A X + \sum_{B=1}^p \{L_A^B(X) U_B - F'^B H_B X\},$$

$$(1.12) \quad D_X^N F'^B = g(H_B U_A - H_A U_B, X).$$

Differentiating (1.9) covariantly, we have

$$n D_X^N \mu = \sum_{A=1}^p \left\{ X (\text{trace } H_A) N_A + \sum_{B=1}^p (\text{trace } H_A) L_A^B(X) N_B \right\},$$

from which we know that the mean curvature vector field is parallel with respect to the normal connection if and only if

$$(1.13) \quad X (\text{trace } H_A) = \sum_{B=1}^p L_A^B(X) \text{trace } H_B,$$

because of the fact that  $L_A^B(X) = -L_A^B(X)$  for any  $X \in T(M)$ .

**2. Real submanifolds of the complex projective space.** Let the ambient manifold  $M'$  be the complex projective space  $P^{(n+p)/2}(C)$  with the Fubini-Study metric of constant holomorphic sectional curvature 4. Since the curvature tensor  $R'(X', Y')Z'$  of the ambient manifold satisfies

$$(2.1) \quad R'(X', Y')Z' = G(Y', Z')X' - G(X', Z')Y' \\ + G(JY', Z')JX' - G(JX', Z')JY' - 2G(JX', Y')JZ'$$

the Codazzi, Ricci equations become respectively

$$(2.2) \quad (\nabla_X H_A)Y - (\nabla_Y H_A)X = \sum_{B=1}^p \{L_A^B(X)H_B Y - L_A^B(Y)H_B X\} \\ - u^A(Y)FX + u^A(X)FY - 2g(FX, Y)U_A,$$

$$(2.3) \quad R^N(X, Y)N_A = \sum_{B=1}^p \{g((H_A H_B - H_B H_A)X, Y) \\ + u^A(Y)u^B(X) - u^A(X)u^B(Y) - 2g(FX, Y)F_A'^B\}N_B,$$

where  $R^N(X, Y)$  denotes the normal curvature of  $M'$ . If we choose an orthonormal basis  $\{E_1, \dots, E_n\}$  of  $T_x(M)$  at a point  $x \in M$ , it follows from (1.4) and (2.3) that

$$(2.4) \quad \sum_{i=1}^n R^N(FE_i, E_i)N_A = \sum_{B=1}^p \{\text{trace}(H_A H_B - H_B H_A)F + 2g(FU_A, U_B) \\ - 2(\text{trace } F^2)F_A'^B\}N_B.$$

Now recall the fact that an  $n + p + 1$ -dimensional sphere of radius 1 is a principal circle bundle over the complex projective space and let  $\pi$  be the Hopf-fibration. We construct the circle bundle over the submanifold  $M$  in such a way that the following diagram commutes:

$$\begin{array}{ccc} \pi^{-1}(M) & \longrightarrow & S^{n+p+1} \\ \downarrow & & \downarrow \\ M & \longrightarrow & P^{(n+p)/2}(C). \end{array}$$

In the previous paper [5] the author proved that the normal connection of  $\pi^{-1}(M)$  in  $S^{n+p+1}$  is flat if and only if the following two conditions are satisfied on  $M$ :

$$(2.5) \quad R^N(X, Y)N_A = -2g(FX, Y) \sum_{B=1}^p F_A'^B N_B,$$

$$(2.6) \quad D_X^N F_A'^B = 0.$$

Moreover the author proved that if  $n > p + 2$ , (2.5) implies (2.6). Thus we call the normal connection of  $M$  in  $P^{(n+p)/2}(C)$  *lift-flat* normal connection or briefly *L-flat* normal connection if it satisfies (2.5) and (2.6). It is easily checked that the totally geodesic complex submanifold  $P^{n/2}(C)$  of  $P^{(n+p)/2}(C)$  is a submanifold with *L-flat* normal connection. We can also check the fact that if the submanifold  $M$  is a complex submanifold, the normal connection is *L-flat* if

and only if the Weingarten maps  $H_A$  and  $H_B$  commute for any pair of  $A$  and  $B = 1, 2, \dots, p$ .

**3. Submanifolds with  $L$ -flat normal connection.** We put

$$(3.1) \quad f = \sum_{A=1}^p u^A(U_A).$$

Then from (1.3) we have

$$(3.2) \quad f = \text{trace } F^2 + n,$$

which shows that  $f$  is a globally defined function on the submanifold  $M$ . Using the formulas which are stated in §1, we now calculate the Laplacian  $\Delta f$ .

By means of (1.10) and (3.2), it follows that

$$(3.3) \quad \frac{1}{2} Yf = \frac{1}{2} Y(\text{trace } F^2) = \text{trace } (\nabla_Y F)F = 2 \sum_{A=1}^p g(FH_A Y, U_A),$$

from which

$$(3.4) \quad \begin{aligned} & \frac{1}{4} (\nabla_X \nabla_Y f - \nabla_{\nabla_X Y} f) = \frac{1}{4} (\nabla_X (Yf) - (\nabla_X Y)f) \\ & = \sum_{A=1}^n \{g((\nabla_X F)H_A Y, U_A) + g(F(\nabla_X H_A)Y, U_A) \\ & \quad + g(FH_A Y, \nabla_X U_A)\} \\ & = \sum_{A,B=1}^p \{g(H_A U_B, Y)g(H_B U_A, X) - g(H_A H_B X, Y)g(U_A, U_B) \\ & \quad - g((\nabla_X H_A)F U_A, Y) - g(F^2 H_A Y, H_A X) \\ & \quad - L_A^B(X)g(H_A F U_B, Y) - F_A^B g(X, H_B F H_A Y)\}, \end{aligned}$$

because of (1.10) and (1.11).

On the other hand, from (2.2), it follows that

$$\begin{aligned} (\nabla_X H_A)F U_A &= (\nabla_{F U_A} H_A)X + \sum_{B=1}^p \{L_A^B(X)H_B F U_A - L_A^B(F U_A)H_B X\} \\ & \quad + u^A(X)F^2 U_A - 2g(FX, F U_A)U_A. \end{aligned}$$

Substituting this into (3.4), we find

$$\begin{aligned} & \frac{1}{4} (\nabla_X \nabla_Y f - \nabla_{\nabla_X Y} f) = \sum_{A,B=1}^p \{g(H_A U_B, Y)g(H_B U_A, X) \\ & \quad - g(H_A H_B X, Y)g(U_A, U_B) - g((\nabla_{F U_A} H_A)X, Y) \\ & \quad + L_A^B(F U_A)g(H_B X, Y) - g(U_A, X)g(F^2 U_A, Y) \\ & \quad - 2g(F^2 U_A, X)g(U_A, Y) + g(H_A^2 Y, X) \\ & \quad - g(H_A U_B, Y)g(H_A U_B, X) - F_A^B g(X, H_B F H_A Y)\}, \end{aligned}$$

from which

$$(3.5) \quad \begin{aligned} \frac{1}{4} \Delta f &= \sum_{A, B=1}^p \{g(H_A U_B, H_B U_A - H_A U_B) - \text{trace}(H_A H_B)g(U_A, U_B) \\ &\quad - (\nabla_{FU_A}(\text{trace } H_A) - L_A^B(FU_A) \text{trace } H_B) \\ &\quad + 3g(FU_A, FU_A) + \text{trace } H_A^2 - F_A'^B \text{trace } H_A H_B F\}. \end{aligned}$$

Now we rewrite (3.5), using the normal curvature  $R^N$ . By means of (2.4) and the fact that  $F$  is a skew-symmetric linear transformation, we have

$$(3.6) \quad \begin{aligned} \frac{1}{4} \Delta f &= \sum_{A, B=1}^p \left\{ g(H_A U_B, H_B U_A - H_A U_B) - \text{trace}(H_A H_B)g(U_A, U_B) \right. \\ &\quad - (\nabla_{FU_A}(\text{trace } H_A) - L_A^B(FU_A) \text{trace } H_B) \\ &\quad + 3g(FU_A, FU_A) + \text{trace } H_A^2 \\ &\quad - F_A'^B \left( \frac{1}{2} \sum_{i=1}^n G(R^N(FE_i, E_i)N_A, N_B) - g(FU_A, U_B) \right. \\ &\quad \left. \left. + (\text{trace } F^2)F_A'^B \right) \right\}. \end{aligned}$$

Before we prove Theorem 1, we prove the following more general result.

**THEOREM 2.** *Let  $M$  be a real submanifold with  $L$ -flat normal connection of a complex projective space. If the mean curvature vector field is parallel with respect to the normal connection, then the submanifold  $M$  is an  $F$ -submanifold of rank  $\geq n - p$ . Particularly, if  $F$  is of almost everywhere rank  $n$ ,  $M$  is a totally geodesic complex submanifold and consequently a complex projective linear subspace.*

We begin with the following

**LEMMA.** *When the normal connection of  $M$  in  $P^{(n+p)/2}(C)$  is  $L$ -flat, the function  $f$  is constant.*

*Proof.* It follows from (1.5) and (3.3) that

$$\begin{aligned} \frac{1}{4} Yf &= - \sum_{A=1}^p g(H_A Y, FU_A) = \sum_{A, B=1}^p F_A'^B g(H_A U_B, Y) \\ &= \frac{1}{2} \sum_{A, B=1}^p F_A'^B g(H_A U_B - H_B U_A, Y). \end{aligned}$$

Thus, (1.12) and (2.6) imply that  $f$  is constant.

*Proof of Theorem 3.* Since the  $p \times p$  matrix (trace  $H_A H_B$ ) is symmetric and can be assumed to be diagonal for a suitable choice of  $N_1, N_2, \dots, N_p$ ,

$$\sum_{A,B=1}^p \text{trace}(H_A H_B)g(U_A, U_B) = \sum_{A=1}^p (\text{trace } H_A^2)g(U_A, U_A).$$

Moreover the conditions of the theorem, (1.13), (2.5), (2.6), and (3.6) imply that

$$\begin{aligned} \nabla_{FU_A}(\text{trace } H_A) &= \sum_{B=1}^p L_A^B(FU_A) \text{trace } H_B, \\ \sum_{i=1}^n G(R^N(FE_i, E_i)N_A, N_B) &= -2(\text{trace } F^2)F_A'^B, \end{aligned}$$

and

$$\sum_{B=1}^p F_A'^B g(FU_A, U_B) = -g(FU_A, FU_A).$$

Thus we have

$$(3.7) \quad \frac{1}{4}\Delta f = \sum_{A=1}^p \{\text{trace } H_A^2 - (\text{trace } H_A^2)g(U_A, U_A) + 2g(FU_A, FU_A)\} = 0,$$

from which  $FU_A = 0$ ,  $A = 1, 2, \dots, p$ , because of  $g(U_A, U_A) \leq 1$ . Thus  $M$  is an  $F$ -submanifold of rank  $\geq n - p$ . To prove the last part of the theorem we assume that the rank of  $F$  is almost everywhere  $n$ . Then we have  $U_A = 0$  for  $A = 1, 2, \dots, p$ . This means that the submanifold is a complex submanifold. Again we use (3.7) and get that  $M$  is totally geodesic. This completes the proof.

*Proof of Theorem 1.* It is well known [6] that a complex submanifold is a minimal submanifold and the induced complex structure  $F$  is of rank  $n$ . Hence a complex submanifold with  $L$ -flat normal connection satisfies the conditions of Theorem 3. This completes the proof of Theorem 1.

Finally we point out that, as an application of our discussions the formula (3.6) gives another proof of the following known [1].

**THEOREM 3.** *There does not exist a complex submanifold with flat normal connection of a complex projective space.*

*Proof.* Since a complex submanifold is invariant under  $J$  it follows that  $U_A = 0, u^A = 0$  for  $A = 1, 2, \dots, p$ . Hence (3.6) becomes

$$\begin{aligned} \frac{1}{4}\Delta f &= \sum_{A=1}^p \text{trace } H_A^2 - \sum_{A,B=1}^p (\text{trace } F^2)F_A'^B F_A'^B \\ &= \sum_{A=1}^p \text{trace } H_A^2 + np > 0, \end{aligned}$$



because of (1.6). On the other hand, (3.3) implies that the function  $f$  is constant and consequently  $\Delta f = 0$ . This is contradiction. This completes the proof.

#### REFERENCES

1. B. Y. Chen and K. Ogiue, *Some extrinsic results for Kaehler submanifolds*, Tamkang J. Math., **4** (1973), 207-213.
2. H. B. Lawson Jr., *Rigidity theorems in rank-1 symmetric spaces*, J. of Differential Geometry, **4** (1970), 349-357.
3. M. Okumura, *On some real hypersurfaces of a complex projective spaces*, Trans. Amer. Math. Soc., **212** (1975), 355-364.
4. M. Okumura, *Submanifolds of real codimension of a complex projective space*, Atti della Accademia Nazionale dei Lincei, **58** (1975), 544-555.
5. ———, *Normal curvature and real submanifold of the complex projective space*, Geometriae Dedicata.
6. J. A. Schouten and K. Yano, *On invariant subspaces in the almost complex  $X_{2n}$* , Indagatione Math., **17** (1955), 261-269.
7. K. Yano and S. Ishihara, *The  $f$ -structure induced on submanifolds of complex and almost complex spaces*, Kōdai Math. Sem. Rep., **18** (1966), 120-160.

Received August 9, 1977 and in revised form March 26, 1978.

SAITAMA UNIVERSITY  
URAWA, JAPAN

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)

University of California  
Los Angeles, CA 90024

CHARLES W. CURTIS

University of Oregon  
Eugene, OR 97403

C. C. MOORE

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University  
Stanford, CA 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).

8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1978 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

Su-Shing Chen, <i>Weak rigidity of compact negatively curved manifolds</i> . . . . .	273
Heinz Otto Cordes and D. A. Williams, <i>An algebra of pseudodifferential operators with nonsmooth symbol</i> . . . . .	279
Herbert Paul Halpern, <i>Normal expectations and integral decomposition of type III von Neumann algebras</i> . . . . .	291
G. Hochschild, <i>On representing analytic groups with their automorphisms</i> . . . . .	333
Dean G. Hoffman and David Anthony Klarner, <i>Sets of integers closed under affine operators—the closure of finite sets</i> . . . . .	337
Simeon Ivanov, <i>On holomorphic relative inverses of operator-valued functions</i> . . . . .	345
O. P. Juneja and M. L. Mogra, <i>Radii of convexity for certain classes of univalent analytic functions</i> . . . . .	359
Hadi Kharaghani, <i>The evolution of bounded linear functionals with application to invariant means</i> . . . . .	369
Jack W. Macki, <i>A singular nonlinear boundary value problem</i> . . . . .	375
A. W. Mason and Walter Wilson Stothers, <i>Remarks on a theorem of L. Greenberg on the modular group</i> . . . . .	385
Kevin Mor McCrimmon, <i>Peirce ideals in Jordan algebras</i> . . . . .	397
John C. Morgan, II, <i>On the absolute Baire property</i> . . . . .	415
Gerard J. Murphy, <i>Commutative non-Archimedean <math>C^*</math>-algebras</i> . . . . .	433
Masafumi Okumura, <i>Submanifolds with <math>L</math>-flat normal connection of the complex projective space</i> . . . . .	447
Chull Park and David Lee Skoug, <i>Distribution estimates of barrier-crossing probabilities of the Yeh-Wiener process</i> . . . . .	455
Irving Reiner, <i>Invariants of integral representations</i> . . . . .	467
Phillip Schultz, <i>The typeset and cotypeset of a rank 2 abelian group</i> . . . . .	503
John Brendan Sullivan, <i>Representations of Witt groups</i> . . . . .	519
Chia-Chi Tung, <i>Equidistribution theory in higher dimensions</i> . . . . .	525
Toshio Uda, <i>Complex bases of certain semiproper holomorphic maps</i> . . . . .	549