

# Pacific Journal of Mathematics

**COMPLEX BASES OF CERTAIN SEMIPROPER  
HOLOMORPHIC MAPS**

TOSHIO UDA

## COMPLEX BASES OF CERTAIN SEMI-PROPER HOLOMORPHIC MAPS

TOSHIO UDA

**The existence theorem of complex bases of quasi-proper holomorphic maps was studied by N. Kuhlmann. In this paper the existence of the complex bases in a more general case will be shown.**

**O. Introduction.** In the function theory of several complex variables, the complex bases of holomorphic maps of analytic spaces have been introduced as a generalized concept of Riemann surfaces defined by inverse functions of given holomorphic functions of one complex variable.

Let  $f: X \rightarrow Y$  be a holomorphic map of analytic spaces. How does  $f$  have a complex base? Authors have discussed the sufficient conditions which allow for the existence of a complex base of  $f$  (cf. for example, [3], [5], [6], [7]). If  $f$  is proper, then  $f$  has a complex base ([7]). N. Kuhlmann [3] showed existence theorems in the case of quasi-proper ( $N$ -quasi-proper).  $f$  is called *quasi-proper* (resp.  *$N$ -quasi-proper*) if, for every compact subset  $K$  of  $Y$ , there exists a compact subset  $\tilde{K}$  of  $X$  such that each of the irreducible branches (resp. each of the connected components) of fibres on  $K$  intersects  $\tilde{K}$ .

On this subject, an attempt will be made to abate the condition, so that each of the given unions of connected components of fibres intersects  $\tilde{K}$ . For such holomorphic maps, we shall have an existence theorem of complex bases (of type of N. Kuhlmann's).

**THEOREM.** *Let  $X$  be an irreducible normal analytic space,  $f: X \rightarrow Y$  be a holomorphic map of  $X$  into an analytic space  $Y$  and  $E_f$  be the set of degeneracy of  $f$ . Suppose that  $f$  satisfies (C) and that  $f(E_f)$  is analytic in  $Y$ . Then  $f$  has a complex base  $(\tilde{Z}, \tilde{\varphi})$  and  $\tilde{Z}$  is also normal. Moreover, the natural holomorphic map  $\tilde{\varphi}$  with  $f = \tilde{\varphi} \circ \tilde{\varphi}$  is proper and light, and  $\tilde{\varphi}$  satisfies  $(C_1)$ .*

**1. Preliminaries.** We assume in this paper that all analytic spaces are reduced and have countable bases of open sets.

Let  $f: X \rightarrow Y$  and  $f_1: X \rightarrow Y_1$  be holomorphic maps of analytic spaces.  $f_1$  is said to *strictly depend* on  $f$ , if  $f_1$  is constant on each connected component of fibres of  $f$ .  $f_1$  is said to be *analytically related* to  $f$ , if  $f$  and  $f_1$  strictly depend on each other. A pair  $(Z, \varphi)$  is called a *complex base* of  $f$ , if  $Z$  is an analytic space, and

if  $\varphi: X \rightarrow Z$  is a surjective holomorphic map which is analytically related to  $f$ , and if, for each holomorphic map  $h: X \rightarrow T$  which strictly depends on  $f$ , there exists a unique holomorphic map  $\psi: Z \rightarrow T$  with  $h = \psi \circ \varphi$ .

A holomorphic map  $f: X \rightarrow Y$  is said to be *semi-proper*, if, for each compact subset  $K$  of  $Y$ , there exists a compact subset  $\tilde{K}$  of  $X$  such that  $f^{-1}(y) \cap \tilde{K} \neq \emptyset$ , for  $y \in K \cap f(X)$ ;  $f$  is said to be *quasi-proper* if  $B \cap \tilde{K} \neq \emptyset$ , for each irreducible branch  $B$  of  $f^{-1}(y)$ . N. Kuhlmann modified this definition ([3]);  $f$  is said to be *N-quasi-proper*, if  $N \cap \tilde{K} \neq \emptyset$ , for each connected component  $N$  of  $f^{-1}(y)$ . He showed the existence of complex bases of *N-quasi-proper* holomorphic maps in [3].

Now, we consider a more general case in which each of the given unions of connected components of  $f^{-1}(y)$  ( $y \in K \cap f(X)$ ) intersects  $\tilde{K}$ .

DEFINITION. A holomorphic map  $f: X \rightarrow Y$  is said to *satisfy (C)* if  $f$  has the following property;

(C) Given an analytic set  $A$  in  $Y$  and a commutative diagram of holomorphic maps

$$\begin{array}{ccc} X - f^{-1}(A) & \xrightarrow{\varphi} & Z \\ & \searrow (f, h) & \downarrow \psi \\ & & (Y - A) \times T \end{array}$$

where  $\psi$  is light (that is, each fibre is discrete) and  $h: X \rightarrow T$  strictly depends on  $f$  and  $(f, h)$  is a holomorphic map given by  $x \mapsto (f(x), h(x))$ , and if  $K$  is a compact subset of  $(Y - A) \times T$ , then there exists a compact subset  $\tilde{K}$  of  $X - f^{-1}(A)$  such that  $\varphi^{-1}(p) \cap \tilde{K} \neq \emptyset$ , for  $p \in \psi^{-1}(K) \cap \varphi(X - f^{-1}(A))$ .

If  $f$  satisfies (C), then  $f$  satisfies the following (C<sub>1</sub>) (we take  $A = \emptyset$  and  $h = f$ );

(C<sub>1</sub>) Given a compact subset  $K$  of  $Y$  and a commutative diagram of holomorphic maps

$$\begin{array}{ccc} X & \xrightarrow{\varphi} & Z \\ & \searrow f & \downarrow \psi \\ & & Y \end{array}$$

where  $\psi$  is light, then there exists a compact subset  $\tilde{K}$  of  $X$  such that  $\varphi^{-1}(p) \cap \tilde{K} \neq \emptyset$  for  $p \in \psi^{-1}(K) \cap \varphi(X)$ .

Note that in such cases,  $f$ ,  $\varphi$ , and  $(f, h)$  (in the two diagrams above) are naturally semi-proper, and that if  $\varphi$  is surjective,  $\psi$  is

proper; in this case  $\psi$  is finite! Every  $N$ -quasi-proper holomorphic map satisfies (C); for, if  $f: X \rightarrow Y$  is  $N$ -quasi-proper, so is  $(f, h)$ . Thus we have the following inclusion:

$$\begin{aligned} \text{proper} &\implies \text{quasi-proper} \implies N\text{-quasi-proper} \\ &\implies (C) \implies (C_1) \implies \text{semi-proper} . \end{aligned}$$

LEMMA 1.1. ([4]) *Let  $Y$  be a normal analytic space and  $f: X \rightarrow Y$  be a proper modification map. If  $f$  is nowhere degenerate, then  $f$  is a biholomorphic map.*

LEMMA 1.2. *Let  $X$  be an irreducible normal analytic space and  $f: X \rightarrow Y$  be a nowhere degenerate holomorphic map. If  $f$  satisfies  $(C_1)$ , then  $f$  has a complex base  $(Z, \varphi)$ , and the natural holomorphic map  $\psi: Z \rightarrow Y$  with  $f = \psi \circ \varphi$  is proper and light.*

*Proof.*  $f$  is nowhere degenerate, and so  $f$  has a complex base  $(Z, \varphi)$  (cf. [6]).  $f$  is semi-proper, so  $f(X)$  is analytic in  $Y$  (cf. [1]). Thus we may assume that  $f$  is surjective. Since  $f$  and  $\varphi$  are analytically related and  $X$  has a countable basis,  $\psi^{-1}(y) = \varphi(f^{-1}(y))$  is discrete for  $y \in Y$ . Thus  $\psi$  is light and therefore, is proper, as desired.

2. Proof of theorem. We shall prove our theorem by introducing modification of the proof of the theorem of N. Kuhlmann.

We may assume that  $f$  is surjective as in Lemma 1.2, and moreover, that  $Y$  is a connected complex manifold since the set of singular points of  $Y$  is a thin analytic set in  $Y$ . By [1], Proposition 1.24,  $f(E_f)$  is thin of dimension  $\leq \dim Y - 2$ .

Let  $Y' = Y - f(E_f)$ ,  $X' = X - f^{-1}(f(E_f))$  and  $f' = f|X' \rightarrow Y'$ . Since  $f'$  satisfies  $(C_1)$  and is nowhere degenerate,  $f'$  has a complex base  $(Z', \varphi')$  and the natural holomorphic map  $\psi'$  with  $f' = \psi' \circ \varphi'$  is proper and light by Lemma 1.2.  $Z'$  is a normal analytic space. By [5], Satz 1 (or [2], Satz A), we have a (unique up to biholomorphic equivalence) normal analytic space  $\tilde{Z}$  with a holomorphic map  $\tilde{\psi}: \tilde{Z} \rightarrow Y$  which is proper, light and surjective such that  $Z' = \tilde{\psi}^{-1}(Y')$  and  $Z'$  is dense in  $\tilde{Z}$  and  $\psi' = \tilde{\psi}|Z'$ .

We have to show that there exists a surjective holomorphic map  $\tilde{\varphi}: X \rightarrow \tilde{Z}$  such that  $(\tilde{Z}, \tilde{\varphi})$  is a complex base of  $f$ .

(a) A holomorphic map  $\varphi': X' \rightarrow Z'$  can uniquely be extended to the surjective holomorphic map  $\tilde{\varphi}: X \rightarrow \tilde{Z}$  such that  $\tilde{\varphi}$  is analytically related to  $f$  and  $f = \tilde{\psi} \circ \tilde{\varphi}$ : Let  $G \subset X \times Y$  be a graph of  $f: X \rightarrow Y$  and  $G' \subset X' \times Z'$  be a graph of  $\varphi': X' \rightarrow Z'$ . Let  $\iota \times \tilde{\psi}: X \times \tilde{Z} \rightarrow X \times Y$  be a holomorphic map given by  $(x, p) \mapsto (x, \tilde{\psi}(p))$  and

$G_1 := (\iota \times \tilde{\psi})^{-1}(G)$ . There exists an irreducible branch  $\tilde{G}$  of  $G_1$  with  $\tilde{G} \cap (X' \times Z') = G'$ .

The projection  $\pi_1: \tilde{G} \rightarrow X$  onto 1st component  $X$  is a proper light modification map and therefore,  $\pi_1$  is biholomorphic by Lemma 1.1. Let  $\pi_2: \tilde{G} \rightarrow \tilde{Z}$  be the projection onto 2nd component  $\tilde{Z}$  and  $\tilde{\varphi} := \pi_2 \circ \pi_1^{-1}: X \rightarrow \tilde{Z}$ . Then  $f = \tilde{\psi} \circ \tilde{\varphi}$ . And since  $\tilde{\psi}$  is light,  $\tilde{\varphi}$  is analytically related to  $f$ . Uniqueness of  $\tilde{\varphi}$  is obvious.

(b) Let  $h: X \rightarrow T$  be a holomorphic map strictly depending on  $f$ . Then there exists a (unique) holomorphic map  $\psi: \tilde{Z} \rightarrow T$  such that  $h = \psi \circ \tilde{\varphi}$ : Since  $f$  satisfies (C),  $(\tilde{\varphi}, h): X \rightarrow \tilde{Z} \times T$  is semi-proper and therefore,  $G_0 := (\tilde{\varphi}, h)(X)$  is analytic in  $\tilde{Z} \times T$ . Let  $\tilde{\pi}_1$  and  $\tilde{\pi}_2$  be projections of  $G_0$  onto 1st and 2nd components, respectively.  $\pi_1|_{G_0} \cap (Z' \times T) \rightarrow Z'$  is a biholomorphic map. In fact,  $h'|_{X'}$  strictly depends on  $f'$  and  $(Z', \varphi')$  is a complex base of  $f'$ , so there exists a holomorphic map  $\psi': Z' \rightarrow T$  with  $h' = \psi' \circ \varphi'$ . And then,  $G_0 \cap (Z' \times T)$  is a graph of  $\psi'$ .  $h$  strictly depends on  $\tilde{\varphi}$ , so  $\tilde{\pi}_1$  is light.  $f$  satisfies (C<sub>1</sub>) and  $\tilde{\varphi} = \tilde{\pi}_1 \circ (\tilde{\varphi}, h)$ , so  $\tilde{\pi}_1$  is proper. Thus  $\tilde{\pi}_1: G_0 \rightarrow \tilde{Z}$  is a proper light modification map and therefore, it is biholomorphic. Let  $\psi := \tilde{\pi}_2 \circ \tilde{\pi}_1^{-1}: \tilde{Z} \rightarrow T$ , then  $h = \psi \circ \tilde{\varphi}$ .

With (a) and (b) we conclude the proof.

## REFERENCES

1. A. Andreotti and W. Stoll, *Analytic and algebraic dependence of meromorphic functions*, Lecture Notes in Math. 234, Springer (1971).
2. N. Kuhlmann, *Niveaumengenräume holomorpher Abbildungen und nullte Bildgarben*, Manuscripta Math., **1** (1969), 147-189.
3. ———, *Komplexe Basen zu quasieigentlichen holomorphen Abbildungen*, Comm. Math. Helv., **48** (1973), 340-353.
4. R. Remmert, *Holomorphe und meromorphe Abbildungen komplexer Räume*, Math. Ann., **133** (1957), 328-370.
5. K. Stein, *Analytische Zerlegungen komplexer Räume*, Math. Ann., **132** (1956), 68-93.
6. ———, *Die Existenz komplexer Basen zu holomorphen Abbildungen*, Math. Ann., **136** (1958), 1-8.
7. ———, *Maximale holomorphe und meromorphe Abbildungen, I*, Amer. J. Math., **85** (1963), 298-315.

Received September 27, 1977.

AKITA UNIVERSITY  
AKITA, JAPAN

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)

University of California  
Los Angeles, CA 90024

CHARLES W. CURTIS

University of Oregon  
Eugene, OR 97403

C. C. MOORE

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University  
Stanford, CA 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of *Math. Reviews*, Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).

8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1978 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

Su-Shing Chen, <i>Weak rigidity of compact negatively curved manifolds</i> . . . . .	273
Heinz Otto Cordes and D. A. Williams, <i>An algebra of pseudodifferential operators with nonsmooth symbol</i> . . . . .	279
Herbert Paul Halpern, <i>Normal expectations and integral decomposition of type III von Neumann algebras</i> . . . . .	291
G. Hochschild, <i>On representing analytic groups with their automorphisms</i> . . . . .	333
Dean G. Hoffman and David Anthony Klarner, <i>Sets of integers closed under affine operators—the closure of finite sets</i> . . . . .	337
Simeon Ivanov, <i>On holomorphic relative inverses of operator-valued functions</i> . . . . .	345
O. P. Juneja and M. L. Mogra, <i>Radii of convexity for certain classes of univalent analytic functions</i> . . . . .	359
Hadi Kharaghani, <i>The evolution of bounded linear functionals with application to invariant means</i> . . . . .	369
Jack W. Macki, <i>A singular nonlinear boundary value problem</i> . . . . .	375
A. W. Mason and Walter Wilson Stothers, <i>Remarks on a theorem of L. Greenberg on the modular group</i> . . . . .	385
Kevin Mor McCrimmon, <i>Peirce ideals in Jordan algebras</i> . . . . .	397
John C. Morgan, II, <i>On the absolute Baire property</i> . . . . .	415
Gerard J. Murphy, <i>Commutative non-Archimedean <math>C^*</math>-algebras</i> . . . . .	433
Masafumi Okumura, <i>Submanifolds with <math>L</math>-flat normal connection of the complex projective space</i> . . . . .	447
Chull Park and David Lee Skoug, <i>Distribution estimates of barrier-crossing probabilities of the Yeh-Wiener process</i> . . . . .	455
Irving Reiner, <i>Invariants of integral representations</i> . . . . .	467
Phillip Schultz, <i>The typeset and cotypeset of a rank 2 abelian group</i> . . . . .	503
John Brendan Sullivan, <i>Representations of Witt groups</i> . . . . .	519
Chia-Chi Tung, <i>Equidistribution theory in higher dimensions</i> . . . . .	525
Toshio Uda, <i>Complex bases of certain semiproper holomorphic maps</i> . . . . .	549