

Pacific Journal of Mathematics

**PROBABILITY MEASURES AND THE C -SETS OF
SELIVANOVSKIJ**

STEVEN EUGENE SHREVE

PROBABILITY MEASURES AND THE C-SETS OF SELIVANOVSKIJ

STEVEN E. SHREVE

Let X be a Borel space and $\mathcal{S}(X)$ the smallest σ -field containing the Borel subsets of X and closed under operation (A). Then $\mathcal{S}(X)$ is a sub- σ -field of the class of universally measurable subsets of X . Let $P(X)$ be the space of probability measures on the Borel subsets of X and equip $P(X)$ with the weak topology. It is proved that if $A \in \mathcal{S}(X)$, then $\{p \in P(X): P(A) \geq \lambda\}$ is in $\mathcal{S}[P(X)]$ for every real λ . This allows us to show that if X and Y are Borel spaces, f is an $\mathcal{S}(X \times Y)$ measurable extended-real-valued function, and $x \rightarrow q(\cdot | x)$ is $\mathcal{S}(X)$ measurable from X to $P(Y)$, then $x \rightarrow \int f(x, y)q(dy | x)$ is $\mathcal{S}(X)$ measurable on the set where it is defined. (Measurability is relative to the Borel σ -fields in the range spaces.)

1. Introduction. In the study of sequential optimization, it is often convenient to pose problems in Polish spaces or Borel subsets of Polish spaces (hereafter called *Borel spaces*). Such a space together with its Borel subsets is a well behaved measurable space, but there are certain simple operations which lead one out of the Borel sets into larger σ -fields. One of these is infimization: if $f: X \times Y \rightarrow R^*$ is Borel, where X and Y are Borel spaces and $R^* = R \cup \{\pm \infty\}$ is a Borel space with the usual topology, then

$$(1.1) \quad g(x) = \inf_{y \in Y} f(x, y)$$

may fail to be Borel, but will be analytically measurable, i.e., measurable with respect to the σ -field generated by the analytic sets in X [10]. Another such operation is selection: if $B \subset X \times Y$ is Borel and the projection of B onto X is X , there may fail to exist a Borel measurable function $\phi: X \rightarrow Y$ whose graph lies in B [2], but by a famous result due independently to Jankov [6] and von Neumann [11], a selector ϕ which is analytically measurable can always be found. (See also [8] and [3].) Another selection problem is this. Let g be as in (1.1) and $\varepsilon > 0$. Does there exist a measurable $\phi: X \rightarrow Y$ such that

$$(1.1) \quad f[x, \phi(x)] \leq \begin{cases} g(x) + \varepsilon & \text{if } g(x) > -\infty \\ -\frac{1}{\varepsilon} & \text{if } g(x) = -\infty? \end{cases}$$

Such a selector which is Borel measurable need not exist, but one

which is analytically measurable will. However, if we define $I = \{x \in X \mid g(x) = f(x, y) \text{ for some } y \in Y\}$, and require in addition to (1.2) that

$$(1.3) \quad f[x, \phi(x)] = g(x) \quad \forall x \in I,$$

it is unknown if an analytically measurable selector exists, but one which is the composition of two analytically measurable functions does exist [4]. This question is of particular interest in dynamic programming [3], [15], [16].

Once the decision has been made to consider classes of sets and functions larger than the Borel class, three questions arise. The first is whether the sets in the larger class are universally measurable,¹ the second is whether the class of functions is closed under composition, and the third is to determine if the mapping

$$(1.4) \quad x \longrightarrow \int f(x, y)q(dy|x)$$

is in the class when X and Y are Borel spaces, the extended real-valued function f is in the class, and the mapping $x \rightarrow q(\cdot|x)$ is Borel measurable from X to $P(Y)$, the space of probability measures on the Borel subsets of Y . Note that $P(Y)$ with the weak topology is again a Borel space, and we will always understand it to be equipped with this topology. We will generalize the last question somewhat and ask about the measurability of (1.4) when both f and $q(\cdot|x)$ are measurable with respect to some σ -field larger than the Borel σ -field. The significance of the first two questions is apparent; in fact, we must look at the second to discuss the selector given by [4] mentioned above. The third is of interest in dynamic programming and other sequential optimization problems.

In this paper we deal with $\mathcal{S}(X)$, the smallest σ -field on a Borel space X which contains the Borel sets and is closed under operation (A). This σ -field was studied by Selivanovskij [14] and is known to be contained in the σ -field of universally measurable sets (see, e.g., Saks [13]). Furthermore, all the selection results mentioned above and even much stronger ones are true for $\mathcal{S}(X)$ measurable functions [5]. We say that a mapping f from a Borel space X to a Borel space Y is $\mathcal{S}(X)$ measurable if $f^{-1}(B) \in \mathcal{S}(X)$ for every Borel set $B \subset Y$ and show that this implies $f^{-1}(B) \in \mathcal{S}(X)$ for every $B \in \mathcal{S}(Y)$. This allows us to conclude that the composition of two $\mathcal{S}(X)$ measurable functions is again $\mathcal{S}(X)$ measurable, which is the result of Lemma 2 of [5]. Our method of proving this, however,

¹ A subset A of a Borel space X is said to be *universally measurable* if given any probability measure p on the Borel subsets of X , A is in the completion of the Borel σ -field with respect to p .

differs from that of [5] and shows that $\mathcal{S}(X)$ is the smallest σ -algebra which contains the analytic sets and has this property. We show also that if $A \in \mathcal{S}(X)$, then the mapping $p \rightarrow p(A)$ from $P(X)$ to $[0, 1]$ is $\mathcal{S}[P(X)]$ measurable, and consequently the mapping (1.4) is $\mathcal{S}(X)$ measurable on the set where the integral is defined. The method of proof shows in addition that if A is analytic in X and $0 \leq \lambda \leq 1$, then $\{p: p(A) \geq \lambda\}$ is analytic in $P(X)$, which is a result obtained earlier by different methods in [3] and [10]. A proof of this fact by our method can be found in [17], [18].

2. Souslin schemes. Following Meyer [9], we define a *paving* \mathcal{P} of a space X to be a collection of subsets of X . We denote by \mathcal{P}_σ the collection of sets obtainable as countable unions of sets in \mathcal{P} , and by \mathcal{P}_δ the collection obtainable as countable intersections. The σ -field generated by \mathcal{P} will be denoted $\sigma(\mathcal{P})$.

Let \mathcal{N} be the set of infinite sequences and Σ the set of finite sequences of positive integers. For $s = (\sigma_1, \dots, \sigma_n) \in \Sigma$ and $z = (\zeta_1, \zeta_2, \dots) \in \mathcal{N}$, we write $s < z$ if $\sigma_1 = \zeta_1, \dots, \sigma_n = \zeta_n$. A *Souslin scheme* S for \mathcal{P} is a mapping from Σ into \mathcal{P} . The *nucleus* of S is

$$N(S) = \bigcup_{z \in \mathcal{N}} \bigcap_{s < z} S(s).$$

The nucleus $N(S)$ is called the result of *operation* (A) acting on the sets $\{S(s): s \in \Sigma\}$. We denote by $\mathcal{A}(\mathcal{P})$ the collection of nuclei of Souslin schemes for \mathcal{P} . It is well known that

$$(2.1) \quad \mathcal{P} \subset \mathcal{A}(\mathcal{P}),$$

$$(2.2) \quad \mathcal{A}(\mathcal{P})_\delta = \mathcal{A}(\mathcal{P})_\sigma = \mathcal{A}(\mathcal{P}),$$

$$(2.3) \quad \mathcal{A}(\mathcal{P}) = \mathcal{A}[\mathcal{A}(\mathcal{P})].$$

In general, $\mathcal{A}(\mathcal{P})$ is not a σ -field because it is not closed under complementation. However, if the complement of each set in \mathcal{P} is in $\mathcal{A}(\mathcal{P})$, then $\sigma(\mathcal{P}) \subset \mathcal{A}(\mathcal{P})$. Finally, if X is a Borel space and \mathcal{B} is the collection of Borel subsets of X , then the analytic subsets of X are the members of $\mathcal{A}(\mathcal{B})$.

3. The σ -field $\mathcal{S}(X)$. Let \mathcal{F} be the collection of closed sets in a Borel space X and define by transfinite induction:

$$(3.1) \quad \mathcal{C}_0 = \mathcal{F}$$

$$(3.2) \quad \mathcal{C}_\alpha = \mathcal{A}[\sigma(\bigcup_{\beta < \alpha} \mathcal{C}_\beta)],$$

where α ranges over the ordinals less than the first uncountable ordinal Ω . Note that \mathcal{C}_1 is the collection of analytic subsets of X . We set

$$(3.3) \quad \mathcal{S}(X) = \bigcup_{\alpha < \Omega} \mathcal{C}_\alpha .$$

If S is a Souslin scheme for $\mathcal{S}(X)$, then since S has a countable range, there exists an $\alpha < \Omega$ such that S is a Souslin scheme for \mathcal{C}_α . Therefore $N(S) \in \mathcal{C}_{\alpha+1}$, so $\mathcal{S}(X) = \mathcal{A}[\mathcal{S}(X)]$. Since $\mathcal{S}(X)$ is closed under complementation, $\mathcal{A}[\mathcal{S}(X)]$ must contain $\sigma[\mathcal{S}(X)]$, so $\mathcal{S}(X)$ is a σ -field; indeed, it is the smallest σ -field containing \mathcal{S} and closed under operation (A). It is also true that if X is uncountable and $\beta < \alpha < \Omega$, then \mathcal{C}_β is a proper subset of \mathcal{C}_α . Although there seems to be no easily accessible proof of this in the literature, a discussion of the method of proof using universal sets or universal functions can be found in [1] and [7].

If (X, \mathcal{D}) is a measurable space and Y is a Borel space, we say $f: X \rightarrow Y$ is \mathcal{D} measurable if $f^{-1}(B) \in \mathcal{D}$ for every Borel set B in Y . We are interested in the following two theorems.

THEOREM 1. *If X and Y are Borel spaces and $f: X \rightarrow Y$ is $\mathcal{S}(X)$ measurable, then for every $\mathcal{S}(Y)$ set B , we have $f^{-1}(B) \in \mathcal{S}(X)$. Equivalently, if X, Y , and Z are Borel spaces and $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are $\mathcal{S}(X)$ and $\mathcal{S}(Y)$ measurable respectively, then $g \circ f$ is $\mathcal{S}(X)$ measurable. Furthermore, if \mathcal{D} is a σ -field on X containing the analytic sets such that the \mathcal{D} measurability of $f: X \rightarrow X$ and $g: X \rightarrow X$ implies the \mathcal{D} measurability of $g \circ f$, then \mathcal{D} contains $\mathcal{S}(X)$.*

THEOREM 2. *If A is a \mathcal{C}_α set in a Borel space X and $0 \leq \lambda \leq 1$, then $\{p: p(A) > \lambda\}$ and $\{p: p(A) \geq \lambda\}$ are \mathcal{C}_α sets in $P(X)$. In particular, when A is analytic, $\{p: p(A) > \lambda\}$ and $\{p: p(A) \geq \lambda\}$ are analytic.*

COROLLARY. *Let X and Y be Borel spaces, let $f: X \times Y \rightarrow R^*$ be $\mathcal{S}(X \times Y)$ measurable, and let $x \rightarrow q(\cdot | x)$ be an $\mathcal{S}(X)$ measurable map into $P(Y)$. Then the set*

$$C = \left\{ x \in X \mid \int f^+(x, y)q(dy|x) = \int f^-(x, y)q(dy|x) = \infty \right\}$$

is $\mathcal{S}(X)$ measurable, and if for some $a \in R^$ we define*

$$I(x) = \begin{cases} \int f(x, y)q(dy|x), & x \in C, \\ a, & x \in C^c, \end{cases}$$

then I is $\mathcal{S}(X)$ measurable.

Proof of Theorem 1. We show first that if X and Y are Borel

spaces, $f: X \rightarrow Y$ is $\mathcal{S}(X)$ measurable, and $B \in \mathcal{S}(Y)$, then $f^{-1}(B) \in \mathcal{S}(X)$. By definition, $B \in \mathcal{C}_\alpha$ for some $\alpha < \Omega$, and the proof is by transfinite induction on α . If $\alpha = 0$, then $f^{-1}(B) \in \mathcal{S}(X)$. If the result holds for all $\beta < \alpha$, then $f^{-1}(C) \in \mathcal{S}(X)$ for every $C \in \sigma(\bigcup_{\beta < \alpha} \mathcal{C}_\beta)$. Since $B = \bigcup_{z \in \mathcal{N}} \bigcap_{s < z} S(s)$, where S is some Souslin scheme for $\sigma(\bigcup_{\beta < \alpha} \mathcal{C}_\beta)$, we have

$$f^{-1}(B) = \bigcup_{z \in \mathcal{N}} \bigcap_{s < z} f^{-1}[S(s)] \in \mathcal{N}[\mathcal{S}(X)] = \mathcal{S}(X).$$

It is straightforward to verify that the second statement of the theorem is equivalent to the first. Under the assumptions of the last part of the theorem, we show that $\mathcal{N}(\mathcal{D}) = \mathcal{D}$. Since $\mathcal{S}(X)$ is the smallest σ -field containing the analytic sets and closed under operation (A), this will imply $\mathcal{S}(X) \subset \mathcal{D}$. We may restrict attention to uncountable X , and since every such X is Borel isomorphic to the space of sequences of zeroes and ones with the topology of componentwise convergence [12], we may assume X has this form. Let S be a Souslin scheme for \mathcal{D} , let ψ be a one-to-one function from the positive integers onto \mathcal{N} , and let $f: X \rightarrow X$ be defined so that the k th component of $f(x)$ is the indicator of $S[\psi(k)]$. Define $R(k) = \{(\xi_1, \xi_2, \dots) \in X: \xi_k = 1\}$, and define $g(x) = x_1$ if $x \in \bigcup_{z \in \mathcal{N}} \bigcap_{s < z} R[\psi^{-1}(s)]$ and $g(x) = x_2 \neq x_1$ otherwise. It is easy to verify that f is \mathcal{D} measurable, g is measurable with respect to the σ -field generated by the analytic sets, and $f^{-1}[g^{-1}(x_i)] = N(S)$. It follows that $N(S) \in \mathcal{D}$, so \mathcal{D} is closed under operation (A) as claimed.

Proof of Theorem 2. We proceed by transfinite induction to show that if A is a \mathcal{C}_α set in X and $0 \leq \lambda \leq 1$, then $\{p \in P(X): p(A) \geq \lambda\}$ is a \mathcal{C}_α set in $P(X)$. If $\alpha = 0$, this follows from Theorem II.6.1(c) of [12]. If the result holds for all $\beta < \alpha$, then for $C \in \sigma(\bigcup_{\beta < \alpha} \mathcal{C}_\beta)$, the mapping $p \rightarrow p(C)$ is $\sigma(\bigcup_{\beta < \alpha} \mathcal{C}_\beta)$ measurable by a Dynkin system argument. Now let S be a Souslin scheme for $\sigma(\bigcup_{\beta < \alpha} \mathcal{C}_\beta)$ (in X) and set $A = N(S)$. For $(\sigma_1, \dots, \sigma_k) \in \mathcal{N}$, define

$$(3.4) \quad M(\sigma_1, \dots, \sigma_k) = \{(\zeta_1, \zeta_2, \dots) \in \mathcal{N}: \tau_i \leq \sigma_i, \dots, \tau_k \leq \sigma_k\},$$

$$(3.5) \quad R(\sigma_1, \dots, \sigma_k) = \bigcup_{z \in M(\sigma_1, \dots, \sigma_k)} \bigcap_{s < z} S(s),$$

$$(3.6) \quad K(\sigma_1, \dots, \sigma_k) = \bigcup_{\tau_1 \leq \sigma_1, \dots, \tau_k \leq \sigma_k} \bigcap_{j=1}^k S(\tau_1, \dots, \tau_j).$$

As $\sigma_1 \uparrow \infty$, $R(\sigma_1) \uparrow A$, and as $\sigma_k \uparrow \infty$, $R(\sigma_1, \dots, \sigma_k) \uparrow R(\sigma_1, \dots, \sigma_{k-1})$.

Suppose $p \in P(X)$ is such that $p(A) \geq \lambda$ and let n be a positive integer. Choose ζ_1, ζ_2, \dots such that

$$p(A) \leq p^*[R(\bar{\zeta}_1)] + \frac{1}{2n}$$

$$p^*[R(\bar{\zeta}_1, \dots, \bar{\zeta}_{k-1})] \leq p^*[R(\bar{\zeta}_1, \dots, \bar{\zeta}_{k-1}, \bar{\zeta}_k)] + \frac{1}{2^k n},$$

where p^* denotes p outer measure. Then

$$p(A) \leq p^*[R(\bar{\zeta}_1, \dots, \bar{\zeta}_k)] + \frac{1}{n} \text{ for all } k,$$

and since $R(\bar{\zeta}_1, \dots, \bar{\zeta}_k) \subset K(\bar{\zeta}_1, \dots, \bar{\zeta}_k)$ and the latter set is p measurable, we have

$$\lambda \leq p(A) \leq p[K(\bar{\zeta}_1, \dots, \bar{\zeta}_k)] + \frac{1}{n} \text{ for all } k.$$

Thus we may write

$$(3.7) \quad \{p \in P(X) \mid p(A) \geq \lambda\} \subset \bigcap_{n=1}^{\infty} \bigcup_{z \in \mathcal{N}} \bigcap_{s < z} \left\{ p \in P(X) : p[K(s)] \geq \lambda - \frac{1}{n} \right\}.$$

We show the reverse of set containment (3.7). We begin with a proof found in Saks [13] that

$$(3.8) \quad \bigcap_{k=1}^{\infty} K(\zeta_1, \dots, \zeta_k) \subset A \text{ for all } (\zeta_1, \zeta_2, \dots) \in \mathcal{N}.$$

To show (3.8), choose $x \in \bigcap_{k=1}^{\infty} K(\zeta_1, \dots, \zeta_k)$, i.e.,

$$(3.9) \quad x \in \bigcap_{k=1}^{\infty} \bigcup_{\tau_1 \leq \zeta_1, \dots, \tau_k \leq \zeta_k} \bigcap_{j=1}^k S(\tau_1, \dots, \tau_j).$$

An argument by contradiction will be used to show that for some $\bar{\tau}_1 \leq \zeta_1$, we have $x \in S(\bar{\tau}_1)$ and

$$(3.10) \quad x \in \bigcap_{k=2}^{\infty} \bigcup_{\tau_2 \leq \zeta_2, \dots, \tau_k \leq \zeta_k} \bigcap_{j=2}^k S(\bar{\tau}_1, \tau_2, \dots, \tau_j).$$

If no such $\bar{\tau}_1$ existed, then for every $\tau_1 \leq \zeta_1$, there would exist a positive integer $k(\tau_1)$ such that

$$x \notin S(\tau_1) \cap \left[\bigcap_{k=2}^{k(\tau_1)} \bigcup_{\tau_2 \leq \zeta_2, \dots, \tau_k \leq \zeta_k} \bigcap_{j=2}^k S(\tau_1, \tau_2, \dots, \tau_j) \right].$$

If $\bar{k} = \max_{\tau_1 \leq \zeta_1} k(\tau_1)$, then

$$\begin{aligned} x &\notin \bigcup_{\tau_1 \leq \zeta_1} \left\{ S(\tau_1) \cap \left[\bigcap_{k=2}^{\bar{k}} \bigcup_{\tau_2 \leq \zeta_2, \dots, \tau_k \leq \zeta_k} \bigcap_{j=2}^k S(\tau_1, \tau_2, \dots, \tau_j) \right] \right\} \\ &\supset \bigcup_{\tau_1 \leq \zeta_1, \dots, \tau_{\bar{k}} \leq \zeta_{\bar{k}}} \bigcap_{j=1}^{\bar{k}} S(\tau_1, \tau_2, \dots, \tau_j) \\ &= K(\zeta_1, \zeta_2, \dots, \zeta_{\bar{k}}), \end{aligned}$$

and a contradiction is reached. Because (3.10) is of the same form as (3.9), we can apply this argument repeatedly to obtain $\bar{\tau}_1 \leq \zeta_1$, $\bar{\tau}_2 \leq \zeta_2, \dots$ such that

$$x \in \bigcap_{k=1}^{\infty} S(\bar{\tau}_1, \dots, \bar{\tau}_k) \subset A,$$

and (3.8) is proved.

Now choose

$$p \in \bigcap_{n=1}^{\infty} \bigcup_{z \in \mathcal{N}} \bigcap_{s < z} \left\{ q \in P(X) : q[K(s)] \geq \lambda - \frac{1}{n} \right\}.$$

For each n there exists $(\zeta_1, \zeta_2, \dots) \in \mathcal{N}$ such that

$$(3.11) \quad p[K(\zeta_1, \dots, \zeta_k)] \geq \lambda - \frac{1}{n} \quad \text{for all } k.$$

Since the sets $K(\zeta_1, \dots, \zeta_k)$ decrease with increasing k , (3.11) implies

$$p\left[\bigcap_{k=1}^{\infty} K(\zeta_1, \dots, \zeta_k)\right] \geq \lambda - \frac{1}{n},$$

and this together with (3.8) tells us that $p(A) \geq \lambda - 1/n$. Since n is arbitrary, $p(A) \geq \lambda$. This shows that equality holds in (3.7), i.e.,

$$\{p : p(A) \geq \lambda\} = \bigcap_{n=1}^{\infty} N(S_n),$$

where S_n is the Souslin scheme defined by

$$S_n(s) = \left\{ p \in P(X) : p[K(s)] \geq \lambda - \frac{1}{n} \right\}.$$

Since $K(s) \in \sigma(\bigcup_{\beta < \alpha} \mathcal{C}_\beta)$ (in X) for each s , $S_n(s) \in \sigma(\bigcup_{\beta < \alpha} \mathcal{C}_\beta)$ (in $P(X)$) by the induction hypothesis, so $\{p : p(A) \geq \lambda\} \in \mathcal{C}_\alpha$ by (2.2). Note that

$$\{p : p(A) > \lambda\} = \bigcup_{n=1}^{\infty} \left\{ p : p(A) \geq \lambda + \frac{1}{n} \right\},$$

and use (2.2) to conclude that this is also a \mathcal{C}_α set.

Proof of Corollary. Suppose f is the indicator of an $\mathcal{S}(X \times Y)$ set A . For $x \in X$, let $p_x \in P(X)$ be the unit point mass at x , and for $p \in P(X)$, $q \in P(Y)$, let $qp \in P(X \times Y)$ be defined on Borel sets B by

$$(qp)(B) = \int q\{y : (x, y) \in B\} p(dx).$$

Then I is the composition of the mappings $x \rightarrow (p_x, q(\cdot | x))$, $(p, q) \rightarrow qp$ and $r \rightarrow r(A)$, where the last mapping is from $P(X \times Y)$ to $[0, 1]$ and is $\mathcal{S}(X \times Y)$ measurable by Theorem 2. The $\mathcal{S}(X)$ measurability of I follows from Theorem 1. The extension to general f is straightforward.

The author wishes to thank John Addison for pointing out the work of Selivanovskij [14].

REFERENCES

1. J. Addison, *Current problems in descriptive set theory*, Proc. Symposia Pure Math., **13** (1974), 1-10.
2. D. Blackwell, *A Borel set not containing a graph*, Ann. Math. Statist., **39** (1968), 1345-1347.
3. D. Blackwell, D. Freedman, and M. Orkin, *The optimal reward operator in dynamic programming*, Ann. Probability, **2** (1974), 926-941.
4. L. D. Brown and R. Purves, *Measurable selections of extrema*, Ann. Statist., **1** (1973), 902-912.
5. D. Cenzer and R. D. Mauldin, *Measurable parameterizations and selections*, to appear in Trans. Amer. Math. Soc.
6. B. Jankov, *On the uniformisation of A -sets*, Doklady Acad. Nauk. SSSR, **30** (1941), 591-592 (in Russian).
7. K. Kuratowski, *Topology I*, Academic Press, New York, 1966.
8. G. Mackey, *Borel structure in groups and their duals*, Trans. Amer. Math. Soc., **85** (1957), 134-165.
9. P. A. Meyer, *Probability and Potentials*, Blaisdell, Waltham, Mass., 1966.
10. P. A. Meyer and M. Traki, *Reduites et jeux de hasard*, Sem. Prob. VII, Univ. Strasbourg, Lecture Notes in Mathematics 321, 155-171, Springer, Berlin, 1973.
11. J. von Neumann, *On rings of operators. Reduction theory*, Ann. Math., **50** (1949), 401-485.
12. K. Parthasarathy, *Probability Measures on Metric Spaces*, Academic Press, New York, 1967.
13. S. Saks, *Theory of the Integral*, Stechert, New York, 1937.
14. E. Selivanovskij, *Ob odnom klasse effektivnyh mnozestv (mnozestva C)*, Matematicheskij Sbornik, **35** (1928), 379-413.
15. S. Shreve and D. P. Bertsekas, *Alternative theoretical frameworks for finite horizon stochastic optimal control*, SIAM J. Control and Optimization, **16** (1978), 953-978.
16. ———, *Universally measurable policies in dynamic programming*, to appear in Math. Operations Research.
17. Barbaroshie, *On the theory of controlled Markov processes*, Theory Prob. Appl., **22** (1977), 53-69.
18. A. K. Zvonkin, *On sequentially controlled Markov Processes*, Matematicheskij Sbornik, **86** (128) (1971), 611-621 (in Russian).

Received March 20, 1978 and in revised form June 12, 1978. This work was supported in part by the National Science Foundation under Grant NSF ENG 74-19332.

UNIVERSITY OF DELAWARE
NEWARK, DE 19711

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

C. W. CURTIS

University of Oregon
Eugene, OR 97403

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Teófilo Abuabara, <i>A remark on infinitely nuclearly differentiable functions</i>	1
David Fenimore Anderson, <i>Projective modules over subrings of $k[X, Y]$ generated by monomials</i>	5
Joseph Barback and Thomas Graham McLaughlin, <i>On the intersection of regressive sets</i>	19
Murray Bell, John Norman Ginsburg and R. Grant Woods, <i>Cardinal inequalities for topological spaces involving the weak Lindelof number</i>	37
Laurence Richard Boxer, <i>The space of ANRs of a closed surface</i>	47
Zvonko Cerin, <i>Homotopy properties of locally compact spaces at infinity-calmness and smoothness</i>	69
Isidor Fleischer and Ivo G. Rosenberg, <i>The Galois connection between partial functions and relations</i>	93
John R. Giles, David Allan Gregory and Brailey Sims, <i>Geometrical implications of upper semi-continuity of the duality mapping on a Banach space</i>	99
Troy Lee Hicks, <i>Fixed-point theorems in locally convex spaces</i>	111
Hugo Junghenn, <i>Almost periodic functions on semidirect products of transformation semigroups</i>	117
Victor Kaftal, <i>On the theory of compact operators in von Neumann algebras. II</i>	129
Haynes Miller, <i>A spectral sequence for the homology of an infinite delooping</i>	139
Sanford S. Miller, Petru T. Mocanu and Maxwell O. Reade, <i>Starlike integral operators</i>	157
Stanley Stephen Page, <i>Regular FPF rings</i>	169
Ghan Shyam Pandey, <i>Multipliers for C, 1 summability of Fourier series</i>	177
Shigeo Segawa, <i>Bounded analytic functions on unbounded covering surfaces</i>	183
Steven Eugene Shreve, <i>Probability measures and the C-sets of Selivanovskij</i>	189
Tor Skjelbred, <i>Combinatorial geometry and actions of compact Lie groups</i>	197
Alan Sloan, <i>A note on exponentials of distributions</i>	207
Colin Eric Sutherland, <i>Type analysis of the regular representation of a nonunimodular group</i>	225
Mark Phillip Thomas, <i>Algebra homomorphisms and the functional calculus</i>	251
Sergio Eduardo Zarantonello, <i>A representation of H^p-functions with $0 < p < \infty$</i>	271