

Pacific Journal of Mathematics

**APPROXIMATION BY RATIONAL MODULES ON NOWHERE
DENSE SETS**

JAMES LI-MING WANG

APPROXIMATION BY RATIONAL MODULES ON NOWHERE DENSE SETS

JAMES LI-MING WANG

Let X be a compact subset of the complex plane. Let the module $\mathcal{R}(X)\tilde{\mathcal{P}}_m$ be the space of all functions of the form

$$r_0(z) + r_1(z)\bar{z} + \cdots + r_m(z)\bar{z}^m$$

where each r_i is a rational function with poles off X . We prove that $\mathcal{R}(X)\tilde{\mathcal{P}}_1$ is dense in $L^p(X)$ for all $1 \leq p < \infty$ and $\mathcal{R}(X)\tilde{\mathcal{P}}_2$ is dense in $\mathcal{C}(X)$ if X has no interior point. As corollaries, we also prove that $\mathcal{R}(X)\tilde{\mathcal{P}}_2$ is dense in $\text{lip}(\alpha, X)$ for all $0 < \alpha < 1$ and $\mathcal{R}(X)\tilde{\mathcal{P}}_3$ is dense in $D^1(X)$ for the same X .

1. **Introductions.** Let X be a compact subset of the complex plane. Let the module $\mathcal{R}(X)\tilde{\mathcal{P}}_m$ be the space $\mathcal{R} + \mathcal{R}\bar{z} + \cdots + \mathcal{R}\bar{z}^m$

$$= \{r_0(z) + r_1(z)\bar{z} + \cdots + r_m(z)\bar{z}^m\},$$

where each r_i is a rational function with poles off X . In [3, 4], O'Farrell has studied the relation of the problems of approximation by rational modules in different Lipschitz norms, and in the uniform norm, etc., to one another.

In this note, we investigate the problem of determining the set X so that $\mathcal{R}(X)\tilde{\mathcal{P}}_m$ is uniformly dense in $\mathcal{C}(X)$ for each m .

Vitushkin [8] has given a necessary and sufficient condition in terms of analytic capacities for the case $m = 0$. In [3], O'Farrell has given an example of an X such that $\mathcal{R}(X)\tilde{\mathcal{P}}_1$ is uniformly dense in $\mathcal{C}(X)$ whereas $\mathcal{R}(X)$ fails to be dense in $\mathcal{C}(X)$.

It is apparent that if X has interior, then $\mathcal{R}(X)\tilde{\mathcal{P}}_m$ can not be dense in $\mathcal{C}(X)$. Thus we restrict our attention to a compact set X without interior throughout this note. Let $L^p(X) = L^p(\chi_X dm)$, where dm denotes the 2-dimensional Lebesgue measure. We prove the following theorem:

THEOREM. *Let X be a compact set with no interior. Then*

- I. $\mathcal{R}(X)\tilde{\mathcal{P}}_1$ is dense in $L^p(X)$ for all $1 \leq p < \infty$, and
- II. $\mathcal{R}(X)\tilde{\mathcal{P}}_2$ is dense in $\mathcal{C}(X)$.

2. *Proof of theorem.* Let μ be a (finite Borel) measure on X . The Cauchy transform $\hat{\mu}$ is defined by

$$\hat{\mu}(z) = \int_x \frac{d\mu(\zeta)}{\zeta - z}.$$

It is well known that $\hat{\mu}$ is absolutely convergent for almost all z (dm) and $\hat{\mu}$ belongs to L^p_{loc} for $1 \leq p < 2$ (see [2], p. 37). If g is a function on X , we write \hat{g} for $\widehat{gd\mu}$. Sinanjan [6] was the first to prove that $\mathcal{R}(X)$ is dense in $L^p(X)$ when $1 \leq p < 2$. However, it is Brennan's [1] proof that leads to the results we obtain here.

We use the symbol $\bar{\partial}$ for the operator $\partial/\partial x + i(\partial/\partial y)$ and write $g \perp \mathcal{R}(X)\bar{\mathcal{P}}_m$ if $\int fgdm = 0$ for all $f \in \mathcal{R}(X)\bar{\mathcal{P}}_m$. The following lemma is a special case of the key lemma in [3].

LEMMA. *Let μ be a measure on X . Then $\mu \perp \mathcal{R}(X)\bar{\mathcal{P}}_{m+1}$ if and only if $\hat{\mu} \perp \mathcal{R}(X)\bar{\mathcal{P}}_m$.*

Proof. Because $\int fd\mu = \pi^{-1} \int (\bar{\partial}f)\hat{\mu}dm$ for all $f \in \mathcal{R}(X)\bar{\mathcal{P}}_{m+1}$ (see [2]).

Proof of theorem. Of course, the case $1 \leq p < 2$ is contained in Sinanjan's theorem.

Let $g \in L^p(X)$, $q > 1$, $p^{-1} + q^{-1} = 1$, such that $g \perp \mathcal{R}(X)\bar{\mathcal{P}}_1$. Lemma implies $\hat{g} \perp \mathcal{R}(X)$. In particular, $\hat{g} = 0$ if $z \notin X$. Being the convolution of g and $\zeta^{-1} \in L^r_{loc}$ ($1 \leq r < 2$), \hat{g} belongs to $L^s(X)$ for some $s > 2$, by the classical Young's inequality (see [7], p. 271). And being the convolution of \hat{g} and ζ^{-1} , \hat{g} is continuous everywhere in the plane. It follows that $\hat{g} \equiv 0$ since X has no interior. It is well known that this implies $\hat{g} = 0$ in $L^s(X)$, which in turn implies $g = 0$ in $L^q(X)$ and hence $\mathcal{R}(X)\bar{\mathcal{P}}_1$ is dense in $L^p(X)$. Part I is proved.

Let μ be a measure on X such that $\mu \perp \mathcal{R}(X)\bar{\mathcal{P}}_2$. Lemma implies $\hat{\mu} \perp \mathcal{R}(X)\bar{\mathcal{P}}_1$ and Part I implies $\hat{\mu} = 0$ and hence $\mu = 0$. Part II is proved.

3. Some remarks. Let \mathcal{E} and \mathcal{D} be the usual linear topological space of complex-valued \mathcal{C}^∞ functions on the complex plane. Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be two norms on \mathcal{D} and \mathcal{E} respectively. $\mathcal{R}(X)\bar{\mathcal{P}}_m$ can be regarded as a subspace of $\mathcal{D}|_X$ and $\mathcal{E}|_X$. We denote the closure (i.e., the completion) of $\mathcal{R}(X)\bar{\mathcal{P}}_m$ with respect to the norm $\|\cdot\|_j$ (restricted to X) by $[\mathcal{R}(X)\bar{\mathcal{P}}_m]_j$, for $j = 1, 2$. O'Farrell's theorems in [3] can be roughly stated in the following general form:

If $\phi \rightarrow \hat{\phi}$ is continuous from $(\mathcal{D}, \|\cdot\|_1)$ to $(\mathcal{E}, \|\cdot\|_2)$, then

$[\mathcal{R}(X)\bar{\mathcal{P}}_m]_1 = [\mathcal{D}]_1$ implies $[\mathcal{R}(X)\bar{\mathcal{P}}_{m+1}]_2 = [\mathcal{E}]_2$.

As corollaries, we have the following:

COROLLARY. *Let X be a compact set with no interior. Then*

- I. $\mathcal{R}(X)\bar{\mathcal{P}}_2$ is dense in $\text{lip}(\alpha, X)$ for all $0 < \alpha < 1$, and
- II. $\mathcal{R}(X)\bar{\mathcal{P}}_3$ is dense in $D^1(X)$.

For the definitions of $\text{lip}(\alpha, X)$ and $D^1(X)$, we refer the reader to [3].

It is not known whether $\mathcal{R}(X)\bar{\mathcal{P}}_1$ is dense in $\mathcal{C}(X)$ for all X without interior. The annular Swiss Cheese of A. Roth [5] has the property that the continuous function $|z|^2$ is not in the uniform closure of $\mathcal{R}(X)$. However, it is not clear whether or not the function $|z|$ is in the uniform closure of $\mathcal{R}(X)\bar{\mathcal{P}}_1$.

REFERENCES

1. J. Brennan, *Invariant subspaces and rational approximation*, J. Functional Analysis, **7** (1971), 285-310.
2. J. Garnett, *Analytic capacity and measure*, Springer Lecture Notes in Math., No. 297, 1973.
3. A. O'Farrell, *Annihilators of rational modules*, J. Functional Analysis, **19** (1975), 373-389.
4. ———, *Hausdorff content and rational approximation in fractional Lipschitz norms*, Trans. Amer. Math. Soc., **228** (1977), 187-206.
5. A. Roth, *Approximationseigenschaften und Strahlengrenzwerte meromorpher und ganzer Funktionen*, Comment. Math. Helv., **11** (1938), 77-125.
6. S. O. Sinanjan, *The uniqueness property of analytic functions on closed sets without interior points*, Sibirsk Mat. Z., **6** (1965), 1365-1381.
7. E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton, 1970.
8. A. G. Vitushkin, *Analytic capacity of sets and problems in approximation theory*, Russian Math. Survey, **22** (1967), 139-200.

Received May 17, 1978 and in revised form August 10, 1978. Research supported in part by NSF Grant MCS 77-02208.

UNIVERSITY OF ALABAMA
UNIVERSITY, AL 35486

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

C. W. CURTIS

University of Oregon
Eugene, OR 97403

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Jeroen Bruijning and Jun-iti Nagata, <i>A characterization of covering dimension by use of $\Delta_k(X)$</i>	1
John J. Buoni and Albert Jonathan Klein, <i>On the generalized Calkin algebra</i>	9
Thomas Ashland Chapman, <i>Homotopy conditions which detect simple homotopy equivalences</i>	13
John Albert Chatfield, <i>Solution for an integral equation with continuous interval functions</i>	47
Ajit Kaur Chilana and Ajay Kumar, <i>Spectral synthesis in Segal algebras on hypergroups</i>	59
Lung O. Chung, Jiang Luh and Anthony N. Richoux, <i>Derivations and commutativity of rings</i>	77
Michael George Cowling and Paul Rodway, <i>Restrictions of certain function spaces to closed subgroups of locally compact groups</i>	91
David Dixon, <i>The fundamental divisor of normal double points of surfaces</i>	105
Hans Georg Feichtinger, Colin C. Graham and Eric Howard Lakien, <i>Nonfactorization in commutative, weakly selfadjoint Banach algebras</i>	117
Michael Freedman, <i>Cancelling 1-handles and some topological imbeddings</i>	127
Frank E., III Gerth, <i>The Iwasawa invariant μ for quadratic fields</i>	131
Maurice Gilmore, <i>Three-dimensional open books constructed from the identity map</i>	137
Stanley P. Gudder, <i>A Radon-Nikodým theorem for $*$-algebras</i>	141
Peter Wamer Harley, III and George Frank McNulty, <i>When is a point Borel?</i>	151
Charles Henry Heiberg, <i>Fourier series with bounded convolution powers</i>	159
Rebecca A. Herb, <i>Characters of averaged discrete series on semisimple real Lie groups</i>	169
Hideo Imai, <i>On singular indices of rotation free densities</i>	179
Sushil Jajodia, <i>On 2-dimensional CW-complexes with a single 2-cell</i>	191
Herbert Meyer Kamowitz, <i>Compact operators of the form uC_φ</i>	205
Matthew Liu and Billy E. Rhoades, <i>Some properties of the Chebyshev method</i>	213
George Edgar Parker, <i>Semigroups of continuous transformations and generating inverse limit sequences</i>	227
Samuel Murray Rankin, III, <i>Oscillation results for a nonhomogeneous equation</i>	237
Martin Scharlemann, <i>Transverse Whitehead triangulations</i>	245
Gary Joseph Sherman, <i>A lower bound for the number of conjugacy classes in a finite nilpotent group</i>	253
Richard Arthur Shoop, <i>The Lebesgue constants for (f, d_n)-summability</i>	255
Stuart Jay Sidney, <i>Functions which operate on the real part of a uniform algebra</i>	265
Tim Eden Traynor, <i>The group-valued Lebesgue decomposition</i>	273
Tavan Thomas Trent, <i>$H^2(\mu)$ spaces and bounded point evaluations</i>	279
James Li-Ming Wang, <i>Approximation by rational modules on nowhere dense sets</i>	293