

Pacific Journal of Mathematics

**IRREDUCIBLE LENGTHS OF TRIVECTORS OF RANK SEVEN
AND EIGHT**

ROY WESTWICK

IRREDUCIBLE LENGTHS OF TRIVECTORS OF RANK SEVEN AND EIGHT

R. WESTWICK

We determine the irreducible length of complex trivectors of rank less than or equal to eight. The irreducible length is invariant under the action induced by the general linear group on the underlying complex vector space. A classification under this action is available where representatives are explicitly given for each equivalence class and it is the lengths of these representatives which are determined.

In their paper [1], Busemann and Glassco consider the problem of determining the maximal irreducible length (called *length* from now on) $N(F, n, r)$ of r -vectors in $A^r U$ where U is an n -dimensional vector space over the field F . The length of an r -vector is the number of decomposable summands (blades in J. Schouten's book [5] and paper [6]) in a shortest possible representation of that r -vector. In [1] Busemann and Glassco state that "The values $N(C, 7, 3) = 5$, $N(C, 8, 3) = 7$, and $N(C, 9, 3) = 10$ have been claimed but questioned, see Schouten [3, p. 27] and [1]." The purpose of this paper is to show that $N(C, 8, 3) = 5$ (and not 7 as claimed) by determining the lengths of each of the representatives of the Gurevich classification in [2]. For sake of completeness the lengths of the rank 7 trivectors are also included. That $N(C, 7, 4) = 4$ (and not 5) is shown in [7].

Let U be a fixed 8-dimensional vector space over the complex number field and $A^3 U$ the space of trivectors considered. If $X \in A^3 U$ then $[X]$ denotes the intersection of all subspaces W of U for which $X \in A^3 W$. Then $\dim [X]$ is the rank of X . The letters a, b, c, q, r, s, p, t appearing in the Gurevich classification may be assumed to be independent vectors in U . Note that in [2] the "A" has been suppressed.

The equivalence classes are as follows.

- I: 0
- II: [abc]
- III: [aqp] + [brp]
- IV: [aqr] + [brp] + [cpq]
- V: [abc] + [pqr]
- VI: [aqp] + [brp] + [csp]
- VII: [qrs] + [aqp] + [brp] + [csp]
- VIII: [abc] + [qrs] + [aqp]

- IX: $[abc] + [qrs] + [aqp] + [brp]$
 X: $[abc] + [qrs] + [aqp] + [brp] + [csp]$
 XI: $[brp] + [csp] + [aqp] + [crt]$
 XII: $[qrs] + [brp] + [csp] + [aqp] + [crt]$
 XIII: $[abc] + [qrs] + [aqp] + [crt]$
 XIV: $[abc] + [qrs] + [aqp] + [brp] + [crt]$
 XV: $[abc] + [qrs] + [aqp] + [brp] + [csp] + [crt]$
 XIV: $[aqp] + [bst] + [crt]$
 XVII: $[aqp] + [brp] + [bst] + [crt]$
 XVIII: $[qrs] + [aqp] + [brp] + [bst] + [crt]$
 XIX: $[aqp] + [brp] + [csp] + [bst] + [crt]$
 XX: $[qrs] + [aqp] + [brp] + [csp] + [bst] + [crt]$
 XXI: $[abc] + [qrs] + [aqp] + [bst] + [crt]$
 XXII: $[abc] + [qrs] + [aqp] + [brp] + [bst] + [crt]$
 XXIII: $[abc] + [qrs] + [aqp] + [brp] + [csp] + [bst] + [crt]$

Since rank and length remain constant in an equivalence class the terms will be used on the equivalence class itself. There are five equivalence classes of trivectors with rank 7. Three of them, namely VI, VIII, and IX, have length 3 and the other two, VII and X, have length 4. There are thirteen equivalence classes with rank 8. Two of them, XVI and XIX have length 3; one of them, XV has length 5; and the remaining 10 classes have length 4. The results are proved as follows.

Consider first a trivector X of rank 7. If it has length 3 then $X = X_1 + X_2 + X_3$ where each X_i is decomposable. If $[X_1] \cap [X_2] \neq 0$ then we may write $X_1 + X_2 = x_1 \wedge (x_2 \wedge x_3 + x_4 \wedge x_5)$ where x_1, \dots, x_5 are independent vectors in U . Then $X_3 = u \wedge x_6 \wedge x_7$ where $u \in \langle x_1, \dots, x_5 \rangle$ and x_1, \dots, x_7 are independent vectors in U . If u is a multiple of x_1 then X has the form VI where $p = x_1$. If u is not a multiple of x_1 then by rewriting $X_1 + X_2$ we may assume that $u = x_2$. (For if $u = \alpha x_1 + \beta x_2 + w$ where $\beta \neq 0$ and $w \in \langle x_3, x_4, x_5 \rangle$ then $X_1 + X_2 = x_1 \wedge (u \wedge \beta^{-1}x_3 - \beta^{-1}w \wedge x_3 + x_4 \wedge x_5)$ and $-\beta^{-1}w \wedge x_3 + x_4 \wedge x_5$ is decomposable.) Then X has the form VIII. If $[X_i] \cap [X_j] = 0$ for all pairs $i \neq j$ then we may write $X + X_2 = x_1 \wedge x_2 \wedge x_3 + x_4 \wedge x_5 \wedge x_6$ and $X_3 = u \wedge v \wedge x_7$ where $u, v \in \langle x_1, \dots, x_6 \rangle$ and x_1, \dots, x_7 are independent. Then $u = u_1 + u_2, v = v_1 + v_2$ where $u_1, v_1 \in [X_1]$ and $u_2, v_2 \in [X_2]$ so by refactoring X_1 and X_2 we may assume that $u = x_1 + x_4$ and $v = x_2 + x_5$. Then X has the form XIX which is equal to $[b(c + p)\alpha] + [qr(x + p)] + [(q - b)p(\alpha + r)]$ where $x_1 = -b, x_2 = \alpha, x_3 = c + p, x_4 = q, x_5 = r, x_6 = s + p, x_7 = -p$.

Since all possibilities for rank 7 length 3 trivectors have been considered the remaining classes have length at least 4. Equivalence class VII then has length 4 and X, which is equal to $([aqp] + [(b + s)(r - c)p] + [(a + p)bc] + [(p + q)rs])$ has length 4 also. This takes care of the rank 7 trivectors.

Suppose $X = X_1 + X_2 + X_3$ is a trivector of length 3 and rank 8, where each X_i is decomposable. If $[X_1] \cap [X_2] \neq 0$ then $X_1 + X_2 = x_1 \wedge (x_2 \wedge x_3 + x_4 \wedge x_5)$ and for X to have rank 8 it follows that $[X_3] \cap ([X_1] + [X_2]) = 0$. Therefore X has the form $x_1 \wedge (x_2 \wedge x_3 + x_4 \wedge x_5) + x_6 \wedge x_7 \wedge x_8$ or XVI in the Gurevich notation. If $[X_i] \cap [X_j] = 0$ for $i \neq j$ then $X_1 + X_2 = x_1 \wedge x_2 \wedge x_3 + x_4 \wedge x_5 \wedge x_6$ and $X_3 = u \wedge x_7 \wedge x_8$ where $u \in \langle x_1, \dots, x_6 \rangle$ and x_1, \dots, x_8 are independent. By refactoring X_1 and X_2 we may assume that $u = x_1 + x_4$ and so X is of type XIX which is equal to $([aqp] + 1/2[(b + c)(r + s)(p + t)] + 1/2[(b - c)(r - s)(p - t)])$. This takes care of trivectors of rank 8 and length 3. For each of the remaining equivalence classes except XV we exhibit a length 4 representation.

Types XI, XIII, and XVII are already in the form advertised.

$$\text{XII} = [(a - s)qp] + [(q - c)(p + r)s] + [(t + s)cr] + [brp].$$

$$\text{XIV} = [ab(c - p)] + [(a - r)(b + q)p] + [rq(p - s)] + [crt].$$

$$\text{XVIII} = [(t - r)bs] + [crt] + [r(p - s)(b - q)] + [(a - r)qp].$$

$$\text{XX} = [(r + s)(t - r)b] + [(r + s)(r + p)(c - q)] + [(a - r - s)qp] \\ + [r(b - c)(s - p + t)].$$

$$\text{XXI} = [(b - r)(c + s)t] + [(a - t)bc] + [aqp] + [rs(q + t)].$$

$$\text{XXII} = [(a + r)(b + 2q)(p - c + 1/2s)] + [cr(t - 3b - 2q)] \\ + [bs(1/2a + 3/2r + t)] + [(b + q)(a + 2r)(p - 2c + s)].$$

$$\text{XXIII} = [(a + 1/2s + 1/2r)q(b + c + r - s + 1/2p + 1/2t)] \\ + [(b + c + 1/2q)(r + s)(b + c + 1/2p + 1/2t)] \\ + [a(-1/2b + 1/2c + q)(-b - c - r + s + 1/2p - 1/2t)] \\ + [(b - c)(-1/2a + r - s)(-r + s + 1/2p - 1/2t)].$$

The only item that remains to be justified is that XV has length 5. We write the representative in the form $X = X_1 + [crt]$ where $X_1 = [abc] + [qrs] + [aqp] + [brp] + [csp]$. We note that X_1 is of type X, has rank 7, length 4, from which it follows that X has length at most 5. We will show that $(t - u) \wedge X$ has length at least 4 for all $u \in \langle a, b, c, q, r, s, p \rangle$. This will complete the proof because if Y is any rank 8 trivector in $\bigwedge^3 U$ with length 4 then at least one of the terms in any representation of Y as a sum of 4 decomposable trivectors must contain a factor of the form $t - u$ for some $u \in \langle a, b, c, q, r, s, p \rangle$, and for this u , the length of $(t - u) \wedge Y$ is at most 3.

Since $(t - u) \wedge X = t \wedge (X_1 + [ucr]) - u \wedge X_1$ it is sufficient to prove that $X_1 + [ucr]$ has length at least 4 for all $u \in \langle a, b, c, q, r, s, p \rangle$. Let

$$u = \alpha_1 a + \alpha_2 b + \alpha_3 c + \alpha_4 q + \alpha_5 s + \alpha_6 r + \alpha_7 p.$$

After the substitution

$$s \longrightarrow s - \alpha_7 r + \alpha_4 c$$

$$b \longrightarrow b + \alpha_1 r$$

with the other letters remaining unchanged we obtain

$$X_1 + [cr(\alpha_2 b + \alpha_5 s)].$$

If $\alpha_2 = \alpha_5 = 0$ we have X_1 which has length 4. If $\alpha_2 \neq 0$ then $X_1 + [cr(\alpha_2 b + \alpha_5 s)] = [(a + \alpha_2 r)b(c - \alpha_2^{-1}p)] + [(q + \alpha_5 c)rs] + [a(q + \alpha_2^{-1}b)p] + [csp] \sim X_1$ under the substitution

$$a \longrightarrow \alpha_2^{-1/2}(a - r)$$

$$b \longrightarrow \alpha_2^{3/2}b + \alpha_5(c - p)$$

$$c \longrightarrow \alpha_2^{-1}c$$

$$q \longrightarrow \alpha_2^{1/2}q - \alpha_2^{-1}\alpha_5 c$$

$$s \longrightarrow \alpha_2(s - p)$$

$$r \longrightarrow \alpha_2^{-3/2}r$$

$$p \longrightarrow p.$$

If $\alpha_2 = 0$ and $\alpha_5 \neq 0$ then

$$\begin{aligned} X_1 + \alpha_2[crs] &= [abc] + [brp] + [s(q + \alpha_5 c)(r - \alpha_5^{-1}p)] \\ &\quad + [(a + \alpha_5^{-1}s)qp] \sim X_1 \end{aligned}$$

under the substitution

$$a \longrightarrow -\alpha_5 a$$

$$b \longrightarrow -\alpha_5(b + p)$$

$$c \longrightarrow \alpha_5^{-2}c$$

$$q \longrightarrow -\alpha_5^{-1}(q + c)$$

$$r \longrightarrow -\alpha_5^{-1}r$$

$$s \longrightarrow \alpha_5^2 s$$

$$p \longrightarrow p.$$

Lastly, we point out also that $N(C, 9, 3) \leq 9$. This follows from (2.5) of [1] since $N(C, 8, 3) = 5$. The bound 9 is not likely to be the best one however.

REFERENCES

1. H. Busemann and D. E. Glassco, *Irreducible Sums of Simple Multivectors*, Pacific J. Math., **49** (1973), 13-32.
2. G. B. Gurevich, *Foundations of the Theory of Algebraic Invariants*, P. Noordhoff Ltd., Groningen the Netherlands, (1964).
3. ———, *Sur Les Trivecteurs Dans L'Espace a Sept Dimensions*, C. R. Acad. Sc. URSS., **3** (1934), 564-569.
4. ———, *Classification Des Trivecteurs Ayant Le Rang Huit*, C. R. Acad. Sc. USSR., **2** (1935), 353-356.
5. ———, *L'Algebre du Trivecteur*, Abh. Sem. Vektor u. Lensor. Moscow II, (1935), 51-113.
6. J. A. Schouten, *Tensor Analysis for Physicists*, 2nd Edition, Oxford, (1954).
7. ———, *Klassifizierung Der Alternierenden Groszen Dritten Grades in 7 Dimensionen*, Circolo. Mat. di. Palermo. Rend., **55** (1931), 137-156.
8. R. Westwick, *Trivectors in a space of Seven Dimensions*, Canad. Math. Bull., **20** (1977), 401.

Received March 9, 1977 and in revised form April 6, 1978.

UNIVERSITY OF BRITISH COLUMBIA
VANCOUVER, B. C.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, CA 90024

CHARLES W. CURTIS

University of Oregon
Eugene, OR 97403

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1978 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

K. Adachi, <i>On the multiplicative Cousin problems for $N^P(D)$</i>	297
Howard Banilower, <i>Isomorphisms and simultaneous extensions in $C(S)$</i>	305
B. R. Bhonsle and R. A. Prabhu, <i>An inversion formula for a distributional finite-Hankel-Laplace transformation</i>	313
Douglas S. Bridges, <i>Connectivity properties of metric spaces</i>	325
John Patton Burgess, <i>A selection theorem for group actions</i>	333
Carl Claudius Cowen, <i>Commutants and the operator equations $AX = \lambda XA$</i>	337
Thomas Curtis Craven, <i>Characterizing reduced Witt rings. II</i>	341
J. Csima, <i>Embedding partial idempotent d-ary quasigroups</i>	351
Sheldon Davis, <i>A cushioning-type weak covering property</i>	359
Micheal Neal Dyer, <i>Nonminimal roots in homotopy trees</i>	371
John Erik Fornæss, <i>Plurisubharmonic defining functions</i>	381
John Fuelberth and James J. Kuzmanovich, <i>On the structure of finitely generated splitting rings</i>	389
Irving Leonard Glicksberg, <i>Boundary continuity of some holomorphic functions</i>	425
Frank Harary and Robert William Robinson, <i>Generalized Ramsey theory. IX. Isomorphic factorizations. IV. Isomorphic Ramsey numbers</i>	435
Frank Harary and Allen John Carl Schwenk, <i>The spectral approach to determining the number of walks in a graph</i>	443
David Kent Harrison, <i>Double coset and orbit spaces</i>	451
Shiro Ishikawa, <i>Common fixed points and iteration of commuting nonexpansive mappings</i>	493
Philip G. Laird, <i>On characterizations of exponential polynomials</i>	503
Y. C. Lee, <i>A Witt's theorem for unimodular lattices</i>	509
Teck Cheong Lim, <i>On common fixed point sets of commutative mappings</i>	517
R. S. Pathak, <i>On the Meijer transform of generalized functions</i>	523
T. S. Ravisankar and U. S. Shukla, <i>Structure of Γ-rings</i>	537
Olaf von Grudzinski, <i>Examples of solvable and nonsolvable convolution equations in \mathcal{K}'_p, $p \geq 1$</i>	561
Roy Westwick, <i>Irreducible lengths of trivectors of rank seven and eight</i>	575