

Pacific Journal of Mathematics

BOREL BOXES

DOROTHY MAHARAM AND A. H. STONE

BOREL BOXES

DOROTHY MAHARAM AND A. H. STONE

This note gives a simple necessary and sufficient condition for a box, in a product of uncountably many metrizable spaces, to be a Borel set.

Throughout, $X = \prod \{X_a : a \in A\}$ where A is an uncountable index set and each X_a is a nonempty topological space; X is given the usual product topology. The a th co-ordinate of a point x of X is denoted by $x_a (a \in A)$. A sub-product $\prod \{X_a : a \in B\}$, where $B \subset A$, is denoted by X_B . A set of the form $Y = \prod \{Y_a : a \in A\}$, where $Y_a \subset X_a$ for each $a \in A$, will be called a "box" in X . The object of this note is to point out that, when each X_a is metrizable, there is a surprisingly simple characterization of the boxes that are Borel subsets of X . We require the following three lemmas, of which at least the first two are well known. (See for instance [2].) The third is a special case of our main result.

LEMMA 1. *Suppose each X_a is a separable metrizable space. Then each Baire set E in X is of the form $E_0 \times X_{A-B}$, where B is a countable subset of A and E_0 is a Borel subset of the countable product X_B . Conversely, each set of this form is a Baire subset of X .*

LEMMA 2. *Again suppose each X_a is separable and metrizable. Then, given a Borel set D in X , there exist Baire sets V, W in X such that $V \subset D \subset W$ and $W - V$ is meagre (in X).*

LEMMA 3. *Suppose each X_a is compact and metrizable, and p_a is a nonisolated point of $X_a (a \in A)$. Define*

$$D = \{x \in X : \text{for all } a \in A, x_a \neq p_a\} = \prod \{X_a - \{p_a\} : a \in A\}.$$

Then D is not Borel in X .

Proofs. When E is a cozero set, the assertion of Lemma 1 holds by [3], and the general case follows. The converse is elementary.

When D is open, the assertion of Lemma 2 holds; it suffices to take $V =$ union of a maximal collection (necessarily countable) of pairwise disjoint nonempty elementary open sets contained in D (as in [3]), and $W = \bar{V}$. Since the family of sets D for which Lemma 2 holds is a Borel field, it includes all Borel sets.

In Lemma 3, suppose D is Borel, and apply Lemma 2 to obtain Baire sets V, W with $V \subset D \subset W$ and $W - V$ meagre. By Lemma 1, $V = V_0 \times X_{A-B}$ where B is a countable subset of A . If $V \neq \emptyset$, pick $q \in V$ and define $x \in X$ by:

$$x_a = q_a \text{ if } a \in B; x_a = p_a \text{ if } a \in A - B.$$

Since A is uncountable, $x \notin D$. But $x \in V_0 \times X_{A-B} = V \subset D$, a contradiction. This shows $V = \emptyset$, and therefore W is meagre.

On the other hand, $D \subset W = W_0 \times X_{A-C}$ for some countable $C \subset A$. We have $\prod \{X_a - \{p_a\}; a \in C\} \subset W_0$, so that $X_C - W_0 \subset \bigcup \{\{p_a\} \times X_{C-\{a\}}; a \in C\}$, a countable union of closed subsets of X_C that are nowhere dense in X_C (because p_a is not isolated in X_a). Thus $X_C - W_0$ is meagre in X_C , whence $X - W$ is meagre in X . Since W also is meagre, X is meagre, contradicting Baire's theorem.

THEOREM. *Let $Y = \prod_a Y_a$ be a nonempty box in $X = \prod_a X_a$, and suppose each X_a is a first countable Hausdorff space. Then Y is a Borel set in X if, and only if,*

- (i) *for all $a \in A$, Y_a is Borel in X_a ,*
- (ii) *for all but at most a countable set of a 's, Y_a is closed in X_a .*

Proof. Put $B = \{a \in A: Y_a \text{ is not closed in } X_a\}$. Assuming (i) and (ii), we have that B is countable, and $Y = E \times \prod \{Y_a: a \in B\}$ where $E = \prod \{Y_a: a \in A - B\}$ is closed in X_{A-B} and each Y_a is Borel in X_a . As a countable product of Borel sets, Y is Borel.

Conversely, suppose Y is Borel in X . It is easy to see that (i) holds, and that the sub-product $Y_B = \prod \{Y_a: a \in B\}$ is Borel in X_B . Suppose (ii) is false, so that B is uncountable. For each $b \in B$, pick $p_b \in \bar{Y}_b - Y_b$ and pick a sequence of points q_{b1}, q_{b2}, \dots , of Y_b , distinct from p_b and from each other, converging to p_b . Put $Z_b = \{q_{bn}: n = 1, 2, \dots\}$, $T_b = Z_b \cup \{p_b\}$. Thus T_b is a compact metric space (homeomorphic to the subspace $\{0, 1, 1/2, \dots, 1/n, \dots\}$ of the real line), and p_b is a nonisolated point of it. Write $Z_B = \prod \{Z_b: b \in B\}$, $T_B = \prod \{T_b: b \in B\}$; then $Z_B = Y_B \cap T_B$, where T_B is closed and Y_B is Borel in X_B . Thus Z_B is Borel in X_B . But $Z_B = \{t \in T_B: \text{for all } b \in B, t_b \neq p_b\}$, so that Lemma 1 is contradicted. Thus (ii) holds, and the proof is complete.

COROLLARY 1. *If Y is a box in $X = \prod_a X_a$, where each X_a is a separable metric space, then Y is Borel in X if, and only if, Y is a Baire set relative to its closure \bar{Y} in X .*

(From the theorem and Lemma 1.)

COROLLARY 2. *Suppose, for each $a \in A$, X_a and X'_a are non-empty separable metric spaces and f_a is a Borel isomorphism of X_a onto X'_a . Then $\prod_a f_a$ is a Borel isomorphism of X onto $X' (= \prod_a X'_a)$ if, and only if, f_a is a homeomorphism for all but (at most) countably many a 's.*

The "only if" in Corollary 2 is a straightforward deduction from the theorem. The "if" is an easy consequence of the theorem and the following lemma.

LEMMA 4. *If (for $i = 1, 2$) g_i is a Borel measurable map from Z_i to Z'_i , and if at least one of the spaces Z'_1, Z'_2 has a countable base, then $g_1 \times g_2$ is Borel measurable.*

Proof. Let V_1, V_2, \dots , be a countable base for (say) Z'_2 . It is easy to see that an open set G in $Z'_1 \times Z'_2$ can always be written in the form $G = \bigcup \{(U_n \times V_n) : n = 1, 2, \dots\}$, where U_1, U_2, \dots , are suitable open sets in Z'_1 . From this it is clear that $(g_1 \times g_2)^{-1}(G)$ is a Borel set in $Z_1 \times Z_2$.

It would be interesting to know whether the separability hypothesis can be dropped in Corollary 2. It is not needed for the "only if" part of the corollary, and it is at least relatively consistent that separability is unnecessary for the "if" part. This is because the appropriate variant of Lemma 4 (with Z_2, Z'_2 both metrizable, though not necessarily separable) holds in a model of set theory given by Fleissner [1].

REFERENCES

1. W. G. Fleissner, *All Borel additive decompositions can be σ -discretely decomposable*, to appear.
2. Dorothy Maharam, *Realizing automorphisms of category algebras of product spaces*, *General Topology and Appl.*, **10** (1979), 161-174.
3. K. A. Ross and A. H. Stone, *Products of separable spaces*, *Amer. Math. Monthly*, **71** (1964), 398-403.

Received September 27, 1978.

UNIVERSITY OF ROCHESTER
ROCHESTER, NY 14627

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).

8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1979 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Ersan Akyildiz, <i>Vector fields and equivariant bundles</i>	283
Ehrhard Behrends, <i>The centralizer of tensor products of Banach spaces (a function space representation)</i>	291
Geoffrey R. Burton, <i>Congruent sections of a convex body</i>	303
John Warnock Carlson, <i>H-closed and countably compact extensions</i>	317
Robert Charles Carlson, <i>Eigenfunction expansions for selfadjoint integro-differential operators</i>	327
Robert Damiano, <i>Coflat rings and modules</i>	349
Eric Karel van Douwen and Washek (Vaclav) Frantisek Pfeffer, <i>Some properties of the Sorgenfrey line and related spaces</i>	371
Uri Elias, <i>Necessary conditions and sufficient conditions for difocality and disconjugacy of a differential equation</i>	379
V. L. (Vagn Lundsgaard) Hansen, <i>Polynomial covering spaces and homomorphisms into the braid groups</i>	399
Paul Hess, <i>Dedekind's problem: monotone Boolean functions on the lattice of divisors of an integer</i>	411
Alan Hopenwasser and David Royal Larson, <i>The carrier space of a reflexive operator algebra</i>	417
Kyung Bai Lee, <i>Spaces in which compacta are uniformly regular G_δ</i>	435
Claude Levesque, <i>A class of fundamental units and some classes of Jacobi-Perron algorithms in pure cubic fields</i>	447
Teck Cheong Lim, <i>A constructive proof of the infinite version of the Belluce-Kirk theorem</i>	467
Dorothy Maharam and A. H. Stone, <i>Borel boxes</i>	471
Roger McCann, <i>Asymptotically stable dynamical systems are linear</i>	475
Peter A. McCoy, <i>Approximation and harmonic continuation of axially symmetric potentials in E^3</i>	481
Takahiko Nakazi, <i>Extended weak-*Dirichlet algebras</i>	493
Carl L. Prather, <i>On the zeros of derivatives of balanced trigonometric polynomials</i>	515
Iain Raeburn, <i>An implicit function theorem in Banach spaces</i>	525
Louis Jackson Ratliff, Jr., <i>Two theorems on the prime divisors of zeros in completions of local domains</i>	537
Gloria Jean Tashjian, <i>Cartesian-closed coreflective subcategories of Tychonoff spaces</i>	547
Stephen Edwin Wilson, <i>Operators over regular maps</i>	559