

Pacific Journal of Mathematics

ON A THEOREM OF MURASUGI

CAMERON GORDON AND RICHARD A. LITHERLAND

ON A THEOREM OF MURASUGI

C. MCA. GORDON AND R. A. LITHERLAND

1. Let $l = k_1 \cup k_2$ be a 2-component link in S^3 , with k_2 unknotted. The 2-fold cover of S^3 branched over k_2 is again S^3 ; let $k_1^{(2)}$ be the inverse image of k_1 , and suppose that $k_1^{(2)}$ is connected. How are the signatures $\sigma(k_1)$, $\sigma(k_1^{(2)})$ of the knots k_1 and $k_1^{(2)}$ related? This question was considered (from a slightly different point of view) by Murasugi, who gave the following answer [Topology, 9 (1970), 283-298].

THEOREM 1 (*Murasugi*).

$$\sigma(k_1^{(2)}) = \sigma(k_1) + \xi(l).$$

Recall [4] that the invariant $\xi(l)$ is defined by first orienting l , giving, an oriented link \bar{l} , say, and then setting $\xi(l) = \sigma(\bar{l}) + Lk(\bar{k}_1, \bar{k}_2)$, where σ denotes signature and Lk linking number.

In the present note we shall give an alternative, more conceptual, proof of Theorem 1, and in fact obtain it as a special case of a considerably more general result.

The idea of our proof is the following. If $l = l_1 \cup l_2$ is a link, partitioned into two sublinks l_1 and l_2 , then the 2-fold branched covers over l_1 , l_2 , and the whole of l , are all quotients of a $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ -cover branched over l . After possibly multiplying by 2, the diagram consisting of these branched covers bounds a corresponding diagram of 4-manifolds, and the signatures of the various links involved are expressible in terms of the signatures of these 4-manifolds (and the euler numbers of the branch sets); see e.g., [3]. The result is then a consequence of a relation among these 4-manifold signatures (Lemma 1).

This more general setting requires that we consider links in 3-manifolds other than homology spheres; in § 2 we discuss the signature in this context. (It becomes necessary to prescribe a particular 2-fold branched cover. However, we sacrifice some generality inasmuch as we restrict ourselves to oriented, null-homologous links: it would otherwise be necessary to prescribe a framing of the link as well.) In § 3 we set up the diagram of covering spaces, and in § 4 derive the relation between the signatures of the manifolds therein. Section 5 contains some consequences of this, including the appropriate generalization of Theorem 1.

All manifolds of dimensions 3 and 4 are to be oriented; manifolds of dimensions 1 and 2 are oriented only when this is explicitly

stated, and those of dimension 2 need not even be orientable. We make no assumptions on the connectedness of our manifolds. If \bar{l} is an oriented link, we denote the underlying nonoriented link by l .

2. Let $\bar{l} = \bar{k}_1 \cup \dots \cup \bar{k}_m$ be an oriented link in a closed 3-manifold M , and suppose \bar{l} is null-homologous. Let W be a 4-manifold and F a surface in W such that $\partial(W, F) = (M, l)$. Let F' be (the image of) a section of the normal S^1 -bundle of F in W , with $\partial F' = l' = k'_1 \cup \dots \cup k'_m$, say. Orient l' to obtain $\bar{l}' = \bar{k}'_1 \cup \dots \cup \bar{k}'_m$ by requiring $\bar{k}'_i \sim \bar{k}_i$ in a tubular neighborhood of k_i , and define $\bar{\epsilon}(F) = -Lk(\bar{l}, \bar{l}')$. (Note that this is well-defined as \bar{l}, \bar{l}' are both null-homologous in M .)

Now let $p: \tilde{M} \rightarrow M$ be some 2-fold covering of M branched along l , and suppose that p extends to a 2-fold covering $\tilde{W} \rightarrow W$ branched along F . Then


$$\sigma(\bar{l}, p) = \sigma(\tilde{W}) - 2\sigma(W) + \frac{1}{2} \bar{\epsilon}(F)$$

depends only on \bar{l} and p . (If (W_1, F_1) and (W_2, F_2) are two pairs as above, apply the G -signature theorem [1] to the resulting involution on the closed 4-manifold $\tilde{W}_1 \cup \tilde{W}_2$, together with Novikov additivity and the fact that the euler number of the normal bundle of $F_1 \cup F_2$ in $W_1 \cup W_2$ is equal to $\bar{\epsilon}(F_1) - \bar{\epsilon}(F_2)$.)

We remark that if M is a homology sphere, p is unique, and $\sigma(\bar{l}, p)$ is just the signature of \bar{l} . Again, we may take l to be the empty link; $-\sigma(\phi, p)$ is the α -invariant [2] of the nontrivial covering translation of \tilde{M} .

3. Let l_1, l_2 be disjoint links in a 3-manifold M , and write $l = l_1 \cup l_2$. Let $\alpha: H_1(M - l) \rightarrow \mathbf{Z}_2 \oplus \mathbf{Z}_2$ be a homomorphism which sends each meridian of l_1 (resp. l_2) to the nontrivial element of the first (resp. second) \mathbf{Z}_2 . Let W be a 4-manifold and F_1, F_2 disjoint surfaces in W such that $\partial(W, F_1, F_2) = (M, l_1, l_2)$. Write $F = F_1 \cup F_2$, and suppose there exists a homomorphism $\beta: H_1(W - F) \rightarrow \mathbf{Z}_2 \oplus \mathbf{Z}_2$ such that $\alpha = \beta i_*$, where $i: M - l \rightarrow W - F$ is inclusion. (We shall discuss this assumption later.)

Let $\tilde{W} \rightarrow W$ be the branched covering associated with β . The covering translations induce a $\mathbf{Z}_2 \oplus \mathbf{Z}_2$ -action on \tilde{W} . Let g_1 generate the second \mathbf{Z}_2 factor, g_2 the first, and let $g_3 = g_1 g_2$ be the remaining nontrivial element. Setting $W^{(i)} = \tilde{W}/(g_i)$, $i = 1, 2, 3$, we have the following commutative diagram of 2-fold branched coverings.



Here q_i is branched over F_i , $i = 1, 2$, and q is branched over F . If $F_1^{(2)}$ and $F_2^{(1)}$ are the inverse images of F_1 in $W^{(2)}$ and F_2 in $W^{(1)}$, then $q_1^{(2)}$, $q_2^{(1)}$ are branched over $F_1^{(2)}$, $F_2^{(1)}$ respectively. Finally, \tilde{q} is unbranched.

Now suppose that l_1 and l_2 can be oriented to obtain null-homologous links \bar{l}_1 and \bar{l}_2 respectively. Let $\bar{l} = \bar{l}_1 \cup \bar{l}_2$. There are induced orientations of $l_1^{(2)} = \partial F_1^{(2)}$ and $l_2^{(1)} = \partial F_2^{(1)}$, giving null-homologous links $\bar{l}_1^{(2)}$ and $\bar{l}_2^{(1)}$ in $\partial W^{(2)}$ and $\partial W^{(1)}$ respectively.

Writing p 's instead of q 's to denote the restrictions of these coverings to the appropriate boundaries, we have the equations

- (i) $\sigma(\bar{l}_1, p_1) = \sigma(W^{(1)}) - 2\sigma(W) + \frac{1}{2} \bar{e}(F_1)$
- (ii) $\sigma(\bar{l}_2, p_2) = \sigma(W^{(2)}) - 2\sigma(W) + \frac{1}{2} \bar{e}(F_2)$
- (iii) $\sigma(\bar{l}, p) = \sigma(W^{(3)}) - 2\sigma(W) + \frac{1}{2} \bar{e}(F)$
- (iv) $\sigma(\bar{l}_1^{(2)}, p_1^{(2)}) = \sigma(\tilde{W}) - 2\sigma(W^{(2)}) + \frac{1}{2} \bar{e}(F_1^{(2)})$
- (v) $\sigma(\bar{l}_2^{(1)}, p_2^{(1)}) = \sigma(\tilde{W}) - 2\sigma(W^{(1)}) + \frac{1}{2} \bar{e}(F_2^{(1)})$
- (vi) $\sigma(\bar{\phi}, \bar{p}) = \sigma(\tilde{W}) - 2\sigma(W^{(3)}) .$

We now consider the question of the existence of a suitable homomorphism β . Suppose $H_1(W; \mathbf{Z}_2) = 0$. Then (see [3, § 1]) the cohomology exact sequence of the pair $(W, W - F)$, together with duality, gives an exact sequence

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H^1(W - F; \mathbf{Z}_2) & \longrightarrow & H_2(F, \partial F; \mathbf{Z}_2) & \longrightarrow & H_2(W, \partial W; \mathbf{Z}_2) \\
 & & & & \Downarrow & & \\
 & & & & H^0(F; \mathbf{Z}_2) & & .
 \end{array}$$

The existence of $\beta: H_1(W - F) \rightarrow \mathbf{Z}_2 \oplus \mathbf{Z}_2$ taking a meridian of F_i to the nontrivial element of the i th \mathbf{Z}_2 , $i = 1, 2$, is then seen to be equivalent to the condition that $[F_i, \partial F_i] = 0 \in H_2(W, \partial W; \mathbf{Z}_2)$, for $i = 1, 2$. (In particular, the assertion $H^1(W - F; \mathbf{Z}_2) \cong H^0(F; \mathbf{Z}_2)$ in [3, p. 353] is incorrect.)

Now suppose, in addition, that $H_1(M; \mathbf{Z}_2) = 0$. Then β will automatically satisfy $\alpha = \beta i_*$. But β will not in general exist, for it is clear that if $[F_i, \partial F_i] = 0 \in H_2(W, \partial W; \mathbf{Z}_2)$, $i = 1, 2$, then $Lk_{\mathbf{Z}_2}(l_1, l_2) = 0$. However, this condition is also sufficient; that is, given links $l_1, l_2 \subset M$ such that $H_1(M; \mathbf{Z}_2) = 0$ and $Lk_{\mathbf{Z}_2}(l_1, l_2) = 0$, there exist W, F_1, F_2, β as above. To see this, let W be any 4-manifold with $\partial W = M$ and $H_1(W; \mathbf{Z}_2) = 0$, and let E_1, E_2 be connected surfaces in W with $\partial E_i = l_i$ and $[E_i, \partial E_i] = 0 \in H_2(W, \partial W; \mathbf{Z}_2)$, $i = 1, 2$. (For example, we could obtain E_i by starting with a connected surface in M bounded by l_i and pushing its interior slightly into W .) We may assume that E_1 and E_2 intersect transversally in points in $\text{int } W$. Since $Lk_{\mathbf{Z}_2}(l_1, l_2) = 0$, there will be an even number of such intersection points, and these may be removed, a pair at a time, by adding a tube to (say) E_1 along an arc in E_2 connecting the two points in question.

REMARK. Section 5 contains equations, derived from (i)—(vi) above, involving link signatures and linking numbers. Since both are additive under disjoint union, these equations will still be valid if we only assume $\partial(W, F_1, F_2) = k(M, l_1, l_2)$, the disjoint union of k copies of (M, l_1, l_2) , for some $k > 0$. Moreover, we have just seen that this weaker assumption is always satisfied (with $k = 2$) if M is a \mathbf{Z}_2 -homology sphere. For notational simplicity, however, we shall continue to take $k = 1$, without further comment.

4. To deduce relations between the link signatures on the left of equations (i)—(vi), we must find relations between the quantities on the right. The main ingredient is the following.

LEMMA 1.

$$\sigma(\tilde{W}) = \sum_{i=1}^3 \sigma(W^{(i)}) - 2\sigma(W) .$$

Proof. If G is a finite group and N is a G -manifold, then a standard transfer argument shows that

$$(*) \quad \sigma(N) = |G|\sigma(N/G) - \sum_{g \in G - \{1\}} \text{sign}(g, N) .$$

Applying this to the $\mathbf{Z}_2 \oplus \mathbf{Z}_2$ -manifold \tilde{W} , we have

$$(1) \quad \sigma(\tilde{W}) = 4\sigma(W) - \sum_{i=1}^3 \text{sign}(g_i, \tilde{W}) .$$

For $i = 1, 2, 3$, $W^{(i)} = \tilde{W}/(g_i)$ has an action of $(\mathbf{Z}_2 \oplus \mathbf{Z}_2)/(g_i) \cong \mathbf{Z}_2$, generated by h_i , say. Applying (*) again, we get

$$(2) \quad \sigma(W^{(i)}) = 2\sigma(W) - \text{sign}(h_i, W^{(i)}), \quad i = 1, 2, 3.$$

By the proof of the proposition on page 415 of [2]

$$\text{sign}(h_i, W^{(i)}) = \frac{1}{2} \sum_{j \neq i} \text{sign}(g_j, \tilde{W}).$$

Hence

$$(3) \quad \sum_{i=1}^3 \text{sign}(h_i, W^{(i)}) = \sum_{i=1}^3 \text{sign}(g_i, \tilde{W}).$$

The result now follows from equations (1), (2) and (3).

We also need

LEMMA 2.

$$\begin{aligned} \bar{e}(F_1^{(2)}) &= 2\bar{e}(F_1), & \bar{e}(F_2^{(1)}) &= 2\bar{e}(F_2), \\ \bar{e}(F) &= \bar{e}(F_1) + \bar{e}(F_2) - 2Lk(\bar{l}_1, \bar{l}_2). \end{aligned}$$

(Note that $Lk(\bar{l}_1, \bar{l}_2)$ is well-defined, since \bar{l}_1 and \bar{l}_2 are both null-homologous.)

Proof. To prove the first statement, let V_1 be an oriented surface in M with $\partial V_1 = -\bar{l}_1$. Let the inverse image of V_1 in $\partial W^{(2)}$ be $V_1^{(2)}$, a 2-fold branched cover (possibly disconnected) of V_1 . Let \bar{l}'_1 be the (oriented) boundary of a section of the normal 1-sphere bundle of F_1 ; its inverse image $\bar{l}'_1{}^{(2)}$ in $\partial W^{(2)}$ is the boundary of a corresponding section for $F_1^{(2)}$. Then

$$\bar{e}(F_1^{(2)}) = \bar{l}'_1{}^{(2)} \cdot V_1^{(2)} = 2\bar{l}'_1 \cdot V_1 = 2\bar{e}(F_1).$$

Similarly, $\bar{e}(F_2^{(1)}) = 2\bar{e}(F_2)$. Finally, we may assume that \bar{l}'_1 does not meet l_2 , and is homologous to \bar{l}_1 in $M - l_2$. Extending in the obvious way the notation already introduced, we then have

$$\begin{aligned} \bar{e}(F) &= (\bar{l}'_1 \cup \bar{l}'_2) \cdot (V_1 \cup V_2) \\ &= \bar{l}'_1 \cdot V_1 + \bar{l}'_2 \cdot V_2 + \bar{l}'_1 \cdot V_2 + \bar{l}'_2 \cdot V_1 \\ &= \bar{e}(F_1) + \bar{e}(F_2) - 2Lk(\bar{l}_1, \bar{l}_2). \end{aligned}$$

5. From equations (i)-(iv), together with Lemmas 1 and 2, one easily obtains

$$\sigma(\bar{l}_1, p_1) + \sigma(\bar{l}, p) + Lk(\bar{l}_1, \bar{l}_2) = \sigma(\bar{l}_2, p_2) + \sigma(\bar{l}_1^{(2)}, p_1^{(2)}).$$

Now suppose $M = S^3$ and l_2 is the unknot. Then $\partial W^{(2)}$ is also S^3 , and $\sigma(\bar{l}_2, p_2) = 0$, so the above equation becomes

$$\sigma(\bar{l}_1^{(2)}) = \sigma(\bar{l}_1) + \sigma(\bar{l}) + Lk(\bar{l}_1, \bar{l}_2) .$$

If, further, l_1 has only one component, then

$$\sigma(\bar{l}) + Lk(\bar{l}_1, \bar{l}_2) = \xi(l)$$

so we obtain

$$\sigma(\bar{l}_1^{(2)}) = \sigma(l_1) + \xi(l) .$$

Theorem 1 is the special case in which $l_1^{(2)}$ has only one component.

REMARK. Using equations (i), (ii), (iii) and (vi) we obtain instead the relation

$$\sigma(\phi, \tilde{p}) + \sigma(\bar{l}, p) + Lk(\bar{l}_1, \bar{l}_2) = \sigma(\bar{l}_1, p_1) + \sigma(\bar{l}_2, p_2) .$$

If $M = S^3$, this can be written as

$$\sigma(\phi, \tilde{p}) + \xi(l) = \xi(l_1) + \xi(l_2) .$$

REFERENCET

1. M. F. Atiyah and I. M. Singer, *The index of elliptic operators III*, Ann. of Math., (2), **87** (1968), 546-604.
2. F. Hirzebruch, *Free involutions on manifolds and some elementary number theory*, Symposia Mathematica (Istituto Nazionale di Alta Matematica, Roma) Vol. V. Academic Press (1971), 411-419.
3. L. Kauffman and L. Taylor, *Signature of links*, Trans. Amer. Math. Soc., **216** (1976), 351-365.
4. K. Murasugi, *On the signature of links*, Topology, **9** (1970), 283-298.

Received April 17, 1978.

THE UNIVERSITY OF TEXAS
AUSTIN, TX 78712

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Werner Bäni, <i>Subspaces of positive definite inner product spaces of countable dimension</i>	1
Marilyn Breen, <i>The dimension of the kernel of a planar set</i>	15
Kenneth Alfred Byrd, <i>Right self-injective rings whose essential right ideals are two-sided</i>	23
Patrick Cousot and Radhia Cousot, <i>Constructive versions of Tarski's fixed point theorems</i>	43
Ralph S. Freese, William A. Lampe and Walter Fuller Taylor, <i>Congruence lattices of algebras of fixed similarity type. I</i>	59
Cameron Gordon and Richard A. Litherland, <i>On a theorem of Murasugi</i>	69
Mauricio A. Gutiérrez, <i>Concordance and homotopy. I. Fundamental group</i>	75
Richard I. Hartley, <i>Metabelian representations of knot groups</i>	93
Ted Hurley, <i>Intersections of terms of polycentral series of free groups and free Lie algebras</i>	105
Roy Andrew Johnson, <i>Some relationships between measures</i>	117
Oldřich Kowalski, <i>On unitary automorphisms of solvable Lie algebras</i>	133
Kee Yuen Lam, <i>K O-equivalences and existence of nonsingular bilinear maps</i>	145
Ernest Paul Lane, <i>PM-normality and the insertion of a continuous function</i>	155
Robert A. Messer and Alden H. Wright, <i>Embedding open 3-manifolds in compact 3-manifolds</i>	163
Gerald Ira Myerson, <i>A combinatorial problem in finite fields. I</i>	179
James Nelson, Jr. and Mohan S. Putcha, <i>Word equations in a band of paths</i>	189
Baburao Govindrao Pachpatte and S. M. Singare, <i>Discrete generalized Gronwall inequalities in three independent variables</i>	197
William Lindall Paschke and Norberto Salinas, <i>C*-algebras associated with free products of groups</i>	211
Bruce Reznick, <i>Banach spaces with polynomial norms</i>	223
David Rusin, <i>What is the probability that two elements of a finite group commute?</i>	237
M. Shafii-Mousavi and Zbigniew Zielezny, <i>On hypoelliptic differential operators of constant strength</i>	249
Joseph Gail Stampfli, <i>On selfadjoint derivation ranges</i>	257
Robert Charles Thompson, <i>The case of equality in the matrix-valued triangle inequality</i>	279
Marie Angela Vitulli, <i>The obstruction of the formal moduli space in the negatively graded case</i>	281