

Pacific Journal of Mathematics

**A NOTE ON COMPACT OPERATORS WHICH ATTAIN THEIR
NORM**

JOHN M. BAKER

A NOTE ON COMPACT OPERATORS WHICH ATTAIN THEIR NORM

J. M. BAKER

For Banach spaces X having the unit cell of $X^{**}w^*$ -sequentially compact, the compact operators from X into a Banach space Y attain their norm in X^{**} . The same holds for weakly compact operators if, in addition, X has the strict Dunford-Pettis property. For Banach spaces X such that the quotient space X^{**}/X is separable and Y the space of absolutely summable sequences, a proper subset P_o of the finite rank operators from X into Y is exhibited. The set P_o is shown to consist of operators which attain their norm and to be norm-dense in the operator space.

Throughout, X and Y will be Banach spaces and $\mathcal{L}(X, Y)$ the space of bounded linear operators from X into Y . An operator $T \in \mathcal{L}(X, Y)$ attains its norm on the unit cell $S_{X^{**}}$ of X^{**} if $\|T\| = \|T^{**}x^{**}\|$ for some $x^{**} \in X^{**}$ of norm one. For general results on norm attaining operators and their density in $\mathcal{L}(X, Y)$, see [2]. A space X is said to have the strict Dunford-Pettis property [4 p. 137] if for all Banach spaces Y an arbitrary weakly compact operator $T \in \mathcal{L}(X, Y)$ maps weakly Cauchy sequences to strongly Cauchy sequences.

THEOREM 1. *Let X be a Banach space with $S_{X^{**}}$ sequentially compact in the $\sigma(X^{**}, X^*)$ topology. Then*

(i) *if $T \in \mathcal{L}(X, Y)$ is compact, T attains its norm on $S_{X^{**}}$. Thus, every compact operator with reflexive domain X attains its norm on S_X .*

(ii) *if $T \in \mathcal{L}(X, Y)$ is weakly compact and X has the strict Dunford-Pettis property, T attains its norm on $S_{X^{**}}$. In addition, therefore, if Y is reflexive, all operators attain their norms on $S_{X^{**}}$.*

Proof. There is a sequence $\{x_n\}$ in S_X satisfying $\|T\| < \|Tx_n\| + 1/n$. Let J_x be the canonical embedding of X into X^{**} . Since $\{J_x x_n\} \subseteq S_{X^{**}}$ there exists a subsequence $\{x_{n_j}\}$ and an $x^{**} \in S_{X^{**}}$ such that $J_x x_{n_j} \xrightarrow{j} x^{**}$ in the $\sigma(X^{**}, X^*)$ -topology. The sequence $\{x_{n_j}\}$ is weakly Cauchy in X , whence under either hypothesis there exists a subsequence $\{w_j\}$ of $\{x_{n_j}\}$ such that $\{Tw_j\}$ is norm-convergent to some $y \in Y$. Since $\{J_x w_j\}$ is $\sigma(X^{**}, X^*)$ -convergent to x^{**} and $\{Tw_j\}$ is weakly convergent to y , we have $T^{**}x^{**} = J_y y$. Thus,

$$\|Tw_j\| \rightarrow \|y\| = \|T^{**}x^{**}\|, \quad \text{whence } \|T\| = \|T^{**}x^{**}\|. \quad \Delta$$

A consequence of [9, Theorem 3] is that if S_X is an RNP set and $\delta > 0$ every $T \in \mathcal{L}(X, Y)$ may be written as $T = T_1 + T_2$, where T_2 attains its norm on S_X and T_1 is rank one (thus attaining its norm on $S_{X^{**}}$) with $\|T_1\| < \delta$. A similar weaker result comes from [5] (proof of Theorem 1 and Remark p. 142) and Theorem 1.

COROLLARY. *If X is a Banach space with $S_{X^{**}\sigma}(X^{**}, X^*)$ -sequentially compact and $\delta > 0$, every $T \in \mathcal{L}(X, Y)$ may be written as $T = T_1 + T_2$, where both attain their norm on $S_{X^{**}}$ and T_1 is compact with $\|T_1\| < \delta$.*

Let Y be a weakly sequentially complete space and $X = c$ the Banach space of convergent sequences. Every $T \in \mathcal{L}(X, Y)$ is compact [3, p. 515], and since X^* is separable, part (i) of the theorem gives that every operator attains its norm on $S_{X^{**}}$. Moreover, the same can occur under the hypotheses of part (ii). Such cases render the central result Theorem 1 of [5] trivial, making it desirable to find useful subsets of norm attaining operators which are dense in the operator space. Such is the purpose of the remainder of this note for the case of Banach spaces X having $X^{**}/J_x X$ separable and $Y = l_1$, the space of absolutely summable sequences. For such spaces X , $\mathcal{L}(X, l_1)$ consists entirely of compact operators [6, Theorem 5].

LEMMA. *If X is a Banach space for which $X^{**}/J_x X$ is separable, then $S_{X^{**}}$ is $\sigma(X^{**}, X^*)$ -sequentially compact.*

Proof. Let $\{x_n^{**}\} \subseteq S_{X^{**}}$. Since the $\sigma(X^{**}, X^*)$ -sequential closure of X is X^{**} [6], for each positive integer n there exists a sequence $\{x_{ni}\}_{i=1}^\infty$ in S_X such that $J_x x_{ni} \xrightarrow{i} x_n^{**}$ in the $\sigma(X^{**}, X^*)$ topology. Let Z be the closed linear span of the set $\{x_{ni}\}$ and apply the lemma in §1 of [6] to deduce that Z^{**} , whence Z^* , is separable. This gives $S_{Z^{**}}$ to be $\sigma(Z^{**}, Z^*)$ -sequentially compact. The remainder of the proof is straight forward using the Hahn-Banach theorem.

Let $X^* \otimes_\lambda Y$ denote the tensor product of X^* and Y equipped with the least crossnorm λ [8]. The assignment $(\sum f_i \otimes y_i)(x) = \sum f_i(x)y_i$ defines an isometric isomorphism of $X^* \otimes_\lambda Y$ onto the subspace of compact operators in $\mathcal{L}(X, Y)$ of finite rank. In the following we let $\{e_i\}$ be the usual unit vector basis of l_1 and put

$$P_\sigma(X, l_1) = \left\{ \sum_{i=1}^n f_i \otimes e_i : f_i \in X^* \right\}$$

where n is arbitrary and f_i attains its norm on S_X . $P_\sigma(X, l_1)$ is not equivalent to the tensor product of two norm dense subsets.

THEOREM 2. *If $X^{**}/J_\sigma X$ is separable, $P_\sigma(X, l_1)$ is norm dense in $\mathcal{L}(X, l_1)$.*

Proof. Let $X^{**}/J_\sigma X$ be separable, $\varepsilon > 0$ be given, and $T \in \mathcal{L}(X, l_1)$. Since every operator in $\mathcal{L}(X, l_1)$ is compact and l_1 has the approximation property [7, p. 115], we have $\mathcal{L}(X, l_1) = X^* \widehat{\otimes}_\lambda l_1$, where $\widehat{\otimes}$ denotes the closure in $\mathcal{L}(X, l_1)$ of $X^* \otimes_\lambda l_1$. Thus, there exists $T_1 \in X^* \otimes_\lambda l_1$ such that $\|T - T_1\| < \varepsilon/3$, where $T_1 = \sum_{s=1}^k x_s^* \otimes y_s$ for appropriate $x_s^* \in X^*$ and $y_s = (\xi_{1s}, \xi_{2s}, \dots) \in l_1$. Hence, $T_1 x = \sum_{s=1}^k x_s^*(x) y_s = ([\sum_{s=1}^k \xi_{1s} x_s^*](x), [\sum_{s=1}^k \xi_{2s} x_s^*](x), \dots)$.

The series $\sum_{j=1}^\infty \|\sum_{s=1}^k \xi_{js} x_s^*\|$ converges: if $\eta > 0$ and m, n are positive integers, $n > m$, there exists $N > 0$ such that $n, m > N$ imply $\sum_{j=m+1}^n |\xi_{js}| < \eta/k \|x_s^*\|$, for $s = 1, 2, \dots, k$, because for each $s \sum_{j=1}^\infty |\xi_{js}|$ converges. Thus, $\sum_{j=m+1}^n \|\sum_{s=1}^k \xi_{js} x_s^*\| \leq \sum_{s=1}^k \|x_s^*\| \sum_{j=m+1}^n |\xi_{js}| < \eta$.

For each $j = 1, 2, \dots$, there exists a norm attaining $f_j \in X^*$ such that $\|f_j - \sum_{s=1}^k \xi_{js} x_s^*\| < \varepsilon/3^{j+1}[1]$, whence the series $\sum_{j=1}^\infty \|f_j\|$ converges since $\|f_j\| < \varepsilon/3^{j+1} + \|\sum_{s=1}^k \xi_{js} x_s^*\|$. We define $T_2 x = (f_1(x), f_2(x), \dots)$. Since T_2 is clearly a bounded linear operator from X into l_1 , we note $\|T_2 x - T_1 x\| \leq \sum_{j=1}^\infty \|f_j - \sum_{s=1}^k \xi_{js} x_s^*\| \|x\| < \varepsilon \|x\|/3$, whence $\|T_2 - T_1\| < \varepsilon/3$. Since there exists $N > 0$ such that $\sum_{j=N+1}^\infty \|f_j\| < \varepsilon/3$, we have $\|T - \sum_{j=1}^N f_j \otimes e_j\| \leq \|T - T_1\| + \|T_1 - T_2\| + \|T_2 - \sum_{j=1}^N f_j \otimes e_j\| < 2\varepsilon/3 + \sup_{S_X} \sum_{j=N+1}^\infty |f_j(x)| < \varepsilon$.

REFERENCES

1. E. Bishop and R. R. Phelps, *A proof that every Banach space is subreflexive*, Bull. Amer. Math. Soc., **67** (1961), 97-98.
2. J. Diestel and J. J. Uhl, Jr., *Vector measures*, Amer. Math. Soc. Survey, **15** (1977).
3. H. Dunford and J. T. Schwartz, *Linear Operators I*, Inter-Science Pub., N. Y., 1958.
4. A. Grothendieck, *Sur les applications lineaires faiblement compactes d'espaces du type $C(K)$* , Canad. J. Math., **5** (1953), 129-173.
5. J. Lindenstrauss, *On operators which attain their norm*, Israel J. Math., **1** (1963), 139-148.
6. R. D. McWilliams, *On certain Banach spaces which are w^* -sequentially dense in their second duals*, Duke J. Math., **37** (1970), 121-126.
7. H. H. Schaefer, *Topological vector spaces*, 2d. ed., McMillan, N. Y., 1967.
8. R. Shatten, *A theory of cross spaces*, Annals of Math. Studies, **26** (1950).
9. Charles Stegall, *Optimization of functions on certain subsets of Banach spaces*, Math. Annalen, **236** (1978), 171-176.

Received April 4, 1978.

WESTERN CAROLINA UNIVERSITY
CULLOWHEE, NC 28723

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$84.00 a year (6 Vols., 12 issues). Special rate: \$42.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).

8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1979 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Krishnaswami Alladi and Paul Erdős, <i>On the asymptotic behavior of large prime factors of integers</i>	295
Alfred David Andrew, <i>A remark on generalized Haar systems in L_p, $1 < p < \infty$</i>	317
John M. Baker, <i>A note on compact operators which attain their norm</i>	319
Jonathan Borwein, <i>Weak local supportability and applications to approximation</i>	323
Tae Ho Choe and Young Soo Park, <i>Wallman's type order compactification</i>	339
Susanne Dierolf and Ulrich Schwanengel, <i>Examples of locally compact noncompact minimal topological groups</i>	349
Michael Freedman, <i>A converse to (Milnor-Kervaire theorem) $\times R$ etc.</i>	357
George Golightly, <i>Graph-dense linear transformations</i>	371
H. Groemer, <i>Space coverings by translates of convex sets</i>	379
Rolf Wim Henrichs, <i>Weak Frobenius reciprocity and compactness conditions in topological groups</i>	387
Horst Herrlich and George Edison Strecker, <i>Semi-universal maps and universal initial completions</i>	407
Sigmund Nyrop Hudson, <i>On the topology and geometry of arcwise connected, finite-dimensional groups</i>	429
K. John and Václav E. Zizler, <i>On extension of rotund norms. II</i>	451
Russell Allan Johnson, <i>Existence of a strong lifting commuting with a compact group of transformations. II</i>	457
Bjarni Jónsson and Ivan Rival, <i>Lattice varieties covering the smallest nonmodular variety</i>	463
Grigori Abramovich Kolesnik, <i>On the order of Dirichlet L-functions</i>	479
Robert Allen Liebler and Jay Edward Yellen, <i>In search of nonsolvable groups of central type</i>	485
Wilfrido Martínez T. and Adalberto Garcia-Maynez Cervantes, <i>Unicoherent plane Peano sets are σ-unicoherent</i>	493
M. A. McKiernan, <i>General Pexider equations. I. Existence of injective solutions</i>	499
M. A. McKiernan, <i>General Pexider equations. II. An application of the theory of webs</i>	503
Jan K. Pachl, <i>Measures as functionals on uniformly continuous functions</i>	515
Lee Albert Rubel, <i>Convolution cut-down in some radical convolution algebras</i>	523
Peter John Slater and William Yslas Vélez, <i>Permutations of the positive integers with restrictions on the sequence of differences. II</i>	527
Raymond Earl Smithson, <i>A common fixed point theorem for nested spaces</i>	533
Indulata Sukla, <i>Generalization of a theorem of McFadden</i>	539
Jun-ichi Tanaka, <i>A certain class of total variation measures of analytic measures</i>	547
Kalathoor Varadarajan, <i>Modules with supplements</i>	559
Robert Francis Wheeler, <i>Topological measure theory for completely regular spaces and their projective covers</i>	565