

Pacific Journal of Mathematics

AN ANTI-OPEN MAPPING THEOREM FOR FRÉCHET SPACES

STEVEN F. BELLENOT

AN ANTI-OPEN MAPPING THEOREM FOR FRÉCHET SPACES

STEVEN F. BELLENOT

It is well-known that completeness is necessary for the usual open mapping theorem for Fréchet spaces. In contrast, it is shown that, with the obvious exception of ω , each infinite-dimensional Fréchet space has another distinct complete topology with the same continuous dual.

By a space or subspace, we mean an infinite-dimensional locally convex Hausdorff topological vector space over either the real or the complex scalars. Our notation generally follows Robertson and Robertson [7]. In particular, X' and $\sigma(X, X')$ denote the continuous dual and the weak topology on X , respectively. Denote by ω (respectively, ϕ) the space formed by the product (respectively, direct sum) of countably-many copies of the scalar field. We use c_0 , l_1 and l_∞ to denote the Banach sequence spaces (with their usual norms) of, respectively, null sequences, absolutely summable sequences and bounded sequences.

Our main result can be stated as:

THEOREM. *Each Fréchet space $(X, \zeta) \neq \omega$ has a topology η , so that, $\sigma(X, X') < \eta < \zeta$ and the space (X, η) is complete.*

By the open mapping theorem, (X, η) is a complete space which is not barrelled. In Section one we prove the theorem for the special cases of $(X, \zeta) = c_0$ (Case I) and (X, ζ) a nuclear space with a continuous norm (Case II). Then in Section two we reduce the theorem to these special cases.

We will have occasion to use Grothendieck's characterization of the completion of the space (X, ζ) as the set of linear functionals on X' which are $\sigma(X', X)$ -continuous on ζ -equicontinuous sets (see Robertson and Robertson [7], p. 103). Berezanskii's [4] (see also [2, pp. 61-62]) notion of inductive semi-reflexivity is used in Case II. In particular, complete nuclear spaces are inductive semi-reflexive, and the topology constructed from $\{\mu_n\}$ in Case II is complete in any inductive semi-reflexive space. The only other fact used about nuclear spaces is that their topology can be defined by means of (semi-) inner products (see Case II and Schaefer [7] p. 103).

Perhaps it is worth pointing out, that there are always lots of differently-defined complete topologies on each complete separable space (see Bellenot [1], [2] and with Ostling [3]): the difficulty is in

showing that these topologies are really different.

1. Two special cases. First we prove the theorem for the following special cases:

Case I. The Banach space c_0 : Let ξ be the norm topology on c_0 and let \mathcal{U} be a free ultrafilter on the set of positive integers (i.e., $\cap \mathcal{U} = \emptyset$). For each $A \in \mathcal{U}$ and $K > 0$ let

$$E(A, K) = \{x = (x_n) \in l_1: \|x\|_1 \leq K \text{ and } x_m = 0 \text{ for each } m \in A\}.$$

Let η be the topology of uniform convergence of the collection of sets

$$\{E(A, K) \cup \{y^n\}: A \in \mathcal{U}, K > 0, \{y^n\} \text{ a } l_1\text{-norm-null-sequence}\}.$$

Since finite sets are η -equicontinuous and each of the sets above are ξ -equicontinuous, we have $\sigma(c_0, l_1) \leq \eta \leq \xi$.

To see that $\eta < \xi$, note that if $\eta = \xi$ there would be a set $E(A, K) \cup \{y^n\}$ whose polar is contained in the unit ball of c_0 . Since y^n is a l_1 -norm-null-sequence, there is an M , so that $m \geq M$ implies that $|y_m^n| < 2^{-1}$, for each n . (Where y^n is the sequence $\{y_m^n\}_m$.) Since \mathcal{U} is free, A must be infinite and there is a $k \in A$ with $k \geq M$. Consider $x \in c_0$, the vector which is the zero sequence, except that it is 2 in the k th position. Clearly x is not in the unit ball of c_0 , but it is the polar of $E(A, K) \cup \{y^n\}$, a contradiction.

Consider X , the completion of (c_0, η) , as a subspace of the algebraic dual of l_1 . Since each l_1 -norm-null-sequence is η -equicontinuous, $X \subset l_\infty$. Suppose D is a subset of the positive integers with $D \notin \mathcal{U}$. Then, since \mathcal{U} is an ultrafilter, D^c , the complement of D , is an element of \mathcal{U} . Thus the η -topology restricted to the subspace $\{x \in c_0: x_n = 0 \text{ if } n \in D\}$ is the norm topology. It follows that for each $f = (f_n) \in X$, the subsequence $\{f_n: n \in D\}$ is a null-sequence, since f is $\sigma(l_1, c_0)$ -continuous on $E(D^c, 1)$. Let $f = (f_n) \in l_\infty$ with $A = \{n: |f_n| \geq \delta\}$ infinite, for some $\delta > 0$. If $A \notin \mathcal{U}$, then $f \notin X$ by the above, so assume $A \in \mathcal{U}$. Write $A = B \cup C$, a disjoint union of infinite sets, one of them is not in \mathcal{U} , and thus $f \notin X$. Therefore $X = c_0$ and (c_0, η) is complete.

Case II. (X, ξ) is a nuclear Fréchet space with a continuous norm: Let $\|\cdot\|_1 \leq \|\cdot\|_2 \leq \dots$ be a sequence of continuous norms which define the ξ -topology on X . Since X is nuclear, we assume that the unit ball of each $\|\cdot\|_{k+1}$ is precompact in the norm $\|\cdot\|_k$ and that each $\|x\|_k^2 = \langle x, x \rangle_k$, for some continuous inner product $\langle \cdot, \cdot \rangle_k$ on $X \oplus X$. Let $\|\cdot\|_k$ also represent (the possibly infinite-valued) dual norm of $\|\cdot\|_k$ on X' . A sequence $\{a'_n\} \subset X'$ is called k -admissible if

$\{\|a'_n\|_k\}$ is bounded and the semi-norm, $\rho_a(x) = \sup_n |a'_n(x)|$, defined for $x \in X$, is stronger than $\|\cdot\|_1$. That is, there is a constant K , with

$$(*) \quad \|x\|_1 \leq K\rho_a(x), \text{ for each } x \in X.$$

A nonincreasing null-sequence of positive reals $\{\lambda_n\}$ is said to be *k-discriminating*, if for each *k*-admissible sequence $\{a'_n\}$,

$$\limsup_n \|a'_n\|_k \lambda_n^{-1} = \infty.$$

Note that if $\{\lambda_n\}$ is *k-discriminating* and $\{\mu_n\}$ is another nonincreasing null-sequence of positive reals so that $\lim_n \mu_n/\lambda_n = 0$, then $\{\mu_n\}$ is also *k-discriminating*.

First, we prove that for each *k*, there is a *k-discriminating* sequence. To see this, let $\{e_n\} \subset X$ be a sequence orthogonal in $\langle \cdot, \cdot \rangle_k$ and orthonormal in $\langle \cdot, \cdot \rangle_1$. (The $\{e_n\}$ can be chosen inductively, by picking $e_{n+1} \in (\bigcap_1^n \ker f_i) \cap (\bigcap_1^n \ker g_i)$, where f_i and g_i are the continuous linear functionals given by $f_i(x) = \langle e_i, x \rangle_1$ and $g_i(x) = \langle e_i, x \rangle_k$, $i = 1, 2, \dots, n$.) Re-order $\{e_n\}$ so that the sequence $\{\|e_n\|_k\}$ is nondecreasing. We claim that the sequence $\lambda_n = 1/n \|e_n\|_k$ is *k-discriminating*. Suppose not, then there is *k*-admissible $\{a'_n\}$ with

$$(**) \quad \|a'_n\|_k \leq \lambda_n.$$

Let $\delta = 2^{-1}K^{-1}$, where K is the constant in (*).

Inductively choose $f_n \in X$ and an integer-valued function ϕ , so that

- (1) $\|f_n\|_1 = 1$ and $f_n \in \text{span}\{e_j : (n-1)^2 < j \leq n^2\}$;
- (2) $a'_{\phi(j)}(f_n) = 0$ for $j < n$; and
- (3) $|a'_{\phi(n)}(f_n)| \geq \delta$.

If f_j and $\phi(j)$ have been chosen for $j < n$, it is possible to choose f_n satisfying (1) and (2) since condition (2) puts $n-1$ constraints on f_n and f_n is chosen from a $(2n-1)$ -dimensional space. Thus by (*) we can find a $\phi(n)$ such that $2^{-1}\|f_n\|_1 = 2^{-1} \leq K|a'_{\phi(n)}(f_n)|$, i.e., that (3) is satisfied.

Let $A(n) = \{j : (n-1)^2 < j \leq n^2\}$ and suppose $f_n = \sum_{i \in A(n)} \alpha_i e_i$. Since $\{e_i\}$ is orthonormal in $\langle \cdot, \cdot \rangle_1$, condition (1) implies $\sum_{i \in A(n)} |\alpha_i|^2 = 1$. But $\{e_i\}$ is orthogonal in $\langle \cdot, \cdot \rangle_k$ hence

$$\|f_n\|_k = \left[\sum_{i \in A(n)} |\alpha_i|^2 \|e_i\|_k^2 \right]^{1/2} \leq \|e_{n^2}\|_k = n^{-1}\lambda_n^{-1}.$$

Thus by condition (3), we have

$$(***) \quad \delta \leq |a'_{\phi(n)}(f_n)| \leq \|a'_{\phi(n)}\|_k \cdot \|f_n\|_k \leq n^{-1}\lambda_n^{-1} \|a'_{\phi(n)}\|_k.$$

On the other hand, condition (2) implies that ϕ is 1-1 and hence $\phi(n) \geq n$, infinitely often. Thus (**) implies $\|a'_{\phi(n)}\|_k \leq \lambda_{\phi(n)} \leq \lambda_n$, infinitely often. Combining with (***) yields

$$0 < \delta \leq n^{-1} \lambda_n^{-1} \lambda_n = \frac{1}{n}, \text{ for infinitely many } n,$$

a contradiction.

For $k \geq 1$, let $\{\lambda_n^k\}$ be a k -discriminating. The sequence $\lambda_n = n^{-1} \min \{\lambda_n^j : j \leq n\}$ is thus k -discriminating for each $k \geq 1$. Let $\mu_n = \lambda_{n(n+1)}$, and let η be the topology of uniform convergence on sequences $\{a'_n\} \subset X'$ with the property that there is an integer k and constant K with $\|a'_n\|_k \leq K\mu_n$. It is easy to check that $\sigma(X, X') < \eta \leq \xi$. Note that if U is any ξ -neighborhood of the origin and if ρ_{U^0} is the gauge functional of U -polar in X' , then there is an integer k and a constant K so that $a' \in X'$ implies $\|a'\|_k \leq K\rho_{U^0}(a')$. Thus by Bellenot [2, p. 62 and Th. 4.1, p. 64], η is a ξ -rotor topology and (X, η) is complete.

To show that $\eta < \xi$, we will prove that $\|\cdot\|_1$ is not η -continuous on X . By Robertson and Robertson [7, p. 46], the η -neighborhoods of the origin are polars of finite unions of the above sequences (as sets of values in X'). (Note that it is possible for $\lim (\mu_n/\mu_{2n}) = \infty$, and so we must consider finite unions.) Suppose $\|\cdot\|_1$ is η -continuous, then there is a finite number of sequences $\{b'_{n,i}\}_n, 1 \leq i \leq j$, used to define η , so that $\|x\|_1 \leq \sup \{\|b'_{n,i}(x)\| : 1 \leq i \leq j, n = 1, 2, \dots\}$ for each $x \in X$. Let k and K be so that $\|b'_{n,i}\|_k \leq K\mu_n$, for $1 \leq i \leq j$ and each n . Let $\{a'_n\}$ be a listing of values in X' contained in the sequences $\{b'_{n,i}\}_n, 1 \leq i \leq j$, so that $\{\|a'_n\|_k\}$ is nonincreasing. It follows that $\{a'_n\}$ is k -admissible. Since $n > j \geq i \geq 1$ implies $(n+1)(n+2) \geq nj + i$, and $\{\lambda_n\}$ is nonincreasing, $\lambda_{(n+1)(n+2)} \leq \lambda_{nj+i}$. Thus if $m \geq j^2 + j + 1$, then $m = nj + i$ with $n > j \geq i \geq 1$, and

$$\|a'_m\|_k \leq K\mu_{n+1} = K\lambda_{(n+1)(n+2)} \leq K\lambda_m.$$

Hence $\limsup_m (\|a'_m\|_k/\lambda_m) < \infty$ and since $\{\lambda_m\}$ is k -discriminating, $\{a'_n\}$ is not k -admissible. This contradiction completes the proof of the theorem for this case.

2. The general case. The following two lemmas are of a general nature. The first lemma shows that completeness is a "three space property" while the second is used often in the proof of the theorem. The referee has pointed out that Lemma 1 is known, we include a proof for completeness.

LEMMA 1. Let X be a space, Y a closed subspace of X and $Z = X/Y$, the quotient. If Y and Z are complete, then X is complete.

Proof. Let $\phi: X \rightarrow Z$ be the quotient map and let $j: X \rightarrow \hat{X}$ be the injection of X into its completion \hat{X} . Since Z is complete, ϕ

extends to a map $\hat{\phi}: \hat{X} \rightarrow Z$ so that $\hat{\phi} \circ j = \phi$. Furthermore, since Y is complete, $j(Y)$ is closed in \hat{X} , and thus we can construct the quotient $W = \hat{X}/j(Y)$ with quotient map $\psi: \hat{X} \rightarrow W$. Since $\ker \psi \circ j = Y$, there is a map $\theta: Z \rightarrow W$ so that $\theta \circ \phi = \psi \circ j: X \rightarrow W$. Thus $\theta \circ \hat{\phi} \circ j = \psi \circ j$, but since $j(X)$ is dense in \hat{X} and since $\theta \circ \hat{\phi}$ and ψ are continuous, we have $\theta \circ \hat{\phi} = \psi$. Therefore θ and thus j are surjective maps, so that X is complete.

LEMMA 2. *A Fréchet space X satisfies the conclusions of the theorem if X has a closed subspace Y which satisfies the conclusions of the theorem.*

Proof. Let ξ be the topology on X with neighborhood basis of the origin \mathcal{U} . Let η be a topology on Y with $\sigma(Y, Y') \leq \eta < \xi|_Y$ and so that the space (Y, η) is complete. Let \mathcal{V} be the neighborhood basis of the origin for (Y, η) . Let $\mathcal{W} = \{V + U: V \in \mathcal{V}, U \in \mathcal{U}\}$. It is straightforward to check that \mathcal{W} is a neighborhood basis of the origin for a topology ζ on X with the properties:

- (i) $\sigma(X, X') \leq \zeta < \xi$,
- (ii) $\zeta|_Y = \eta$, and
- (iii) $(X, \zeta)/Y \equiv (X, \xi)/Y$.

Thus by (ii), (iii) and Lemma 1, (X, ζ) is complete and by (i) it satisfies the conclusion of the theorem.

Proof of the theorem. Let (X, ξ) be a Fréchet space $\neq \omega$. It follows that ξ is not the weak topology on X . First, we show there is a separable closed subspace Y of X , so that ξ , restricted to Y has a continuous norm. Since ξ is strictly stronger than $\sigma(X, X')$, there exists a continuous semi-norm on (X, ξ) which is not a linear combination of semi-norms $x \rightarrow |\langle x, x' \rangle|$ with $x' \in X'$, and thus from Schaefer [8], corollary on p. 124 it follows that X has a continuous semi-norm ρ so that the dimension of $X/\ker \rho$ is infinite. Let E be the normed space $X/\ker \rho$ with ρ norm and let $\psi: X \rightarrow E$ be the quotient map. Let $\{e_n\} \subset E$ be a linearly independent sequence. Let $\{x_n\} \subset X$ be so that $\psi(x_n) = e_n$, and let Y be the closed linear span of $\{x_n\}$ in (X, ξ) . Since $\rho(\sum_1^n \alpha_i x_i) = \rho(\sum_1^n \alpha_i e_i)$, for all scalar sequences $\{\alpha_i\}$, ψ , restricted to Y , is an isometry of Y with semi-norm ρ into a subspace of E with norm ρ . Thus by Lemma 2, we assume that (X, ξ) is separable and has a continuous norm.

Suppose (X, ξ) is a Banach space. In the notation of Bellenot and Ostling [3], since X is separable and complete, we have $\xi = \xi_M$. Furthermore, Theorem 3.1 of that same paper shows (X, ξ_{SW}) is complete, where ξ_{SW} is the topology of uniform convergence on ξ -equicontinuous $\sigma(X', X)$ -null sequences. Clearly, $\sigma(X, X') < \xi_{SW} \leq \xi$,

and if $\xi_{sw} < \xi$, then we are done. If $\xi_{sw} = \xi$ and since (X, ξ) is a Banach space, there must be a $\sigma(X', X)$ -null sequence $\{a_n\} \subset X'$, whose polar in X is contained in the unit ball of X . It is easy to check that the map, $T: X \rightarrow c_0$, which sends $x \in X$ to the sequence $\{a'_n(x)\} \in c_0$, is an isomorphism of X onto a closed subspace of c_0 . (These results are known, see the author [1].) A classical result of Banach (see Lindenstrauss and Tzafriri [6, p. 53]) says that X must have a subspace isomorphic to c_0 . An application of Lemma 2 and Case I completes the proof if (X, ξ) is a Banach space.

If (X, ξ) is not a Banach space, then X is not a subspace of $B \oplus \omega$, for any Banach space B . Thus a result of Bessaga, Pelczyński and Rolewicz [5] show that (X, ξ) has a nuclear subspace Y . Thus Case II and Lemma 2 completes the proof of the theorem.

REMARKS. It is possible that the following statement is true:

- (*) Each complete space (X, ξ) with $\xi \neq \sigma(X, X')$, has another complete topology η with $\sigma(X, X') < \eta < \xi$.

There are three places in the proof of the theorem where metrizability was used. The most subtle use of the metric was in Lemma 2. If (X, ξ) is not Fréchet, it is possible that X/Y is not complete (Schaefer [6, Ex. 11, p. 192]) and hence Lemma 1 cannot be used to show (X, ζ) is complete. (The author thanks E. G. Ostling for pointing this out to the author.) Thus it is possible that (*) could be true for separable X , but false in general.

If (X, ξ) is separable and complete, then, as in the proof of the theorem (X, ξ_{sw}) is complete (see Bellenot and Ostling [3]). In this case $\xi = \xi_{sw}$ implies that (X, ξ) is a closed subspace of a product of copies of the Banach space c_0 . In order to handle this in the manner of Case I, one must extend this case to include each (X, ξ) which is not inductively semi-reflexive, but for which $\xi = \xi_{sw}$. Examples of spaces which fall into this extended case and which may fail (*) are the spaces (X, ξ_{sw}) where (X, ξ) is any separable nonreflexive Banach space.

The proof that the topology constructed in Case II is complete works for any inductive semi-reflexive space. However, to show that this constructed topology was different from the given topology made strong use of the metrizability. In fact, if $(X, \xi) = \phi$, then for any positive nonincreasing null-sequence, $\{\mu_n\}$, the topology constructed in Case II will be the ξ -topology. It is open question if ϕ is the only such exception among complete separable spaces with a continuous norm. (Weak topologies are also exceptions.) In any case the space ϕ is perhaps the most likely counter-example (among the inductively semi-reflexive spaces) to (*).

REFERENCES

1. S. F. Bellenot, Ph. D. Thesis (Claremont Graduate School 1974).
2. ———, *Prevarieties and intertwined completeness of locally convex spaces*, Math. Ann., **217** (1975), 59–67.
3. S. F. Bellenot and E. G. Ostling, *Completeness and intertwined completeness of locally convex spaces*, Math. Proc. Camb. Phil. Soc., **81** (1977), 25–30.
4. I. A. Berezanskii, *Inductively reflexive, locally convex spaces*, Soviet Math. Doklady, (Translations from Russian) **9** (2) (1968), 1080–1082.
5. C. Bessaga, A. Pelczynski and S. Rolewicz, *On diametral approximative dimension and linear homogeneity of F -spaces*, Bull. Acad. Polon. Sci., **9** (1961), 677–683.
6. J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces I, Sequence Spaces*, Springer Verlag, Berlin, 1977.
7. A. P. Robertson and W. Robertson, *Topological Vector Spaces*, 2nd ed., Cambridge University Press, Cambridge, 1973.
8. H. H. Schaefer, *Topological Vector Spaces*, Springer Verlag, Berlin, 1971.

Received July 14, 1978 and in revised form March 1, 1979. Partially supported by NSF Grant MCS-76-06726-A01.

FLORIDA STATE UNIVERSITY
TALLAHASSEE, FL 32306

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$84.00 a year (6 Vols., 12 issues). Special rate: \$42.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1979 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 83, No. 2

April, 1979

Patrick Robert Ahern, <i>On a theorem of Hayman concerning the derivative of a function of bounded characteristic</i>	297
Walter Allegretto, <i>Finiteness of lower spectra of a class of higher order elliptic operators</i>	303
Leonard Asimow, <i>Superharmonic interpolation in subspaces of $C_c(X)$</i>	311
Steven F. Bellenot, <i>An anti-open mapping theorem for Fréchet spaces</i>	325
B. J. Day, <i>Locale geometry</i>	333
John Erik Fornæss and Steven Krantz, <i>Continuously varying peaking functions</i>	341
Joseph Leonide Gerver, <i>Long walks in the plane with few collinear points</i>	349
Joseph Leonide Gerver and Lawrence Thom Ramsey, <i>On certain sequences of lattice points</i>	357
John R. Graef, Yuichi Kitamura, Takaši Kusano, Hiroshi Onose and Paul Winton Spikes, <i>On the nonoscillation of perturbed functional-differential equations</i>	365
James A. Huckaba and James M. Keller, <i>Annihilation of ideals in commutative rings</i>	375
Anzelm Iwanik, <i>Norm attaining operators on Lebesgue spaces</i>	381
Surjit Singh Khurana, <i>Pointwise compactness and measurability</i>	387
Charles Philip Lanski, <i>Commutation with skew elements in rings with involution</i>	393
Hugh Bardeen Maynard, <i>A Radon-Nikodým theorem for finitely additive bounded measures</i>	401
Kevin Mor McCrimmon, <i>Peirce ideals in Jordan triple systems</i>	415
Sam Bernard Nadler, Jr., Joseph E. Quinn and N. Stavrakas, <i>Hyperspaces of compact convex sets</i>	441
Ken Nakamura, <i>An explicit formula for the fundamental units of a real pure sextic number field and its Galois closure</i>	463
Vassili Nestoridis, <i>Inner functions invariant connected components</i>	473
Vladimir I. Olikier, <i>On compact submanifolds with nondegenerate parallel normal vector fields</i>	481
Lex Gerard Oversteegen, <i>Fans and embeddings in the plane</i>	495
Shlomo Reisner, <i>On Banach spaces having the property $G.L$</i>	505
Gideon Schechtman, <i>A tree-like Tsirelson space</i>	523
Helga Schirmer, <i>Fix-finite homotopies</i>	531
Jeffrey D. Vaaler, <i>A geometric inequality with applications to linear forms</i>	543
William Jennings Wickless, <i>T as an \mathcal{G} submodule of G</i>	555
Kenneth S. Williams, <i>The class number of $Q(\sqrt{-p})$ modulo 4, for $p \equiv 3 \pmod{4}$ a prime</i>	565
James Chin-Sze Wong, <i>On topological analogues of left thick subsets in semigroups</i>	571