

# Pacific Journal of Mathematics

**CONTINUOUSLY VARYING PEAKING FUNCTIONS**

JOHN ERIK FORNAESS AND STEVEN KRANTZ

## CONTINUOUSLY VARYING PEAKING FUNCTIONS

JOHN ERIK FORNAESS AND STEVEN GEORGE KRANTZ

**Let  $X$  be a compact metric space,  $A \subseteq C(X)$  a closed subalgebra. Let  $\mathcal{P} \subseteq X$  be the set of peak points for  $A$ . It is shown that there is a continuous function  $\Phi: \mathcal{P} \rightarrow A$  such that  $\Phi(x)$  peaks at  $x$  for all  $x \in \mathcal{P}$ .**

0. Let  $X$  be a compact Hausdorff space,  $C(X)$  the continuous functions on  $X$  under the uniform norm, and  $A$  a closed subspace of  $C(X)$  containing 1. Let  $\mathcal{P}$  be the set of peak points for  $A$ . Clearly if  $X$  has more than one point and  $x \in \mathcal{P}$  then there are infinitely many functions in  $A$  which peak at  $x$ . Can one construct a function

$$\Phi: \mathcal{P} \longrightarrow A$$

so that  $\Phi(x)$  peaks at  $x$  and  $\Phi$  has some regularity properties?

In [4], using the von Neumann selection principle, it was shown that for  $X = \overline{\mathcal{D}} \subset \subset \mathbb{C}^n$  with smooth boundary,  $A = A(\mathcal{D})$  (the analytic functions on  $\mathcal{D}$  which extend continuously to  $\overline{\mathcal{D}}$ ), one can choose  $\Phi$  to be measurable. The same argument is valid under much more general circumstances.

In the present note we prove that, for quite general  $X$  and for  $A$  an algebra,  $\Phi$  can be chosen to be continuous. This generalizes results in [1, Theorem 3.1] and [2, Proposition 4].

1. Throughout the discussion,  $X$  will be a fixed compact metric space with metric  $d$ . We let  $C(X)$  denote the continuous, complex-valued functions on  $X$  with the uniform norm and  $A \subseteq C(X)$  will be a closed complex linear subspace. If  $x \in X$ ,  $r > 0$ , then  $B(x, r) = \{t \in X: d(x, t) < r\}$ .

**DEFINITION.** A point  $x \in X$  is said to be a *peak point* for  $A$  if there is an  $f \in A$  with  $f(x) = 1$  and, for all  $y \in X \sim \{x\}$ ,  $|f(y)| < 1$ . The function  $f$  is said to peak at  $x$ .

We let  $\mathcal{P}(A)$  denote the set of peak points for  $A$ .

**THEOREM.** *Let  $X$  be a compact metric space,  $A \subseteq C(X)$  a closed subalgebra (with or without 1). Then there is a continuous map*

$$\Phi: \mathcal{P}(A) \longrightarrow A$$

*such that  $\Phi(x)$  peaks at  $x$  for each  $x \in \mathcal{P}(A)$ .*

The remainder of the paper is devoted to the proof of the theorem. We proceed via a sequence of lemmas. The plan of the proof is as follows.

For each  $k \in \{1, 2, \dots\}$  we will construct a continuous function

$$\Phi_k: \mathcal{P}(A) \longrightarrow A$$

such that for each  $x \in \mathcal{P}(A)$  we have

- (i)  $\|\Phi_k(x)\| = 1$ ;
- (ii)  $[\Phi_k(x)](x) = 1$ ;
- (iii) if  $t \in X \sim B(x, 1/k)$  then  $|[\Phi_k(x)](t)| \leq 1 - 1/(k + 2)$ .

Once the  $\{\Phi_k\}$  are constructed, the proof is immediate. For let  $\Phi = \sum_{i=1}^{\infty} 2^{-i} \Phi_i$ . Then  $\Phi$  is continuous and for each  $x \in \mathcal{P}(A)$  we have  $\Phi(x) \in A$  and  $[\Phi(x)](x) = 1$ . Moreover, if  $t \neq x$  and  $k > 1/d(x, t)$  then

$$\begin{aligned} |[\Phi(x)](t)| &\leq \sum_{i \neq k} 2^{-i} |[\Phi_i(x)](t)| + 2^{-k} |[\Phi_k(x)](t)| \\ &\leq 1 - 2^{-k} + 2^{-k}(1 - 1/(k + 2)) < 1. \end{aligned}$$

So  $\Phi(x)$  peaks at  $x$ . Thus it remains to construct the  $\Phi_k$ .

**LEMMA 1.** *Let  $x_0 \in \mathcal{P}(A)$ . Let  $p$  be a strictly positive continuous function on  $X$  with  $p(x_0) = 1$ . Then there is an  $f \in A$  with  $f(x_0) = 1$  and  $|f(x)| \leq p(x)$  for all  $x \in X$ .*

*Proof.* This is a special case of Theorem 12.5 of Gamelin [3], p. 58.

**COROLLARY 2.** *With hypotheses as in Lemma 1, there is a  $g \in A$  such that  $g(x_0) = 1$ ,  $|g(x)| < p(x)$  for all  $x \in X \sim \{x_0\}$ .*

*Proof.* Immediate.

**LEMMA 3.** *Let  $x_0 \in \mathcal{P}(A)$ . Let  $\psi \in A$  peak at  $x_0$ . There is a map*

$$\Psi: \mathcal{P}(A) \cap \{|\psi(x)| > 1/2\} \longrightarrow A$$

so that

- (i)  $\Psi(x)$  peaks at  $x$  for each  $x \in \mathcal{P}(A) \cap \{|\psi(x)| > 1/2\}$ ,
- (ii)  $\Psi(x_0) = \psi$ ,
- (iii)  $\Psi$  is continuous at  $x_0$ .

*Proof.* For each  $x \in \mathcal{P}(A) \sim \{x_0\}$  choose, by Corollary 2, a function  $\varphi_x \in A$  such that  $\varphi_x(x) = 1$  and

$$(*) \quad |\varphi_x(t)| < \min \{ (2 - |\psi(x)| - |\psi(t)|) / 2(1 - |\psi(x)|), 1 \} \\ \text{for all } t \in X \sim \{x\}.$$

Now for each  $x \in \mathcal{P}(A) \cap \{|\psi(x)| > 1/2\}$  we define

$$\Psi(x) = \begin{cases} [2(1 - |\psi(x)|)\varphi_x + \overline{\text{sgn } \psi(x)\psi}]/[2 - |\psi(x)|] & \text{if } x \neq x_0, |\psi(x)| > 1/2, \\ \psi & \text{if } x = x_0. \end{cases}$$

Here  $\text{sgn } z \equiv z/|z|$ , any  $z \in C \sim \{0\}$ .

Clearly if  $x \neq x_0$  and  $x$  is sufficiently close to  $x_0$  then  $|\psi(x)| > 1/2$  and we have

$$\begin{aligned} \|\Psi(x) - \psi\| &\leq \|\Psi(x) - \overline{\text{sgn } \psi(x)} \cdot \psi\| + \|\overline{\text{sgn } \psi(x)} \cdot \psi - \psi\| \\ &\leq \|[2(1 - |\psi(x)|)\varphi_x + \overline{\text{sgn } \psi(x)} \cdot \psi]/[2 - |\psi(x)|] - \overline{\text{sgn } \psi(x)} \cdot \psi\| \\ &\quad + \|\psi(1 - \overline{\text{sgn } \psi(x)})\| \\ &\leq \{[2(1 - |\psi(x)|)\|\varphi_x - \overline{\text{sgn } \psi(x)} \cdot \psi\| \\ &\quad + (1 - |\psi(x)|)\|\overline{\text{sgn } \psi(x)} \cdot \psi\|]/[2 - |\psi(x)|] + |1 - \overline{\text{sgn } \psi(x)}|\} \\ &\leq 5(1 - |\psi(x)|) + |1 - \overline{\text{sgn } \psi(x)}| \\ &\longrightarrow 0 \quad \text{as } x \longrightarrow x_0. \end{aligned}$$

It remains to verify that  $\Psi(x)$  peaks at  $x$  when  $|\psi(x)| > 1/2$ . For such  $x$ , we have  $[\Psi(x)](x) = 1$ . Further, if  $t \neq x$  then by (\*) we have

$$2(1 - |\psi(x)|)|\varphi_x(t)| < 2 - |\psi(x)| - |\psi(t)|$$

or

$$|2(1 - |\psi(x)|)\varphi_x(t)| + |\psi(t)| < 2 - |\psi(x)|$$

whence

$$|2(1 - |\psi(x)|)\varphi_x(t) + \overline{\text{sgn } \psi(x)\psi}(t)| < 2 - |\psi(x)|$$

or

$$|[\Psi(x)](t)| < 1.$$

LEMMA 4. Fix a positive integer  $k$ . There is a sequence  $\{\Phi_k^j\}_{j=1}^\infty$  of functions,

$$\Phi_k^j: \mathcal{P}(A) \longrightarrow A$$

satisfying, for each  $z \in \mathcal{P}(A)$  and every  $j$ ,

- (i)  $\|\Phi_k^j(x)\| = 1$ ;
- (ii)  $[\Phi_k^j(x)](x) = 1$ ;
- (iii)  $\limsup_{\substack{(A) \ni y \rightarrow x}} \|\Phi_k^j(x) - \Phi_k^j(y)\| \leq 4^{-j} \cdot (1/k)$ ;

(iv) for every  $t \in X \sim B(x, (1 - 2^{-j}) \cdot (1/k))$ ,

$$|[\Phi_k^j(x)](t)| \leq (1 - 2/(k + 2)) + \sum_{i=1}^j 2^{-i} \cdot (1/(k + 2)) ;$$

(v)  $\|\Phi_k^j(x) - \Phi_k^{j-1}(x)\| \leq 2^{-j} \cdot (1/k), j \geq 2$ .

*Proof.* This lemma is the heart of the matter. We construct the  $\Phi_k^j$  inductively on  $j$ . First consider  $j = 1$ . For each  $x \in \mathcal{P}(A)$  construct, by Lemma 1, a function  $\varphi_x \in A$  which satisfies  $\varphi_x(x) = 1$  and

$$|\varphi_x(t)| \leq \min \{1 - 8kd(x, t)/(k + 2), 1 - 2/(k + 2)\} .$$

Using  $\psi = \varphi_x$ , construct a function

$$(*) \quad \Psi_x^1: \mathcal{P}(A) \cap \{|\psi(x)| > 1/2\} \longrightarrow A$$

satisfying the conclusions of Lemma 3. Choose  $r_x^1, 0 < r_x^1 < 1/4k$  so that  $t \in B(x, r_x^1)$  implies that  $|\varphi_x(t)| > 1/2$  and

$$\|\Psi_x^1(x) - \Psi_x^1(t)\| < 4^{-2} \cdot (1/(k + 2)) .$$

Now observe that if  $y \in B(x, r_x^1)$  and  $t \notin B(y, 1/2k)$  then

$$d(x, t) \geq d(y, t) - d(y, x) \geq 1/4k .$$

Therefore for such  $y, t$  we have

$$(**) \quad \begin{aligned} |[\Psi_x^1(y)](t)| &\leq |[\Psi_x^1(x)](t)| + |[\Psi_x^1(x)](t) - [\Psi_x^1(y)](t)| \\ &\leq |\varphi_x(t)| + 4^{-2} \cdot (1/(k + 2)) \\ &\leq (1 - 2/(k + 2)) + 2^{-1} \cdot (1/(k + 2)) . \end{aligned}$$

Now since  $\mathcal{P}(A)$  is a metric space, it is paracompact ([5], p. 160, Cor. 35). Hence there is a locally finite refinement  $\mathcal{U}^1 = \{U_\omega^1\}_{\omega \in \Omega_1}$  of the covering  $\{B(x, r_x^1)\}_{x \in \mathcal{P}(A)}$  of  $\mathcal{P}(A)$ . Let  $x_\omega, \omega \in \Omega_1$ , be chosen so that  $U_\omega^1 \subseteq B(x_\omega, r_{x_\omega}^1)$ . Let  $B_\omega^1$  denote  $B(x_\omega, r_{x_\omega}^1)$ . We may assume that  $\bar{U}_\omega^1 \subseteq B_\omega^1$ . Let  $\{\chi_\omega^1\}$  be a continuous partition of unity sub-ordinary to  $\mathcal{U}^1$  and define

$$\Phi_k^1 = \sum_{\omega \in \Omega} \chi_\omega^1 \Psi_{x_\omega}^1 .$$

Then conclusions (i) and (ii) are immediate. Conclusion (iv) follows from (\*\*). Conclusion (v) is vacuous for  $j = 1$ . It remains to verify (iii).

Fix  $x \in \mathcal{P}(A)$ . Then there is a neighborhood  $W$  of  $x$  and  $\{\omega_1, \dots, \omega_m\} \subseteq \Omega_1$  so that  $W \cap \text{supp } \chi_\omega \neq 0$  only if  $\omega \in \{\omega_1, \dots, \omega_m\}$ . Of course  $m$  may depend on  $x$ . Letting  $x_i$  denote  $x_{\omega_i}, i = 1, \dots, m$ , we have that

$$\begin{aligned}
\limsup_{\mathcal{S}(A) \ni y \rightarrow x} \|\Phi_k^i(x) - \Phi_k^i(y)\| &\leq \sum_{i=1}^m \limsup_{\mathcal{S}(A) \cap W \ni y \rightarrow x} |\chi_{\omega_i}^1(x) - \chi_{\omega_i}^1(y)| \|\Psi_{x_i}(y)\| \\
&+ \sum_{i=1}^m \chi_{\omega_i}^1(x) \limsup_{\mathcal{S}(A) \cap W \ni y \rightarrow x} \|\Psi_{x_i}^1(x) - \Psi_{x_i}^1(y)\| \\
&\leq 0 + \sum_{i=1}^m \chi_{\omega_i}^1(x) \limsup_{\mathcal{S}(A) \cap B_{\omega_i} \ni y \rightarrow x} \|\Psi_{x_i}^1(x) - \Psi_{x_i}^1(x_i)\| \\
&+ \sum_{i=1}^m \chi_{\omega_i}^1(x) \limsup_{\mathcal{S}(A) \cap B_{\omega_i} \ni y \rightarrow x} \|\Psi_{x_i}^1(x_i) - \Psi_{x_i}^1(y)\| \\
&\leq 2 \cdot 4^{-2}/(k+2) \leq 4^{-1} \cdot (1/k).
\end{aligned}$$

Now suppose that  $\Phi_k^1, \dots, \Phi_k^j$  have been constructed so that (i)-(v) are satisfied. Let  $x \in \mathcal{S}(A)$ . Using  $\psi = \Phi_k^j(x)$ , we construct a function

$$\Psi_x^{j+1}: \mathcal{S}(A) \cap \{|\psi(x)| > 1/2\} \longrightarrow A$$

satisfying the conclusions of Lemma 3. Choose  $r_x^{j+1}$ ,  $0 < r_x^{j+1} < 2^{-j-1} \cdot (1/k)$  so that  $t \in B(x, r_x^{j+1})$  implies that  $|\Phi_k^j(x)(t)| > 1/2$  and both

$$\|\Psi_x^{j+1}(x) - \Psi_x^{j+1}(t)\| \leq 4^{-j-2} \cdot (1/(k+2))$$

(\*\*\*) and

$$\|\Phi_k^j(x) - \Phi_k^j(t)\| \leq (4/3) \cdot 4^{-j} \cdot (1/k).$$

If now  $y \in B(x, r_x^{j+1})$ ,  $t \in B(y, (1 - 2^{-j-1}) \cdot (1/k))$  then

$$d(x, t) \geq d(y, t) - d(y, x) \geq (1 - 2^{-j})(1/k).$$

Hence for such  $y, t$  we have

$$\begin{aligned}
|[\Psi_x^{j+1}(y)](t)| &\leq |[\Psi_x^{j+1}(x)](t)| + |[\Psi_x^{j+1}(x)](t) - [\Psi_x^{j+1}(y)](t)| \\
&\leq |[\Phi_k^j(x)](t)| + 4^{-j-1} \cdot (1/(k+2)) \\
&\leq (1 - 2/(k+2)) + \sum_{i=1}^j 2^{-i} \cdot (1/(k+2)) + 2^{-j-1} \cdot (1/(k+2)) \\
&= (1 - 2/(k+2)) + \sum_{i=1}^{j+1} 2^{-i} \cdot (1/(k+2)).
\end{aligned}$$

Choose a locally finite refinement  $\mathcal{U}^{j+1} = \{U_\omega^{j+1}\}_{\omega \in \Omega_{j+1}}$  of the covering  $\{B(x, r_x^{j+1})\}_{x \in (A)}$  of  $\mathcal{S}(A)$ . Let  $\{x_\omega\}_{\omega \in \Omega_{j+1}}$  be chosen so that  $U_\omega^{j+1} \subseteq B(x_\omega, r_{x_\omega}^{j+1}) \equiv B_\omega^{j+1}$ , each  $\omega \in \Omega_{j+1}$ . We may assume that  $\bar{U}_\omega^{j+1} \subseteq B_\omega^{j+1}$ . Let  $\{\chi_\omega^{j+1}\}$  be a continuous partition of unity subordinate to  $\mathcal{U}^{j+1}$ . Define

$$\Phi_k^{j+1} = \sum_{\omega \in \Omega_{j+1}} \chi_\omega^{j+1} \Psi_{x_\omega}^{j+1}.$$

It follows as in the case  $j = 1$  that (i), (ii), (iii), and (iv) hold. To verify (v) fix  $x \in \mathcal{S}(A)$ . Let  $\omega_1, \dots, \omega_m$  satisfy the property that

$\mathcal{X}_\omega(x) \neq 0$  iff  $\omega \in \{\omega_1, \dots, \omega_m\}$ . Let  $x_i$  denote  $x_{\omega_i}$ ,  $i = 1, \dots, m$ . Then

$$\begin{aligned} \|\Phi_k^{j+1}(x) - \Phi_k^j(x)\| &\leq \left\| \sum_{l=1}^m \mathcal{X}_{\omega_l}^{j+1}(x) [\Psi_{x_l}^{j+1}(x) - \Psi_{x_l}^j(x)] \right\| \\ &\quad + \left\| \sum_{l=1}^m \mathcal{X}_{\omega_l}^{j+1}(x) [\Psi_{x_l}^{j+1}(x_l) - \Phi_k^j(x_l)] \right\| \\ &\quad + \left\| \sum_{l=1}^m \mathcal{X}_{\omega_l}^{j+1}(x) [\Phi_k^j(x_l) - \Phi_k^j(x)] \right\| \\ &\leq 4^{-j-2}(1/(k+2)) + 0 + (4/3)4^{-j}(1/k) \leq 2^{-j}(1/k). \end{aligned}$$

The induction is complete.

LEMMA 5. For  $k \in \{1, 2, \dots\}$  there exist functions

$$\Phi_k: \mathcal{S}(A) \longrightarrow A$$

such that

- (i)  $\|\Phi_k(x)\| = 1$  for all  $x \in \mathcal{S}(A)$ ,
- (ii)  $[\Phi_k(x)](x) = 1$ ;
- (iii)  $\Phi_k$  is continuous;
- (iv)  $|\Phi_k(x)(t)| \leq 1 - 1/(k+2)$  for all  $x \in \mathcal{S}(A)$ ,  $t \in X \sim B(x, 1/k)$ .

*Proof.* Let  $\Phi_k^j$  be as in Lemma 4 and define  $\Phi_k = \lim_{j \rightarrow \infty} \Phi_k^j$ . That the limit exists follows from (v) of Lemma 4. The conclusions (i)-(iv) of the present lemma now follow from the corresponding parts of Lemma 4.

By the discussion preceding Lemma 1, the proof of the theorem is complete.

REMARK. Our proof yields something more general. Indeed, instead of assuming  $X$  to be metric, one need only assume that the relative topology on  $\mathcal{S}$  has a  $\sigma$ -locally finite base. By [5], p. 128, this is equivalent to assuming that  $\mathcal{S}$  is metric, hence paracompact, and the proof goes through as before.

The referee has kindly observed that given our Lemma 3, one can use Theorem 3.1'' of [6] to prove that if  $X$  is compact Hausdorff and  $A$  is separable then the theorem holds. This is a weaker result than the one outlined in the preceding paragraph. Moreover, the proof using [6] is not essentially shorter than the elementary one presented here, and the construction of  $\Phi$  as the uniform limit of discontinuous functions has intrinsic interest.

REMARK. It would be interesting to know whether, in the presence of differentiable structure in  $X$  and  $A$ , the peaking functions may be chosen to vary differentiably.

## REFERENCES

1. E. Bedford and J. E. Fornaess, *A construction of peak functions on weakly pseudoconvex domains*, Ann. of Math., **107** (1978), 555-568.
2. I. Graham, *Boundary behavior of the Caratheodory and Kobayashi metrics on strongly pseudoconvex domains in  $C^n$* , Trans. Amer. Math. Soc., **207** (1975), 219-240.
3. T. W. Gamelin, *Uniform Algebras*, Englewood Cliffs, Academic Press, 1969.
4. N. P. Jewell and S. G. Krantz, *Toeplitz operators and related function algebras on certain pseudoconvex domains*, Trans. Amer. Math. Soc., **252** (1979), 297-312.
5. J. L. Kelley, *General Topology*, New York, Van Nostrand, 1955.
6. E. Michael, *Continuous selections*, I, Ann. of Math., **63** (1956), 361-382.

Received October 30, 1978 and in revised form January 8, 1979. The work of the first author was supported in part under NSF Grant No. MCS 78-01488. The work of the second author was supported in part under NSF Grant No. MCS 77-02213.

PRINCETON UNIVERSITY

PRINCETON, NJ 08540

AND

UNIVERSITY OF CALIFORNIA

LOS ANGELES, CA 90024



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California  
Los Angeles, CA 90024

HUGO ROSSI

University of Utah  
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University  
Stanford, CA 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$84.00 a year (6 Vols., 12 issues). Special rate: \$42.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).

8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1979 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

Patrick Robert Ahern, <i>On a theorem of Hayman concerning the derivative of a function of bounded characteristic</i> .....	297
Walter Allegretto, <i>Finiteness of lower spectra of a class of higher order elliptic operators</i> .....	303
Leonard Asimow, <i>Superharmonic interpolation in subspaces of <math>C_c(X)</math></i> .....	311
Steven F. Bellenot, <i>An anti-open mapping theorem for Fréchet spaces</i> .....	325
B. J. Day, <i>Locale geometry</i> .....	333
John Erik Fornæss and Steven Krantz, <i>Continuously varying peaking functions</i> .....	341
Joseph Leonide Gerver, <i>Long walks in the plane with few collinear points</i> .....	349
Joseph Leonide Gerver and Lawrence Thom Ramsey, <i>On certain sequences of lattice points</i> .....	357
John R. Graef, Yuichi Kitamura, Takaši Kusano, Hiroshi Onose and Paul Winton Spikes, <i>On the nonoscillation of perturbed functional-differential equations</i> .....	365
James A. Huckaba and James M. Keller, <i>Annihilation of ideals in commutative rings</i> .....	375
Anzelm Iwanik, <i>Norm attaining operators on Lebesgue spaces</i> .....	381
Surjit Singh Khurana, <i>Pointwise compactness and measurability</i> .....	387
Charles Philip Lanski, <i>Commutation with skew elements in rings with involution</i> .....	393
Hugh Bardeen Maynard, <i>A Radon-Nikodým theorem for finitely additive bounded measures</i> .....	401
Kevin Mor McCrimmon, <i>Peirce ideals in Jordan triple systems</i> .....	415
Sam Bernard Nadler, Jr., Joseph E. Quinn and N. Stavrakas, <i>Hyperspaces of compact convex sets</i> .....	441
Ken Nakamura, <i>An explicit formula for the fundamental units of a real pure sextic number field and its Galois closure</i> .....	463
Vassili Nestoridis, <i>Inner functions invariant connected components</i> .....	473
Vladimir I. Olikier, <i>On compact submanifolds with nondegenerate parallel normal vector fields</i> .....	481
Lex Gerard Oversteegen, <i>Fans and embeddings in the plane</i> .....	495
Shlomo Reisner, <i>On Banach spaces having the property G.L.</i> .....	505
Gideon Schechtman, <i>A tree-like Tsirelson space</i> .....	523
Helga Schirmer, <i>Fix-finite homotopies</i> .....	531
Jeffrey D. Vaaler, <i>A geometric inequality with applications to linear forms</i> .....	543
William Jennings Wickless, <i>T as an <math>\mathcal{Q}</math> submodule of G</i> .....	555
Kenneth S. Williams, <i>The class number of <math>Q(\sqrt{-p})</math> modulo 4, for <math>p \equiv 3 \pmod{4}</math> a prime</i> .....	565
James Chin-Sze Wong, <i>On topological analogues of left thick subsets in semigroups</i> .....	571