

Pacific Journal of Mathematics

NORM ATTAINING OPERATORS ON LEBESGUE SPACES

ANZELM IWANIK

NORM ATTAINING OPERATORS ON LEBESGUE SPACES

ANZELM IWANIK

Let X and Y be Lebesgue spaces (AL-spaces). Then the norm attaining operators mapping X to Y are dense in the space of all linear bounded operators from X to Y .

For any two real Banach spaces X and Y by $B(X, Y)$ we denote the Banach space of all bounded linear operators from X to Y . In [7] Uhl proved that for any strictly convex Banach space Y the norm attaining operators are (norm) dense in $B(L^1[0, 1], Y)$ if and only if Y has the Radon-Nikodym property. The question of whether the norm attaining operators are dense in $B(L^1[0, 1], L^1[0, 1])$ has remained unsolved (cf. [7], p. 299). Here we answer this question in the affirmative. In fact we prove a slightly more general result.

First we introduce some notations. Let I stand for the unit interval. For any function μ defined on the product algebra in $I \times I$ by $\mu^i (i = 1, 2)$ we denote the corresponding marginal functions defined on the Borel subsets of I :

$$\begin{aligned}\mu^1(A) &= \mu(A \times I) , \\ \mu^2(B) &= \mu(I \times B) .\end{aligned}$$

The vector lattice of all finite signed Borel measures on $I \times I$ will be denoted by M . Given any two finite positive Borel measures m_1, m_2 on I we write $M(m_1, m_2)$ for the set of all measures μ in M such that $|\mu|^i$ is absolutely continuous with respect to $m_i (i = 1, 2)$ and

$$\frac{d|\mu|^i}{dm_i} \in L^\infty(m_i) .$$

The measures m_1 and m_2 will be fixed throughout the rest of the paper.

Let us recall that $B(L^1(m_1), L^1(m_2))$ is a Banach lattice under its canonical order (see [5], IV Theorem 1.5 (ii)).

The forthcoming theorem establishes an isomorphism between $M(m_1, m_2)$ and $B(L^1(m_1), L^1(m_2))$, and extends a corresponding result of J. R. Brown on doubly stochastic operators ([1], p. 18). As was kindly indicated by the referee, our Theorem 1 is also related

to N. J. Kalton's representation of the endomorphisms from L^p to L^p for $0 < p \leq 1$ ([3], Theorem 3.1).

By $\langle \cdot, \cdot \rangle$ we denote the canonical bilinear form on $L^\infty(m_2)^* \times L^\infty(m_2)$.

THEOREM 1. *The space $M(m_1, m_2)$ is a vector lattice ideal in M and to each $\mu \in M(m_1, m_2)$ there corresponds a unique operator $T_\mu \in B(L^1(m_1), L^1(m_2))$ such that*

$$\langle T_\mu f, h \rangle = \int f(x)h(y)d\mu(x, y)$$

for all $f \in L^1(m_1)$ and $h \in L^\infty(m_2)$. Moreover, the mapping $\mu \rightarrow T_\mu$ is a vector lattice isomorphism of $M(m_1, m_2)$ onto $B(L^1(m_1), L^1(m_2))$ and

$$\|T_\mu\| = \left\| \frac{d|\mu|^1}{dm_1} \right\|_\infty$$

for every $\mu \in M(m_1, m_2)$.

Proof. First we note that $M(m_1, m_2)$ is a vector subspace of M . Since $\nu \in M(m_1, m_2)$ whenever $0 \leq \nu \in M$ and $\nu \leq \mu \in M(m_1, m_2)$, we observe that $M(m_1, m_2)$ is a lattice ideal (and clearly a sublattice) in M . If $\mu \in M(m_1, m_2)$ then it is easy to see that the bilinear form

$$[f, h] = \int f(x)h(y)d\mu(x, y)$$

is well-defined and continuous on $L^1(m_1) \times L^\infty(m_2)$. Therefore there exists a unique operator $T_\mu \in B(L^1(m_1), L^\infty(m_2)^*)$ such that

$$[f, h] = \langle T_\mu f, h \rangle$$

(see e.g., [5], IV §2). Clearly the mapping $\mu \rightarrow T_\mu$ is one-to-one and $\mu \geq 0$ if and only if T_μ is a positive operator in the Banach lattice sense. Moreover, for an arbitrary $\nu \geq 0$ in $M(m_1, m_2)$ and for any $h \in L^\infty(m_2)$ we have $\langle T_\nu 1, h \rangle = \int h d\nu^2$, so

$$T_\nu 1 = \frac{d\nu^2}{dm_2} \in L^1(m_2),$$

whence $T_\nu f \in L^1(m_2)$ for any $f \in L^\infty(m_1)$. Consequently, $T_\nu \in B(L^1(m_1), L^1(m_2))$ by continuity. Since every $\mu \in M(m_1, m_2)$ is a difference of two positive measures in $M(m_1, m_2)$ and $\mu \rightarrow T_\mu$ is a linear map, we have $T_\mu \in B(L^1(m_1), L^1(m_2))$ for all $\mu \in M(m_1, m_2)$.

We now show that $\mu \rightarrow T_\mu$ is an "onto" mapping. Since $B(L^1(m_1), L^1(m_2))$ is a Banach lattice, it suffices to prove that every

positive operator $T \in B(L^1(m_1), L^1(m_2))$ is of the form T_μ . Given any such T we define a set function

$$\mu(A \times B) = \langle T\chi_A, \chi_B \rangle$$

on all Borel rectangles in $I \times I$. Evidently μ extends uniquely to a finitely additive positive measure (denoted also by μ) on the product algebra. The marginal measures $\mu^1(A) = \int_A T^*1 dm_1$ and $\mu^2(B) = \int_B T1 dm_2$ are finite, positive, and countably additive, so they are compact by the classical result of Ulam. Since μ is a nondirect product of μ^1 and μ^2 , it is countably additive by Theorem 1 (i) in [4]. The unique extension of μ to a finite positive (countably additive) Borel measure on $I \times I$ is again denoted by μ . By a standard approximation argument,

$$\int f(x)h(y)d\mu(x, y) = \langle Tf, h \rangle$$

for all $f \in L^1(m_1)$ and $h \in L^\infty(m_2)$. Therefore $T = T_\mu$. Finally, we note that for every $\mu \in M(m_1, m_2)$

$$\begin{aligned} \|T_\mu\| &= \|T_{|\mu^1|}\| = \sup \|T_{|\mu^1|}f\|_1 = \sup \langle T_{|\mu^1|}f, 1 \rangle \\ &= \sup \int f(x)d|\mu^1|(x) = \sup \int f(x)\frac{d|\mu^1|}{dm_1}(x)dm_1(x) \\ &= \left\| \frac{d|\mu^1|}{dm_1} \right\|_\infty, \end{aligned}$$

where the suprema are taken over all nonnegative functions $f \in L^1(m_1)$ with $\|f\|_1 \leq 1$.

COROLLARY 1. *Let $\nu \in M(m_1, m_2)$. If there exists a function $g \in L^\infty(m_2)$ with $|g| = 1$ such that the Radon-Nikodym derivative of the marginal measure $(g(y)d\nu(x, y))^1$ with respect to m_1 equals*

$$\left\| \frac{d|\nu^1|}{dm_1} \right\|_\infty$$

on a set B of positive m_1 measure, then the operator T_ν attains its norm on the unit ball in $L^1(m_1)$.

Proof. We put $d\lambda(x, y) = g(y)d\nu(x, y)$. Then

$$\begin{aligned} \langle T_\nu(\chi_B/m_1(B)), g \rangle &= \frac{1}{m_1(B)} \int \chi_B(x)d\lambda(x, y) \\ &= \frac{1}{m_1(B)} \int_B \frac{d\lambda^1}{dm_1} dm_1 = \left\| \frac{d|\nu^1|}{dm_1} \right\|_\infty, \end{aligned}$$

implying $\|T_*(\mathcal{X}_B/m_1(B))\|_1 = \|T_*\|$ by Theorem 1.

The algebra of sets generated by all dyadic-rational rectangles in $I \times I$ will be denoted by \mathcal{A} . The σ -algebra generated by \mathcal{A} coincides with the Borel algebra in $I \times I$.

THEOREM 2. *The norm attaining operators are dense in $B(L^1(m_1), L^1(m_2))$.*

Proof. Let $T \in B(L^1(m_1), L^1(m_2))$. By Theorem 1 we have $T = T_\mu$ for some measure μ in $M(m_1, m_2)$. Without any loss of generality we may assume

$$\left\| \frac{d|\mu|^1}{dm_1} \right\|_\infty = 1.$$

Given $0 < \varepsilon < 1$, the set

$$D = \left\{ x \in I: \frac{d|\mu|^1}{dm_1}(x) > 1 - \frac{\varepsilon}{4} \right\}$$

is of positive m_1 measure, say, $m_1(D) = \delta > 0$. Now let $P, (I \times I) - P$ be the Hahn decomposition for μ with μ^+ concentrated on P (see [2], § 29 Theorem A). Since P is a Borel set, there exists $\tilde{P} \in \mathcal{A}$ such that $|\mu|(P \Delta \tilde{P}) < \delta\varepsilon/4$ ([2], § 13 Theorem D). We define a new measure $\tilde{\mu}$ by

$$d\tilde{\mu} = \mathcal{X}_{\tilde{P}}d\mu^+ - \mathcal{X}_{(I \times I) - \tilde{P}}d\mu^-.$$

Evidently $\tilde{P}, (I \times I) - \tilde{P}$ is the Hahn decomposition for $\tilde{\mu}$ and $d|\mu - \tilde{\mu}| = \mathcal{X}_{P \Delta \tilde{P}}d|\mu|$. Since $|\mu - \tilde{\mu}|(I \times I) < \delta\varepsilon/4$, the Radon-Nikodym derivative of $|\mu - \tilde{\mu}|^1$ with respect to m_1 must be less than $\varepsilon/4$ on some set $C \subset D$ of positive m_1 measure. As $\tilde{P} \in \mathcal{A}$, there exists a natural number n such that \tilde{P} is a union of finitely many squares corresponding to the dyadic partition of I into 2^n subintervals of equal length. Let I_0 be any such open subinterval intersecting C on a set $B = C \cap I_0$ of positive m_1 measure. We let

$$d\nu(x, y) = \mathcal{X}_B(x) \left(\frac{d|\mu|^1}{dm_1} \right)^{-1}(x) d\tilde{\mu}(x, y) + \mathcal{X}_{I-B}(x) d\mu(x, y).$$

Note first that

$$\begin{aligned} d|\nu - \mu| &= \mathcal{X}_B(x) \left(\frac{d|\mu|^1}{dm_1} \right)^{-1}(x) |d(\tilde{\mu} - \mu)(x, y)| \\ &+ \left(1 - \frac{d|\mu|^1}{dm_1}(x) \right) d\mu(x, y) \Big| \leq 2\mathcal{X}_C(x) d|\tilde{\mu} - \mu|(x, y) + \frac{\varepsilon}{2} d|\mu|(x, y). \end{aligned}$$

Therefore

$$\frac{d|\nu - \mu|^1}{dm_1} < 2\frac{\varepsilon}{4} + \frac{\varepsilon}{2} = \varepsilon,$$

whence $\|T_\nu - T_\mu\| = \|T_{\nu-\mu}\| \leq \varepsilon$. Moreover,

$$\frac{d|\mu|^1}{dm_1} = 1 \text{ on } B \text{ and } \leq 1 \text{ elsewhere.}$$

The set $(I_0 \times I) \cap \tilde{P}$ is a finite union of squares of the form $I_0 \times I_k (k = 1, \dots, m)$, where each I_k is an element of the dyadic partition of I into 2^n subintervals of equal length. Therefore $(B \times I) \cap \tilde{P}$ is the finite union of the Borel rectangles $B \times I_k$. We define a function $g \in L^\infty(m_2)$ as follows

$$g(y) = \begin{cases} 1 & \text{if } y \in \cup I_k, \\ -1 & \text{otherwise.} \end{cases}$$

Clearly the Radon-Nikodym derivative of the marginal measure $(g(y)d\nu(x, y))^1$ coincides with

$$\frac{d|\nu|^1}{dm_1} = 1$$

on B . Therefore, by Corollary 1, T_ν attains its norm and the proof is completed.

By the known representation theorems for Lebesgue spaces (see e.g., [5], II 8.5 Corollary and [2], §41 Theorem C, or [6], 26.4.9 Exercise (C)), every separable Lebesgue space (i.e., separable AL-space in terms of [5]) is Banach lattice isomorphic with $L^1(m)$ for some finite positive Borel measure m on I . Therefore we obtain the following corollary to our result:

COROLLARY 2. *Let X and Y be separable Lebesgue spaces. Then the norm attaining operators are dense in $B(X, Y)$.*

After the paper was accepted for publication, the last corollary has been generalized to arbitrary (nonseparable) Lebesgue spaces as a result of the author's conversations with Professors J. Bourgain and H. P. Lotz. The proof is outlined below:

Theorem 1 remains true if we replace (I, m_i) by (J_i, m_i) with J_i compact Hausdorff and m_i a finite regular (compact) positive measure on the Borel σ -algebra \mathcal{B}_i , and with M being the space of all finite signed measures on the product σ -algebra $\mathcal{B}_1 \times \mathcal{B}_2$. Indeed, the marginal measures $\int_A T^*1 dm_1, \int_B T1 dm_2$ are compact since the measures m_i are regular, and so Theorem 1 (i) of [4] is still

applicable. The rest of the proof remains unchanged.

Theorem 2 is valid for the general spaces $L^1(J_i, m_i)$ with essentially the same proof as before, \mathcal{A} being replaced now by the algebra of all finite unions of Borel rectangles in $J_1 \times J_2$.

Now if X_1, X_2 are arbitrary Lebesgue spaces then every $T \in B(X_1, X_2)$ can be approximated by norm attaining operators. Indeed, let (x_n) be a sequence in X_1 such that $\|x_n\| \leq 1$ and $\lim \|Tx_n\| = \|T\|$. The Banach lattice ideal Y_1 spanned by (x_n) is a Lebesgue subspace with a weak order unit. Also the image TY_1 is contained in a Lebesgue subspace $Y_2 \subset X_2$ with a weak order unit. By the Kakutani representation theorem there exist compact spaces J_i with finite regular positive measures m_i such that $Y_i = L^1(J_i, m_i)$. By the above, the restriction T_1 of T to Y_1 can be approximated within a given $\varepsilon > 0$ by a norm attaining operator $T_0 \in B(Y_1, Y_2)$ satisfying $\|T_0\| = \|T\|$. If P denotes the canonical band projection of X_1 onto Y_1 then it is easy to see that $T_0P + T(I - P)$ has norm $\|T_0\|$, is norm attaining, and approximates T within ε .

REFERENCES

1. J. R. Brown, *Approximation theorems for Markov operators*, Pacific J. Math., **16** (1966), 13-23.
2. P. R. Halmos, *Measure Theory*, Springer Verlag, New York-Heidelberg-Berlin, 1974.
3. N. J. Kalton, *The endomorphisms of $L_p(0 \leq p \leq 1)$* , Indiana Univ. Math. J., **27** (1978), 353-381.
4. E. Marczewski and C. Ryll-Nardzewski, *Remarks on the compactness and nondirect products of measures*, Fund. Math., **40** (1953), 165-170.
5. H. H. Schaefer, *Banach Lattices and Positive Operators*, Springer Verlag, Berlin-Heidelberg-New York, 1974.
6. Z. Semadeni, *Banach Spaces of Continuous Functions*, Polish Scientific Publishers, Warsaw, 1971.
7. J. J. Uhl, Jr., *Norm attaining operators on $L^1 [0, 1]$ and the Radon-Nikodym property*, Pacific J. Math., **63** (1976), 293-300.

Received May 30, 1978 and in revised form October 31, 1978.

TECHNICAL UNIVERSITY
WROCLAW, POLAND

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$84.00 a year (6 Vols., 12 issues). Special rate: \$42.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1979 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 83, No. 2

April, 1979

Patrick Robert Ahern, <i>On a theorem of Hayman concerning the derivative of a function of bounded characteristic</i>	297
Walter Allegretto, <i>Finiteness of lower spectra of a class of higher order elliptic operators</i>	303
Leonard Asimow, <i>Superharmonic interpolation in subspaces of $C_c(X)$</i>	311
Steven F. Bellenot, <i>An anti-open mapping theorem for Fréchet spaces</i>	325
B. J. Day, <i>Locale geometry</i>	333
John Erik Fornæss and Steven Krantz, <i>Continuously varying peaking functions</i>	341
Joseph Leonide Gerver, <i>Long walks in the plane with few collinear points</i>	349
Joseph Leonide Gerver and Lawrence Thom Ramsey, <i>On certain sequences of lattice points</i>	357
John R. Graef, Yuichi Kitamura, Takaši Kusano, Hiroshi Onose and Paul Winton Spikes, <i>On the nonoscillation of perturbed functional-differential equations</i>	365
James A. Huckaba and James M. Keller, <i>Annihilation of ideals in commutative rings</i>	375
Anzelm Iwanik, <i>Norm attaining operators on Lebesgue spaces</i>	381
Surjit Singh Khurana, <i>Pointwise compactness and measurability</i>	387
Charles Philip Lanski, <i>Commutation with skew elements in rings with involution</i>	393
Hugh Bardeen Maynard, <i>A Radon-Nikodým theorem for finitely additive bounded measures</i>	401
Kevin Mor McCrimmon, <i>Peirce ideals in Jordan triple systems</i>	415
Sam Bernard Nadler, Jr., Joseph E. Quinn and N. Stavrakas, <i>Hyperspaces of compact convex sets</i>	441
Ken Nakamura, <i>An explicit formula for the fundamental units of a real pure sextic number field and its Galois closure</i>	463
Vassili Nestoridis, <i>Inner functions invariant connected components</i>	473
Vladimir I. Olikier, <i>On compact submanifolds with nondegenerate parallel normal vector fields</i>	481
Lex Gerard Oversteegen, <i>Fans and embeddings in the plane</i>	495
Shlomo Reisner, <i>On Banach spaces having the property $G.L$</i>	505
Gideon Schechtman, <i>A tree-like Tsirelson space</i>	523
Helga Schirmer, <i>Fix-finite homotopies</i>	531
Jeffrey D. Vaaler, <i>A geometric inequality with applications to linear forms</i>	543
William Jennings Wickless, <i>T as an \mathcal{G} submodule of G</i>	555
Kenneth S. Williams, <i>The class number of $Q(\sqrt{-p})$ modulo 4, for $p \equiv 3 \pmod{4}$ a prime</i>	565
James Chin-Sze Wong, <i>On topological analogues of left thick subsets in semigroups</i>	571