

Pacific Journal of Mathematics

POINTWISE COMPACTNESS AND MEASURABILITY

SURJIT SINGH KHURANA

POINTWISE COMPACTNESS AND MEASURABILITY

SURJIT SINGH KHURANA

Among other results it is proved that if (X, \mathfrak{A}, μ) is a probability space, E a Hausdorff locally convex space such that $(E', \sigma(E', E))$ contains an increasing sequence of absolutely convex compact sets with dense union, and $f: X \rightarrow E$ weakly measurable with $f(X) \subset K$, a weakly compact convex subset of E , then f is weakly equivalent to $g: X \rightarrow E$ with $g(X)$ contained in a separable subset of K .

In [8] and [9] some remarkable results are obtained for the pointwise compact subsets of measurable real-valued functions and some interesting applications to strongly measurable Banach space-valued functions are established. In this paper we continue those ideas a little further. We first give a somewhat different proof of ([9], Theorem 1) and then apply it to give a generalization of classical Phillip's theorem ([5]). Also some result about equicontinuous subsets of $C(X)$, the space of all continuous real-valued functions on (X, τ_ρ) (τ_ρ is the lifting topology, [10], p. 59; in [8] this topology is denoted by T_ρ) are obtained.

All locally convex spaces are taken over reals and notations of [6] are used. For a topological space Y , $C(Y)$ (resp. $C_b(Y)$) will denote the set of all (resp. all bounded) real-valued continuous functions of Y . N will denote the set of natural numbers.

In this paper (X, \mathfrak{A}, μ) is a complete probability measure space. Let \mathcal{L} be the set of all real-valued \mathfrak{A} -measurable functions on X , \mathcal{L}^∞ , the essentially bounded elements of \mathcal{L} , and M^∞ , the bounded elements of \mathcal{L} . We fix a lifting, [10], $\rho: \mathcal{L}^\infty \rightarrow M^\infty$ and on X we always take the lifting topology τ_ρ ([10], p. 59). For $f \in \mathcal{L}$, $g \in \mathcal{L}$, we write $f = g$ if $f(x) = g(x)$, $\forall x \in X$, and $f \equiv g$ if $f(x) = g(x)$, a.e. $[\mu]$. For a Hausdorff locally convex space E , a function $f: X \rightarrow E$ is said to be weakly measurable if $h \circ f$ is \mathfrak{A} -measurable, $\forall h \in E'$, the topological dual of E . Two weakly measurable functions $f_i: X \rightarrow E$, $i = 1, 2$, are said to be weakly equivalent if $h \circ f_1 \equiv h \circ f_2$, $\forall h \in E'$. The space \mathcal{L}_1 and norms $\|\cdot\|_1$ and $\|\cdot\|_\infty$ have the usual meanings. We shall call a topological space, countably compact if every sequence in it has a cluster point, and sequentially compact if every sequence has a convergent subsequence.

We start with a different proof of the following result of [9].

THEOREM 1 ([9], *Theorem 1*). *Let H be a subset of \mathcal{L} such that for any $h_1 \in H$, $h_2 \in H$, $h_1 \neq h_2$ implies $h_1 \neq h_2$. Then, with the pointwise topology on H , the following are equivalent:*

- (i) H is sequentially compact;
- (ii) H is compact and metrizable.

If H is convex, then each of (i) and (ii) is also equivalent to:

- (iii) H is compact;
- (iv) H is countably compact.

Proof. By ([6], Theorem 11.2, p. 187) each of (i), (ii), (iii), (iv) implies that H is relatively compact in R^X , with product topology. Thus each of these conditions implies that H is pointwise bounded. Denote by φ the homeomorphism, $[0, \infty) \rightarrow [0, 1]$, $x \rightarrow x/(1+x)$. For any $\alpha \in I$, the directed net of all finite subsets of H , let $h_\alpha = \sup\{h : h \in \alpha\}$, and $p_\alpha = \rho(\varphi \circ h_\alpha)$. $\{p_\alpha\}$ is a monotone bounded net in $C_b(X)$, which is boundedly complete. Let $\sup p_\alpha = p \in C_b(X)$. This means there is an increasing sequence $\{\alpha(n)\} \subset I$ such that $p = \sup p_{\alpha(n)}$ (this follows from the fact that $\mu(p) = \sup \mu(p_\alpha)$). Since $p_\alpha \equiv \varphi \circ h_\alpha$, we get $p_\alpha^{-1}\{1\}$ is μ -null, $\forall \alpha$. From this it follows that $K = p^{-1}\{1\}$ is μ -null. Thus $q = (\varphi^{-1} \circ p)\chi_{X \setminus K}$ is a measurable function such that $|h| \leq q$ a.e. $[\mu]$, $\forall h \in H$.

(i) \Rightarrow (ii) is simple ([8], Prop. 1, p. 197), the metric d of (ii) being defined by $d(f, g) = \|(f - g)/1 + q\|_1$. (ii) \Rightarrow (iii) and (iii) \Rightarrow (iv) are trivial. Now we come to the proof of (iv) \Rightarrow (i). Take a sequence $\{f'_n\} \subset H$. Since $1/(1+q)H$ is relatively weakly compact in $(\mathcal{L}_1, \|\cdot\|_1)$ there exists a subsequence $\{f_n\}$ of $\{f'_n\}$ and an $f_0 \in \mathcal{L}_1$ such that $1/(1+q)f_n \rightarrow f_0$ weakly. Thus there exists a sequence $\{g_n\}$ in the convex hull of $\{f_n : 1 \leq n < \infty\}$ (note $\{g_n\} \subset H$) such that $1/(1+q)g_n \rightarrow f_0$ a.e. $[\mu]$ (because a convergent sequence in $(\mathcal{L}_1, \|\cdot\|_1)$ has a subsequence converging a.e. $[\mu]$). Taking f to be a cluster point of $\{g_n\}$ in H , we get $1/(1+q)f \equiv f_0(\mu)$. We claim $f_n \rightarrow f$ in H . If $f_n \not\rightarrow f$ there exists an $x \in X$, an $\varepsilon > 0$, and a subsequence $\{f''_n\}$ of $\{f_n\}$ such that one of the two following conditions are satisfied:

- (i) $f''_n(x) > f(x) + \varepsilon$, $\forall n$;
- (ii) $f''_n(x) < f(x) - \varepsilon$, $\forall n$.

Since $1/(1+q)f''_n \rightarrow 1/(1+q)f$ weakly, proceeding as before we get a sequence $\{g''_n\}$ in the convex hull of $\{f''_n : 1 \leq n < \infty\}$ such that $1/(1+q)g''_n \rightarrow 1/(1+q)f$ a.e. $[\mu]$. If f'' is a cluster point of $\{g''_n\}$ in H we get $f'' \equiv f(\mu)$ but because of (i) or (ii), $f''(x) \neq f(x)$, a contradiction. This proves that H is sequentially compact.

This result is also proved in [11] by a different method.

By a classical theorem of Phillips [5], if $f: X \rightarrow E$, E being a Banach space, is weakly measurable and $f(X)$ is relatively weakly compact in E , then f is weakly equivalent to a strongly measurable function ([8], Theorem 3, p. 200). What one really needs to do is to find a weakly equivalent function g such that $g(X)$ is separable. The next theorem is a generalization of Phillips' theorem.

THEOREM 2. *Let (E, \mathcal{S}) be a Hausdorff locally convex space such that there exists an increasing sequence $\{A_n\}$ of absolutely convex compact subsets of $(E', \sigma(E', E))$ whose union is dense in $(E', \sigma(E', E))$. Suppose $f: X \rightarrow E$ is weakly measurable and $f(X) \subset K$, for some weakly compact convex subset of E . Then there exists a weakly measurable function $g: X \rightarrow E$, $g \equiv f(w)$ and $g(X) \subset K_0$, a separable closed convex subset of K .*

Proof. Since $(E, \sigma(E, E'))$ can be considered as a subspace of $R^{E'}$, with product topology, f can be considered as $f: X \rightarrow R^{E'}$. For each $h \in E'$, define $g(h) = \rho(h \circ f)$ and let $g: X \rightarrow R^{E'}$, $(g)_h = g(h)$, $\forall h \in E'$. g is evidently continuous. If $g(x_0) \notin K$ for some $x_0 \in X$, there exists, by separation theorem ([6], p. 65), an $h \in E'$ such that $h \circ g(x_0) > \sup(K)$. This is a contradiction since $h \circ f \leq \sup h(K)$ implies $\rho(h \circ f) \leq \sup h(K)$. Evidently $g \equiv f(w)$. Fix $n \in N$. By Theorem 1, $B_n = \{h \circ g: h \in A_n\}$, with the topology of pointwise convergence on X , is a compact metric space. We metrize E by the seminorms p_n , $p_n(x) = \sup \{ |h(x)| : h \in A_n \}$. We denote this metric topology by \mathcal{S}_0 . For each n , $E_n = (C(B_n), \|\cdot\|)$ is a separable Banach space (here $\|\cdot\|$ is sup norm), and so $F = \prod_{n=1}^{\infty} E_n$ is a separable Frechet space. Let X_0 be the quotient space obtained from X by the equivalent relation, $x \equiv y \Leftrightarrow g(x) = g(y)$. Each $x \in X_0$ gives rise to $x \in C(B_n)$, $x(t) = t(x)$ for each $t \in B_n$, for every n . Thus X_0 can be embedded in F , $x_0 \rightarrow (x_0, x_0, \dots) \in F$. Taking, on X_0 , the topology induced by F , we easily verify that $g: X_0 \rightarrow (E, \mathcal{S}_0)$ is continuous and so $(g(X), \mathcal{S}_0)$ is separable. Let $K_0 =$ the closed convex hull, in (E, \mathcal{S}_0) , of a countable dense subset of $(g(X), \mathcal{S}_0)$. If $g(X) \not\subset K_0$, by separation theorem, there exists an $h \in E'$ and $x_0 \in X$ such that $h \circ g(x_0) > \sup h(K_0)$. Since $(E, \mathcal{S}_0)' \supset \bigcup_{n=1}^{\infty} A_n$, $q \circ g(x_0) \leq \sup q(K_0)$, $\forall q \in \bigcup_{n=1}^{\infty} A_n$. Now there exists a net $\{h_\alpha\} \subset \bigcup_{n=1}^{\infty} A_n$ such that $h_\alpha \rightarrow h$ uniformly on each compact convex subset of $(E, \sigma(E, E'))$. From this it follows $h \circ g(x_0) \leq \sup h(K_0)$, a contradiction. This proves the result.

REMARK 3. If E is metrizable then $(E', \sigma(E', E))$ contains a sequence of compact absolutely convex sets whose union is E' . If Y is a completely regular Hausdorff space containing a σ -compact dense set and $E = C_i(Y)$ with strict topology β_0, β_1 , then it is

proved in ([3], Theorem 3) that $(E', \sigma(E', E))$ has an increasing sequence of absolutely convex compact sets with dense union — here E is not metrizable.

REMARK 4. The function $g: X \rightarrow (E, \sigma(E, E'))$, obtained in this theorem, is measurable in the sense of ([2], Def. 4, p. 89).

The next theorem, in some sense, is a generalization of ([9], Theorem 3).

THEOREM 5. *Let E be a Hausdorff locally convex space such that there exist, in $(E', \sigma(E', E))$, an increasing sequence $\{A_n\}$ of absolutely convex compact sets whose union is E' . Suppose $g: X \rightarrow E$ is weakly measurable such that $g \circ f \neq 0$ implies $g \circ f \neq 0$, for every $f \in E'$. Then $g(X)$ is contained in a separable subspace of E .*

Proof. In the notations of Theorem 2, $B_n = \{h \circ g: h \in A_n\}$ are compact and metrizable, with the topology of pointwise convergence, and \mathcal{T}_0 is the metric topology, on E , of uniform convergence on A_n . Proceeding exactly as in Theorem 2, we prove that $g(X)$ is a separable subset of (E, \mathcal{T}_0) . Let $F = (E, \mathcal{T}_0)'$ and $E_0 =$ the closed separable subspace, in (E, \mathcal{T}_0) , generated by a countable dense subset of $(g(X), \mathcal{T}_0)$. If $g(x_0) \notin E_0$ for some $x_0 \in X$ there exists, by separation theorem, an $h \in E'$ such that $h \circ g(x_0) > 0$ and $h \equiv 0$ on E_0 . Since $E' = \bigcup_{n=1}^{\infty} A_n \subset F$, $h \circ g(x_0) \leq \sup(h \circ g(X)) \leq \sup h(E_0) = 0$, a contradiction. This proves the result.

In the next theorem we do not assume H to be uniformly bounded ([8], Theorem 4, p. 203).

THEOREM 6. *Let H be a pointwise bounded subset of $C(X)$. If H is equicontinuous then, with the topology of pointwise convergence on X , its closure in $C(X)$ is compact and metrizable. Conversely if H is sequentially compact then there is a μ -null set A such that H is equicontinuous at each point of the open set $X \setminus A$ of (X, τ_ρ) .*

Proof. If H is equicontinuous then its pointwise closed convex hull H_0 , in R^X , lies in $C(X)$ and is compact and convex, and so the result follows from Theorem 1.

Conversely suppose H is sequentially compact. Then, by Theorem 1, H is compact and metrizable. By the generalized Egoroff's theorem ([4], p. 198) there exists a \mathfrak{U} -partition of $X = \bigcup_{i=0}^{\infty} X_i$, with $\mu(X_0) = 0$ and $\mu(X_i) > 0$, $\forall i \geq 1$ such that $H|_{X_i}$ is compact in the topology of uniform convergence on X_i , $\forall i \geq 1$.

$Y_i = X_i \cap \rho(X_i)$, $i \geq 1$, are nonvoid, disjoint, open subsets of (X, τ_ρ) and $\mu(A) = 0$, where $A = X \setminus \bigcup_{i=1}^{\infty} Y_i$. By the Ascoli Theorem ([1], Ch. X, §2.5), $H|_{Y_i}$ are equicontinuous for each i . The result follows now.

REFERENCES

1. N. Bourbaki, *General Topology*, Hermann, Paris, 1966. (Translation by Addison-Wesley, Massachusetts.)
2. N. Dinculeanu, *Vector Measures*, Pergamon Press, New York, 1967.
3. S. S. Khurana, *Dunford-Pettis property*, J. Math. Anal. Appl., **65** (1978), 361-364.
4. P. A. Meyer, *Représentation intégral des fonctions excessives*, Résultats de Mokobodzki, Deminaire de Probabilités V, Univ. de Strasbourg, Springer-Verlag, Berlin-New York, Lecture Notes, **198** (1971), 196-208.
5. R. S. Phillips, *On weakly compact sets of a Banach space*, Amer. J. Math., **65** (1943), 108-136.
6. H. H. Schaefer, *Topological Vector Spaces*, Springer-Verlag, 1971.
7. A. Ionescu Tulcea, *Topologies compatible with liftings*, Bull. de la Math. Grèce, **8** (1967), 116-126.
8. ———, *On pointwise convergence, compactness and equicontinuity in the lifting topology I*, Zeit. Wahrscheinlichkeitstheorie und Verw. Gebiete, **26** (1973), 197-205.
9. ———, *On pointwise convergence, compactness and equicontinuity II*, Adv. Math., **12** (1974), 171-177.
10. A. Ionescu Tulcea, C. Ionescu Tulcea, *Topics in the Theory of Liftings*, Springer-Verlag, New York, 1969.
11. L. Schwartz, *Certaines propriétés des mesures sur les espaces de Banach*, Sem. Maurey-Schwartz, 1975-76, Ecole Polytechnique, No. 23.

Received August 12, 1978.

THE UNIVERSITY OF IOWA
IOWA CITY, IA 52242

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$84.00 a year (6 Vols., 12 issues). Special rate: \$42.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).

8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1979 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Patrick Robert Ahern, <i>On a theorem of Hayman concerning the derivative of a function of bounded characteristic</i>	297
Walter Allegretto, <i>Finiteness of lower spectra of a class of higher order elliptic operators</i>	303
Leonard Asimow, <i>Superharmonic interpolation in subspaces of $C_c(X)$</i>	311
Steven F. Bellenot, <i>An anti-open mapping theorem for Fréchet spaces</i>	325
B. J. Day, <i>Locale geometry</i>	333
John Erik Fornæss and Steven Krantz, <i>Continuously varying peaking functions</i>	341
Joseph Leonide Gerver, <i>Long walks in the plane with few collinear points</i>	349
Joseph Leonide Gerver and Lawrence Thom Ramsey, <i>On certain sequences of lattice points</i>	357
John R. Graef, Yuichi Kitamura, Takaši Kusano, Hiroshi Onose and Paul Winton Spikes, <i>On the nonoscillation of perturbed functional-differential equations</i>	365
James A. Huckaba and James M. Keller, <i>Annihilation of ideals in commutative rings</i>	375
Anzelm Iwanik, <i>Norm attaining operators on Lebesgue spaces</i>	381
Surjit Singh Khurana, <i>Pointwise compactness and measurability</i>	387
Charles Philip Lanski, <i>Commutation with skew elements in rings with involution</i>	393
Hugh Bardeen Maynard, <i>A Radon-Nikodým theorem for finitely additive bounded measures</i>	401
Kevin Mor McCrimmon, <i>Peirce ideals in Jordan triple systems</i>	415
Sam Bernard Nadler, Jr., Joseph E. Quinn and N. Stavrakas, <i>Hyperspaces of compact convex sets</i>	441
Ken Nakamura, <i>An explicit formula for the fundamental units of a real pure sextic number field and its Galois closure</i>	463
Vassili Nestoridis, <i>Inner functions invariant connected components</i>	473
Vladimir I. Olikier, <i>On compact submanifolds with nondegenerate parallel normal vector fields</i>	481
Lex Gerard Oversteegen, <i>Fans and embeddings in the plane</i>	495
Shlomo Reisner, <i>On Banach spaces having the property G.L.</i>	505
Gideon Schechtman, <i>A tree-like Tsirelson space</i>	523
Helga Schirmer, <i>Fix-finite homotopies</i>	531
Jeffrey D. Vaaler, <i>A geometric inequality with applications to linear forms</i>	543
William Jennings Wickless, <i>T as an \mathcal{Q} submodule of G</i>	555
Kenneth S. Williams, <i>The class number of $Q(\sqrt{-p})$ modulo 4, for $p \equiv 3 \pmod{4}$ a prime</i>	565
James Chin-Sze Wong, <i>On topological analogues of left thick subsets in semigroups</i>	571