

Pacific Journal of Mathematics

**ASYMPTOTIC STABILITY AND NONEXISTENCE OF GLOBAL
SOLUTION FOR A SEMILINEAR PARABOLIC EQUATION**

CHIA-VEN PAO

ASYMPTOTIC STABILITY AND NON-EXISTENCE OF GLOBAL SOLUTION FOR A SEMI-LINEAR PARABOLIC EQUATION

C. V. PAO

The aim of this paper is to investigate the existence and nonexistence of a nonnegative global solution for a semilinear parabolic system in a bounded domain. It is shown that for a certain class of initial functions the corresponding solution of the initial boundary value problem has a finite escape time, while for another class of initial functions a unique solution exists for all time and diminishes to zero. This result leads to an explicit estimate for the stability and the instability regions of the trivial steady-state solution. In the case of nonexistence of global solutions, an estimate for the finite escape time is also given.

I. Introduction. The problem of nonexistence of global solution for the semilinear parabolic equation

$$(1.1) \quad u_t - \nabla \cdot (D(x)\nabla u) = \alpha u^{1+\alpha} \quad (t \in (0, T], x \in \Omega)$$

has been given considerable attention in recent years (cf. [4-6, 13, 15]). In most of the above references the spatial domain Ω was taken as the whole space R^n so that the system under consideration by these authors is a Cauchy problem. In this paper, we consider a bounded domain Ω in R^n together with the following boundary and initial conditions:

$$(1.2) \quad \beta \frac{\partial u}{\partial \nu} + u = 0 \quad (t \in (0, T], x \in \partial\Omega)$$

$$(1.3) \quad u(0, x) = u_0(x) \quad (x \in \Omega).$$

In the above system, ∇ is the gradient operator, D is a positive function on $\bar{\Omega} \equiv \Omega \cup \partial\Omega$, α, β are constants with $\alpha > 0, \beta \geq 0$, $\partial\Omega$ is the boundary of Ω and ν is the outward normal vector on $\partial\Omega$. This system arises from models of simultaneous diffusion and recombination of electrons and ions as well as models of reactor dynamics where positive feedback is allowed when $a > 0$ (cf. [1, 8, 15]). We assume that Ω is sufficiently smooth, D is continuously differentiable in $\bar{\Omega}$ and u_0 is continuous nonnegative on $\bar{\Omega}$ and satisfies the boundary condition (1.2). The constant, a , may assume positive or negative values and its magnitude plays an important role in relation to the stability region of the trivial steady state solution.

The nonexistence of global solution to the Cauchy problem (1.1) and (1.3) for the case $a = D = 1$ in the whole space R^n has been discussed by Fujita [4, 5], Hayakawa [6], Sugitani [16] and more recently by Portnoy [13]. A similar problem for a bounded domain Ω has been investigated by Kaplan [7], Friedman [2] and Pao [11]. (See also Levine and Payne [9].) The purpose of this paper is to investigate the existence and nonexistence of a nonnegative global solution to the system (1.1)-(1.3) and to give a threshold result on the asymptotic behavior of the solution. Specifically, we show that for a certain class of initial functions u_0 a unique nonnegative solution $u(t, x)$ to (1.1)-(1.3) exists on some finite interval $[0, T_0)$ and

$$\lim(\max_x u(t, x)) = \infty \quad \text{as } t \longrightarrow T_0;$$

and for another class of initial functions, the solution u exists on $[0, \infty)$ and $\lim u(t, x) = 0$ on $\bar{\Omega}$ as $t \rightarrow \infty$. Moreover, we establish in the first case an explicit upper bound for the finite escape time T_0 while in the second case we give an estimate for the stability region as well as the instability region of the zero solution. We also show that when $a \leq 0$ the nonnegative solution u diminishes to zero for every nonnegative initial function u_0 . Thus if we increase the value of a from negative to positive then the zero steady state solution changes from global asymptotic stability to regional asymptotic stability; and for initial functions outside certain region the corresponding solutions diverge to infinite in finite time. As is to be expected our estimate demonstrates that larger initial function u_0 requires less time for the blowing-up property of the solution. Hence the consideration of a bounded domain instead of the whole space R^n leads to more delicate asymptotic behavior of the solution. It is interesting to note that this qualitative behavior also has a simple and natural physical interpretation and the results can be extended to more general systems (see the remarks in §2). In the present paper, however, we limit our discussion to the system (1.1)-(1.3).

2. The main results. In order to obtain our main results on the existence and nonexistence of a global solution we as in [10, 11] use the notion of upper and lower solutions for parabolic systems. By an upper solution we mean a smooth function $\tilde{u}(t, x)$ satisfying the inequalities

$$(2.1) \quad \begin{aligned} \tilde{u}_t - \nabla \cdot (D(x)\nabla \tilde{u}) &\geq a\tilde{u}^{1+\alpha} & (t \in (0, T], x \in \Omega) \\ \beta \frac{\partial \tilde{u}}{\partial \nu} + \tilde{u} &\geq 0 & (t \in (0, T], x \in \partial\Omega), \\ \tilde{u}(0, x) &\geq u_0(x) & (x \in \Omega) \end{aligned}$$

and a lower solution is a smooth function $\mathbf{u}(t, x)$ which satisfies all the reversed inequalities in (2.1). Moreover we require that $\mathbf{u} \leq \tilde{\mathbf{u}}$ on $[0, T] \times \bar{\Omega}$. Here by a smooth function it meant a continuous function w on $[0, T] \times \bar{\Omega}$ which is continuously differentiable in $t \in (0, T]$, twice continuously differentiable in $x \in \Omega$ and $\partial u / \partial \nu$ exists on $(0, T] \times \partial \Omega$.

Suppose that upper and lower solutions $\mathbf{u}, \tilde{\mathbf{u}}$ exist and $\mathbf{u} \leq \tilde{\mathbf{u}}$. Then by starting from the initial iteration $\mathbf{u}^{(0)} \equiv \tilde{\mathbf{u}}$ and $\mathbf{u}^{(0)} \equiv \mathbf{u}$, respectively, we can construct two sequences from the recurrence relation

$$(2.2) \quad \begin{aligned} \mathbf{u}_i^{(k)} - \mathcal{F}(D(x)\mathcal{F}\mathbf{u}^{(k)}) &= a(\mathbf{u}^{(k-1)})^{1+\alpha} & (t \in (0, T], x \in \Omega) \\ \beta \frac{\partial \mathbf{u}^{(k)}}{\partial \nu} + \mathbf{u}^{(k)} &= 0 & (t \in (0, T], x \in \partial \Omega) \\ \mathbf{u}^{(k)}(0, x) &= u_0(x) & (x \in \Omega) \end{aligned} \quad k = 1, 2, \dots$$

Denote these two sequences, respectively, by $\{\bar{\mathbf{u}}^{(k)}\}$ and $\{\underline{\mathbf{u}}^{(k)}\}$ (so that $\bar{\mathbf{u}}^{(0)} = \tilde{\mathbf{u}}, \underline{\mathbf{u}}^{(0)} = \mathbf{u}$). Then by the properties of upper and lower solutions one can easily show that (i) $\bar{\mathbf{u}}^{(k+1)} \leq \bar{\mathbf{u}}^{(k)}$, (ii) $\underline{\mathbf{u}}^{(k)} \leq \underline{\mathbf{u}}^{(k+1)}$ and (iii) $\underline{\mathbf{u}}^{(k)} \leq \bar{\mathbf{u}}^{(k)}$ for every $k = 0, 1, 2, \dots$. These inequalities imply that the pointwise limits

$$\lim \bar{\mathbf{u}}^{(k)} = \bar{\mathbf{u}} \quad \text{and} \quad \lim \underline{\mathbf{u}}^{(k)} = \underline{\mathbf{u}} \quad \text{as} \quad k \longrightarrow \infty$$

exist and satisfy the relation $\mathbf{u} \leq \underline{\mathbf{u}} \leq \bar{\mathbf{u}} \leq \tilde{\mathbf{u}}$ on $[0, T] \times \bar{\Omega}$. In fact, $\underline{\mathbf{u}}$ coincides with $\bar{\mathbf{u}}$ and is the unique solution of (1.1)-(1.3) (cf. [10, 12]). Therefore if we can find a pair of upper and lower solution with $\mathbf{u} \leq \tilde{\mathbf{u}}$, then a unique solution $u(t, x)$ to (1.1)-(1.3) exists and

$$(2.3) \quad \mathbf{u}(t, x) \leq u(t, x) \leq \tilde{\mathbf{u}}(t, x) \quad \text{on} \quad [0, T] \times \bar{\Omega}.$$

Since for $u_0 \geq 0$ the function $\mathbf{u} = 0$ satisfies all the reversed inequalities in (2.1), the existence of a nonnegative solution u satisfying $0 \leq u(t, x) \leq \tilde{\mathbf{u}}(t, x)$ is insured if one can find an upper solution $\tilde{\mathbf{u}}$ which is nonnegative. We shall do this for our global existence problem.

In the proof of the theorems the construction of $\tilde{\mathbf{u}}$ is based on the linear eigenvalue problem

$$(2.4) \quad \begin{aligned} \mathcal{F} \cdot (D(x)\mathcal{F}\phi) + \lambda\phi &= 0 & (x \in \Omega) \\ \beta \frac{\partial \phi}{\partial \nu} + \phi &= 0 & (x \in \partial \bar{\Omega}). \end{aligned}$$

It is well-known that the least eigenvalue λ_0 is positive and its corresponding eigenfunction $\psi_1(x)$ is positive in Ω . In fact, if $\beta > 0$ the

maximum principle implies that $\psi(x) > 0$ on $\bar{\Omega}$. We normalize ψ so that $\max \psi(x) = 1$. For convenience we set $\psi_m = \min \psi(x)$. Notice that $\psi_m > 0$ when $\beta > 0$. Our main results are stated in the following two theorems.

THEOREM 1. *Let $\beta > 0$, $a > 0$ and $\eta \equiv (\lambda_0/a)^{1/\alpha} \psi_m^{-1}$. Then for any $u_0(x) \geq b\psi(x)$ with $b > \eta$ there exists a constant $T_0 < \infty$ such that a unique solution $u(t, x)$ to (1.1)-(1.3) exists on $[0, T_0) \times \bar{\Omega}$ and satisfies*

$$(2.5) \quad \lim_{t \rightarrow T_0} \max_{x \in \bar{\Omega}} u(t, x) = \infty .$$

Moreover,

$$(2.6) \quad T_0 \leq (\alpha \lambda_0)^{-1} \ln [a(b\psi_m)^\alpha / (a(b\psi_m)^\alpha - \lambda_0)] .$$

THEOREM 2. *Let $\lambda \in (0, \lambda_0)$ and $\rho_\lambda = ((\lambda_0 - \lambda)/a)^{1/\alpha}$ when $a > 0$. Then for any $u_0(x) \leq \rho_\lambda \psi(x)$, a unique nonnegative solution $u(t, x)$ to (1.1)-(1.3) exists on $[0, \infty) \times \bar{\Omega}$ and satisfies*

$$(2.7) \quad 0 \leq u(t, x) \leq \rho_\lambda e^{-\lambda t} \psi(x) \quad (t > 0, x \in \bar{\Omega}) .$$

In the case of $a \leq 0$ the above inequality holds for every $\rho_\lambda \equiv \rho < \infty$ whenever $u_0(x) \leq \rho \psi(x)$.

REMARK. (a) The results in Theorems 1 and 2 show that when $a > 0$ a stability region of the zero solution is given by the set $A_1 \equiv \{u_0; 0 \leq u_0 \leq \rho_\lambda \psi\}$ while an instability region is $A_2 \equiv \{u_0; u_0 \geq b\psi\}$. By choosing λ sufficiently small the value of ρ_λ may be taken arbitrarily close to $(\lambda_0/a)^{1/\alpha}$. Since λ_0 is directly related to the size (for a fixed geometry) of the domain Ω it follows that smaller domain tends to stabilize the equilibrium solution $u = 0$ while large domain tends to destabilize the zero solution. This is, of course, to be expected physically since energy leaks faster in smaller domains than larger ones.

(b) Theorems 1 and 2 remain true when the operator $\nabla \cdot (D(x)\nabla u)$ is replaced by a more general uniformly elliptic operator in the form

$$Lu \equiv \sum_{i,j=1}^n a_{ij}(x) u_{x_i x_j} + \sum_{i=1}^n b_i(x) u_{x_i} .$$

In this situation the least eigenvalue λ_0 and its corresponding eigenfunction ψ should be with respect to the operator L (under the same boundary condition). Notice that $\lambda_0 > 0$ (or at least there exists a eigenvalue λ_0 which is real positive) and the corresponding eigenfunction ψ is positive (cf. [14]).

The results in Theorems 1 and 2 can be given a physical inter-

pretation such as in the process of diffusion or heat conduction. When the spatial domain is bounded, energy (or heat) leaks through the boundary surface $\partial\Omega$. Thus for the class of initial functions $u_0 \leq \rho_\lambda \psi$, energy leaks away through the boundary surface before it gets large so that the effect due to the feedback term $au^{1+\alpha}$ (which acts as a source for $a > 0$) is diminishing. But when the initial function u_0 is large to the extent of $u_0 > \eta\psi$, the feedback term dominates the leakage initially and its effect tends to grow and eventually leads to the unboundedness of the solution in finite time. It is interesting to note that our estimate indicates that the finite escape time T_0 is larger for smaller values of a and becomes smaller as a gets larger. In fact, if we increase the value of a from $a \leq 0$ to $a > 0$ then the system (1.1)-(1.3) changes from global asymptotic stability to regional asymptotic stability, and as a increases, the stability region gets smaller while the instability region becomes larger.

It is to be noted that there remains a gap between the stability region A_1 and the instability region A_2 (by a factor of ψ_m^{-1}). This gap can be closed if one can show that for u_0 not in these two regions the corresponding solution lies either in A_1 or in A_2 after a finite time.

3. Proofs of the main theorems.

Proof of Theorem 1. We first construct a function u satisfying the reversed inequalities in (2.1). Let $u = e^{-\lambda_0 t} p(t) \psi(x)$ with $p(0) \leq b$, where $p(t)$ is a continuously differentiable function to be chosen. Since $u(0, x) = p(0) \psi(x) \leq u_0(x)$ and by (2.4) $\Delta \cdot (D\nabla u) = -\lambda_0 u$ and $\beta \partial u / \partial \nu + u = 0$ we see that u is a lower solution if

$$(3.1) \quad e^{-\lambda_0 t} P'(t) \psi(x) \leq \alpha (e^{-\lambda_0 t} P(t) \psi(x))^{1+\alpha},$$

where $P' \equiv dP/dt$. Clearly, the above inequality holds if

$$(3.2) \quad P' \leq \alpha \psi_m^\alpha e^{-\alpha \lambda_0 t} P^{1+\alpha}.$$

It is easily seen that the function $P(t)$ given by

$$(3.3) \quad P(t) = [b^{-\alpha} + \alpha \lambda_0^{-1} \psi_m^\alpha (1 - e^{-\alpha \lambda_0 t})]^{-1/\alpha}, \quad (t \in [0, T_1])$$

satisfies the requirement in (3.2) for $t \in [0, T_1)$, where T_1 is given by the right side of (2.6). Hence the function $u = e^{-\lambda_0 t} P(t) \psi(x)$ with $P(t)$ given by (3.3) is a lower solution on $[0, T_2] \times \bar{\Omega}$ for every $T_2 < T_1$. Using this function as the initial iteration in (2.2) with $T < T_1$ we obtain a monotone nondecreasing sequence which converges to a solution u of (1.1)-(1.3) on $[0, T] \times \bar{\Omega}$ provided that this sequence is

uniformly bounded from above. However, this may not be the case for every $T < T_1$ unless u has the finite escape time exactly at T_1 . This is due to the fact that u is a lower bound of u when the solution exists. To overcome this difficulty we take an arbitrarily large constant $M > u_0(x)$ and define a function $f(u)$ by

$$(3.4) \quad f(u) = au^{1+\alpha} \text{ when } u \leq M \text{ and } f(u) = aM^{1+\alpha} \text{ when } u > M .$$

Choose T_2 sufficiently close to T_1 such that $u(T_2, x) \geq M$ for some $x \in \bar{\Omega}$. Then by replacing the right-side of the first equation in (2.2) by $f(u^{(k-1)})$ and using the initial iteration $u^{(0)} = u$ with $T = T_2$ we again obtain a monotone nondecreasing sequence for the "modified" system (1.1)-(1.3) (i.e., with $au^{1+\alpha}$ replaced by $f(u)$). We denote this sequence also by $\{u^{(k)}\}$. Now since $f(u^{(k)})$ is uniformly bounded, an application of the well-known estimate for linear parabolic systems implies that $\{u^{(k)}\}$ is bounded (e.g., see [3] p. 146). It follows from the monotone property that $\{u^{(k)}\}$ converges to a unique solution $u^*(t, x)$ of the modified system (1.1)-(1.3) and

$$(3.5) \quad u^*(t, x) \geq u(t, x) \equiv e^{-\lambda_0 t} P(t) \psi(x) \quad (t \in [0, T_2] \times \bar{\Omega}) .$$

But $u(T_2, x) \geq M$ for some $x \in \bar{\Omega}$ and $u_0(x) < M$, there exists $T_3 \leq T_2$ such that $u^*(t, x) \leq M$ for $(t, x) \in [0, T_3] \times \bar{\Omega}$ and $u^*(T_3, x) = M$ for some $x \in \bar{\Omega}$. By the definition of $f(u)$, u^* is the solution of (1.1)-(1.3) with $T = T_3$. Since M can be chosen arbitrarily large the above conclusion implies that the solution of (1.1)-(1.3) must be unbounded on $[0, T_0] \times \bar{\Omega}$ for some $T_0 \leq T_1$. For if it were bounded (say, by K) on $[0, T] \times \bar{\Omega}$ for every $T \leq T_1$, then by choosing $M > K$ in the definition of $f(u)$ we obtain a solution u^* of (1.1)-(1.3) such that $u^*(t, x) = M$ at some point $(t, x) \in [0, T_3] \times \bar{\Omega}$ where $T_3 < T_1$. Clearly this is impossible. The proof of the theorem is completed.

Proof of Theorem 2. For the case $a > 0$ it suffices to show that $\tilde{u} \equiv \rho_\lambda e^{-\lambda t} \psi(x)$ is an upper solution on $[0, T] \times \bar{\Omega}$ for every finite T . In view of (2.1) and since the last two conditions are satisfied, this will follow if

$$(3.6) \quad (\lambda_0 - \lambda) \rho_\lambda e^{-\lambda t} \psi(x) \geq a(\rho_\lambda e^{-\lambda t} \psi(x))^{1+\alpha} .$$

The above inequality is clearly satisfied for any ρ_λ satisfying $\rho_\lambda^\alpha \leq (\lambda_0 - \lambda)/a$. This proves (2.7). When $a \leq 0$, the sequence obtained from (2.2) with $u^{(0)} = \tilde{u} \equiv \rho e^{-\lambda t} \psi(x)$ is no longer monotone nonincreasing and thus we need to modify our construction. To achieve this, we replace the first equation in (2.2) by

$$(3.7) \quad \begin{aligned} u_t^{(k)} - \mathcal{V} \cdot (D(x) \mathcal{V} u^{(k)}) + Qu^{(k)} \\ = a(u^{(k-1)})^{1+\alpha} + Qu^{(k-1)} , \quad (t \in (0, T], x \in \Omega) \end{aligned}$$

where Q is a constant satisfying $Q \geq -a\rho^\alpha$ while $T < \infty$ is arbitrary. It is easily seen by using the initial iteration $u^{(0)} = \rho e^{-\lambda t} \psi(x)$ that the sequence constructed from (3.7) together with the last two equations in (2.2) is nonnegative and is monotone nonincreasing (cf. [10]). Therefore it converges to a unique solution $u(t, x)$ of the system (1.1)-(1.3) such that $0 \leq (t, x) \leq \rho e^{-\lambda t} \psi(x)$ on $[0, T] \times \bar{D}$. This proves the theorem.

REFERENCES

1. C. Y. Chan, *Existence, uniqueness, upper and lower bounds of solutions of nonlinear space-time nuclear reactor kinetics*, SIAM J. Appl. Math., **27** (1974), 72-82.
2. A. Friedman, *Remarks on nonlinear parabolic equations*, Proc. Amer. Math. Soc.; Symp. Appl. Math., **17** (1965), 3-23.
3. ———, *Partial Differential Equations of Parabolic Type*, Prentice Hall, Englewood Cliffs, N.J., 1964.
4. H. Fujita, *On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$* , J. Fac. Sci. Univ. Tokyo Sect. I, **13** (1966), 109-124.
5. ———, *On some nonexistence and nonuniqueness theorems for nonlinear parabolic equations*, Proc. Symp. Pure Math. Vol. XVIII, Part 1, Amer. Math. Soc., (1970), 105-113.
6. K. Hayakawa, *On nonexistence of global solutions of some semi-linear parabolic differential equations*, Proc. Japan Acad., **49** (1973), 503-505.
7. S. Kaplan, *On the growth of solutions of quasi-linear parabolic equations*, Comm. Pure Appl. Math., **16** (1963), 305-330.
8. W. E. Kastenberg and P. L. Chambré, *On the stability of nonlinear space-dependent reactor kinetics*, Nucl. Sci. Engrg., **31** (1968), 67-79.
9. H. A. Levine and L. E. Payne, *Nonexistence of global weak solutions for classes of nonlinear wave and parabolic equations*, J. Math. Anal. App., **55** (1976), 329-334.
10. C. V. Pao, *Positive solutions of a nonlinear boundary value problem of parabolic type*, J. Differential Equations, **22** (1967), 145-163.
11. ———, *Nonexistence of global solutions and bifurcation analysis for a boundary-value problem of parabolic type*, Proc. Amer. Math. Soc., (to appear).
12. ———, *Successive approximations of some nonlinear initial boundary value problems*, SIAM J. Math. Anal., **46** (1974), 91-102.
13. S. L. Portnoy, *On solutions to $u_t = \Delta u + u^2$ in two dimensions*, J. Math. Anal. Appl., **55** (1976), 291-294.
14. M. H. Protter and H. F. Weinberger, *On the spectrum of general second order operators*, Bull. Amer. Math. Soc., **72** (1966), 251-255.
15. G. Rosen, *Solutions to the nonlinear recombination equation for the infinite spatial domain*, SIAM J. Appl. Math., **29** (1975), 146-151.
16. S. Sugitani, *On non-existence of global solutions for some nonlinear integral equations*, Osaka J. Math., **12** (1975), 45-51.

Received March 4, 1977.

NORTH CAROLINA STATE UNIVERSITY
 RALEIGH, NC 27650
 AND
 UNIVERSITÄT GRAZ
 GRAZ, AUSTRIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Michael James Beeson, <i>Goodman's theorem and beyond</i>	1
Robert S. Cahn and Michael E. Taylor, <i>Asymptotic behavior of multiplicities of representations of compact groups</i>	17
Douglas Michael Campbell and Vikramaditya Singh, <i>Valence properties of the solution of a differential equation</i>	29
J.-F. Colombeau, Reinhold Meise and Bernard Perrot, <i>A density result in spaces of Silva holomorphic mappings</i>	35
Marcel Ern�, <i>On the relativization of chain topologies</i>	43
Le Baron O. Ferguson, <i>Uniform and L_p approximation for generalized integral polynomials</i>	53
Kenneth R. Goodearl and David E. Handelman, <i>Homogenization of regular rings of bounded index</i>	63
Friedrich Haslinger, <i>A dual relationship between generalized Abel-Gon�arov bases and certain Pincherle bases</i>	79
Miriam Hausman, <i>Generalization of a theorem of Landau</i>	91
Makoto Hayashi, <i>2-factorization in finite groups</i>	97
Robert Marcus, <i>Stochastic diffusion on an unbounded domain</i>	143
Isabel Dotti de Miatello, <i>Extension of actions on Stiefel manifolds</i>	155
C. David (Carl) Minda, <i>The hyperbolic metric and coverings of Riemann surfaces</i>	171
Somashekhar Amrith Nainpally and Mohan Lal Tikoo, <i>On T_1-compactifications</i>	183
Chia-Ven Pao, <i>Asymptotic stability and nonexistence of global solution for a semilinear parabolic equation</i>	191
Shigeo Segawa, <i>Harmonic majoration of quasibounded type</i>	199
Sze-Kai Tsui and Steve Wright, <i>The splitting of operator algebras</i>	201
Bruce Williams, <i>Hopf invariants, localization and embeddings of Poincar� complexes</i>	217
Leslie Wilson, <i>Nonopenness of the set of Thom-Boardman maps</i>	225
Alicia B. Winslow, <i>There are 2^c nonhomeomorphic continua in $\beta R^n - R^n$</i>	233