

Pacific Journal of Mathematics

UNBOUNDED MULTIPLIERS ON COMMUTATIVE BANACH ALGEBRAS

JAMES ALAN WOOD

UNBOUNDED MULTIPLIERS ON COMMUTATIVE BANACH ALGEBRAS

JAMES WOOD

In this paper the notion of an unbounded multiplier on a commutative Banach algebra is introduced. It is proven that, as in the case of bounded multipliers, unbounded multipliers also have Gelfand transforms. Some of the properties of these transforms are then developed. The final result of the paper is a new characterization of the bounded multipliers on A where A is a Banach algebra of the type described below.

1. Introduction. In general, by a multiplier on a commutative Banach algebra A one means a bounded linear operator $T: A \rightarrow A$ such that $T(xy) = xT(y)$ for all $x, y \in A$. There is an extensive literature on the subject. One can consult, for example, [2] and [3]. There does not seem to be, however, a systematic treatment of unbounded multipliers although such multipliers occur quite naturally. For example, consider the Banach algebra $L^1(-\infty, \infty)$ with the convolution product and define T by $T(f) = f'$, where the domain $\mathcal{D}(T)$ of T is the set $\{f \mid f \in L^1(-\infty, \infty) \text{ and } f \text{ is absolutely continuous}\}$. It is easy to check that $T(fg) = fT(g)$ almost everywhere for all $f \in L^1(-\infty, \infty)$ and for all $g \in \mathcal{D}(T)$. However, T is not bounded. (Although it is closed.)

As another example, take the Banach algebra $C_0(-\infty, \infty)$ of all complex valued continuous functions on the real line which vanish at infinity with T defined by $T(f)(x) = xf(x)$. The domain $\mathcal{D}(T)$ can be taken to be the set of all functions in $C_0(-\infty, \infty)$ with compact support. Finally, let $\{T_t \mid t \geq 0\}$ be a semi-group of class C_0 of bounded multipliers on the Banach algebra A . Then the infinitesimal generator T_0 of $\{T_t \mid t \geq 0\}$ is an (in general) unbounded multiplier, for if $x \in \mathcal{D}(T_0)$ and $y \in A$ then

$$\begin{aligned} T_0(xy) &= \lim_{h \downarrow 0} \frac{1}{h} [T(xy) - xy] \\ &= y \lim_{h \downarrow 0} \frac{1}{h} [T(x) - x] = yT_0(x). \end{aligned}$$

It is the purpose of this present paper to study some of the properties of unbounded multipliers in the general setting of a commutative Banach algebra. In particular, henceforth A always denotes a regular, commutative semi-simple Banach algebra. We also assume for the rest of the paper that A has a bounded

approximate identity $\{e_n\}$ and that the Gelfand transform of each $\{e_n\}$ has compact support. By $\Delta(A)$ we shall mean the maximal ideal space of A with the Gelfand topology and the symbol \hat{x} denotes the Gelfand transform of $x \in A$.

2. Definitions and general properties.

DEFINITION 2.1. Let A be a Banach algebra. The function T defined on the dense ideal $\mathcal{D}(T) \subseteq A$ is called a multiplier of A if and only if for all $x \in \mathcal{D}(T)$ and all $y \in A$, $T(xy) = yT(x)$.

The first theorem shows that, as in the bounded case, every multiplier has a Gelfand transform.

THEOREM 2.1. *Given a multiplier T on A there is a unique complex valued continuous function T^\wedge on $\Delta(A)$ such that for all $\tau \in \Delta(A)$ and all $x \in \mathcal{D}(T)$, $T(x)^\wedge(\tau) = T^\wedge(\tau)x^\wedge(\tau)$.*

Proof. First of all observe that for each $\tau \in \Delta(A)$ there is an $x \in \mathcal{D}(T)$ with $x^\wedge(\tau) \neq 0$. For otherwise, if for some τ_0 , $x^\wedge(\tau_0) = 0$ for all $x \in \mathcal{D}(T)$, then $\mathcal{D}(T)$ is contained in some maximal ideal which is a contradiction. Now if $\tau_0 \in \Delta(A)$, choose $x \in \mathcal{D}(T)$ so that $x^\wedge(\tau_0) \neq 0$ and define $T^\wedge(\tau_0)$ by $T^\wedge(\tau_0) = T(x)^\wedge(\tau_0)/x^\wedge(\tau_0)$. The function T^\wedge is independent of x , for if $y \in \mathcal{D}(T)$ and $y^\wedge(\tau_0) \neq 0$, then from $T(xy) = xT(y) = yT(x)$ follows $x^\wedge(\tau)T(y)^\wedge(\tau) = y^\wedge(\tau)T(x)^\wedge(\tau)$ for all $\tau \in \Delta(A)$, so that $T(y)^\wedge(\tau_0)/y^\wedge(\tau_0) = T(x)^\wedge(\tau_0)/x^\wedge(\tau_0)$. It is clear that T^\wedge is continuous. To show uniqueness suppose there is a continuous function ϕ on $\Delta(A)$ such that $T(x)^\wedge(\tau) = \phi(\tau)x^\wedge(\tau)$ for all $x \in \mathcal{D}(T)$ and all $\tau \in \Delta(A)$. Then $T^\wedge(\tau)x^\wedge(\tau) = \phi(\tau)x^\wedge(\tau)$ for all $x \in \mathcal{D}(T)$ and $\tau \in \Delta(A)$ and since for each τ we can find an x such that $x^\wedge(\tau) \neq 0$, it follows that $T^\wedge = \phi$.

COROLLARY 2.1. *A multiplier is linear on its domain.*

Proof. Suppose $x, y \in \mathcal{D}(T)$ and let $z = \alpha x + \beta y$ where α and β are complex numbers. Then from $z^\wedge(\tau) = \alpha x^\wedge(\tau) + \beta y^\wedge(\tau)$ we get successively

$$T^\wedge(\tau)z^\wedge(\tau) = \alpha T^\wedge(\tau)x^\wedge(\tau) + \beta T^\wedge(\tau)y^\wedge(\tau)$$

$$T(z)^\wedge(\tau) = \alpha T(x)^\wedge(\tau) + \beta T(y)^\wedge(\tau)$$

$$T(z)^\wedge(\tau) = (\alpha T(x) + \beta T(y))^\wedge(\tau)$$

$$T(z) = \alpha T(x) + \beta T(y),$$

the last equation following from the semi-simplicity of the algebra.

THEOREM 2.2. *Every multiplier is closable.*

Proof. Let T be a multiplier and assume $\{x_n\}$ is a sequence in $\mathcal{D}(T)$ with $\lim x_n = 0$ and $\lim T(x_n) = y$. Since $T(x_n) \rightarrow y$, $T^\wedge(\tau)x_n^\wedge(\tau) \rightarrow y^\wedge(\tau)$. But $T^\wedge(\tau)x_n^\wedge(\tau) \rightarrow 0$ also and thus $y^\wedge(\tau) = 0$ for all $\tau \in \Delta(A)$ so that $y = 0$.

THEOREM 2.3. *If S and T are multipliers such that $S^\wedge(\tau) = T^\wedge(\tau)$ for all τ and in addition S and T are closed then $S = T$.*

Proof. The algebra has a bounded approximate identity $\{e_\alpha\}$. Now for each α and any $\varepsilon > 0$ there is an $e'_\alpha \in \mathcal{D}(S)$ such that $\|e_\alpha - e'_\alpha\| < \varepsilon$. Thus $\{e'_\alpha\}$ is an approximate identity in $\mathcal{D}(S)$. Similarly we can construct an approximate identity e''_α in $\mathcal{D}(T)$. Now letting $f_\alpha = e'_\alpha e''_\alpha$, we have that $f_\alpha \in \mathcal{D}(S) \cap \mathcal{D}(T)$ and since $S^\wedge(\tau) = T^\wedge(\tau)$ we get $S^\wedge(\tau)e'_\alpha(\tau)e''_\alpha(\tau) = T^\wedge(\tau)e'_\alpha(\tau)e''_\alpha(\tau)$ so that $S(f_\alpha) = T(f_\alpha)$. Now let $f \in \mathcal{D}(S)$. Then $ff_\alpha \rightarrow f$ and $S(ff_\alpha) = f_\alpha S(f) \rightarrow S(f)$. But $S(ff_\alpha) = fS(f_\alpha) = fT(f_\alpha)$ so that $T(ff_\alpha) \rightarrow S(f)$ also. Therefore, since T is closed, $f \in \mathcal{D}(T)$ and $T(f) = S(f)$. By reversing the roles of S and T we get the assertion of the theorem.

COROLLARY 2.2. *If S and T are multipliers such that $S^\wedge = T^\wedge$, then S and T have the same closure.*

Proof. If S_0 and T_0 denote the closures of S and T respectively, then $S_0^\wedge = S^\wedge$ and $T_0^\wedge = T^\wedge$. Therefore $S_0^\wedge = T_0^\wedge$ so that by the previous theorem $S_0 = T_0$.

The previous theorem and corollary provide us in an obvious way with a means to define an equivalence relation on the set of multipliers. We say that S is equivalent to T , $S \sim T$, if and only if $S^\wedge = T^\wedge$. Thus the set of multipliers can be split up into equivalence classes such that all multipliers in the same class have the same closure and the same transform.

The next theorem concerns inverses of multipliers.

THEOREM 2.4. *Let T be a multiplier. Then T^{-1} exists and is a multiplier if and only if T^\wedge never vanishes.*

Proof. Assume T^{-1} exists as a multiplier. Then for each $x \in \mathcal{D}(T^{-1})$ there is a $z \in \mathcal{D}(T)$ such that $T^\wedge(\tau)z^\wedge(\tau) = x^\wedge(\tau)$. If $T^\wedge(\tau_0) = 0$ for some τ_0 then $x^\wedge(\tau_0) = 0$ for all $x \in \mathcal{D}(T^{-1})$ which contradicts the fact that $\mathcal{D}(T^{-1})$ is dense. On the other hand assume that T^\wedge never vanishes. Then the set $I = \{y \in A \mid x \in \mathcal{D}(T) \text{ with } T^\wedge(\tau)x^\wedge(\tau) = y^\wedge(\tau)\}$ is an ideal in A . If I is not dense, then its

closure is a closed proper ideal and hence contained in a maximal ideal. Therefore, there is a τ_0 such that $y^\wedge(\tau_0) = 0$ for all $y \in I$ and this in turn implies $T^\wedge(\tau_0)x^\wedge(\tau_0) = 0$ for all $x \in \mathcal{D}(T)$. Thus $x^\wedge(\tau_0) = 0$ for all $x \in \mathcal{D}(T)$ which is a contradiction.

It is possible to characterize those functions on $\Delta(A)$ which are transforms of multipliers. This is done in the following theorem.

THEOREM 2.5. *In order that T^\wedge be the transform of a multiplier it is necessary and sufficient that T^\wedge belong locally to A^\wedge at each point of $\Delta(A)$.*

Proof. If T^\wedge belongs locally to A^\wedge at each point of $\Delta(A)$, then $T^\wedge e_\alpha^\wedge$ belongs locally to A^\wedge at each point $\Delta(A) \cup \{\infty\}$. By a theorem due to F. Birtel, ([1], p. 818), this implies that $T^\wedge e_\alpha^\wedge$ belongs to A^\wedge . Therefore, for any $x \in A$, $x^\wedge T^\wedge e_\alpha^\wedge$ belongs to A^\wedge and we define $T(xe_\alpha) = y_\alpha$ where $y_\alpha^\wedge = x^\wedge T^\wedge e_\alpha^\wedge$. If we then let $\mathcal{D}(T)$ be $\{xe_\alpha \mid x \in A\}$, we have $T(zxe_\alpha) = y_\alpha$ where $y_\alpha^\wedge = z^\wedge x^\wedge T^\wedge e_\alpha^\wedge$. But $x^\wedge T^\wedge e_\alpha^\wedge = T(xe_\alpha)^\wedge$ and so $y_\alpha = zT(xe_\alpha)$. Thus T determines a multiplier. Conversely, if T^\wedge is the transform of a multiplier and $p \in \Delta(A)$, there exists an $x \in \mathcal{D}(T)$ such that $x^\wedge(\tau) \neq 0$ on some compact neighborhood V of p . There is then a $y \in A$ such that $y^\wedge(\tau) = 1/x^\wedge(\tau)$ for all $\tau \in V$ so that $x^\wedge(\tau)y^\wedge(\tau) = 1$ on V . But $xy \in \mathcal{D}(T)$, so that if we let $z = T(xy)$, then $z^\wedge(\tau) = T^\wedge(\tau)x^\wedge(\tau)y^\wedge(\tau)$ for all $\tau \in \Delta(A)$ and T^\wedge belongs locally to A^\wedge at p .

COROLLARY 2.3. *If T^\wedge is the transform of a multiplier, then T^\wedge is the transform of a bounded multiplier if and only if $|p(T(x))|/|px| \leq M$ for all $p \in A^*$ and all $x \in \mathcal{D}(T)$. (Here A^* is the dual of A and px is the functional defined on A by $px(y) = p(xy)$ for all $y \in A$.)*

Proof. The proof follows immediately by combining the above theorem and the following theorem proved in [4].

THEOREM. *Let \hat{F} be a complex valued function on $\Delta(A)$. In order that \hat{F} determine a bounded multiplier of A , it is necessary and sufficient that \hat{F} belong locally to $\Delta(A)$ at each point of $\Delta(A)$ and that $|p(\hat{F}(x))|/|px| \leq M$, for all $p \in A^*$ and for all $x \in A_F$. (A_F is the set $\{x \mid \hat{F}\hat{x} = \hat{y}$ for some $y \in A\}$.)*

3. Further results. In Theorem 2.5 we gave a necessary and sufficient condition that a function on $\Delta(A)$ be the transform of a multiplier. The next theorem gives a different necessary and sufficient condition that a function be the transform of a multiplier.

This identity, as it turns out, allows us to characterize those functions which are transforms of bounded multipliers on the Banach algebra A .

THEOREM 3.1. *The function λ belongs locally to A^\wedge at each point of $\Delta(A)$ if and only if there is an approximate identity $\{e_\alpha\}$ whose Gelfand transforms have compact support such that for each α , $e_\alpha^\wedge(\tau)\lambda(\tau) = f_\alpha^\wedge(\tau)$ for some $f_\alpha \in A$.*

Proof. By referring to the proof of Theorem 2.5 one can easily see that if λ belongs locally to A^\wedge then $\lambda(\tau)e_\alpha^\wedge(\tau)$ belongs locally to A^\wedge .

Conversely, given any $\tau_0 \in \Delta(A)$ there exists a neighborhood V of τ_0 such that $e_{\alpha_0}^\wedge(\tau) \neq 0$ for some α_0 and all $\tau \in V$. Thus there is a $g_{\alpha_0}^\wedge$ such that $g_{\alpha_0}^\wedge(\tau) = 1/e_{\alpha_0}^\wedge(\tau)$ for all $\tau \in V$. But there is an f_{α_0} such that $e_{\alpha_0}^\wedge(\tau)\lambda(\tau) = f_{\alpha_0}^\wedge(\tau)$ for some τ . Hence, for $\tau \in V$ we have $\lambda(\tau) = f_{\alpha_0}^\wedge(\tau)g_{\alpha_0}^\wedge(\tau)$ so that λ belongs locally to A^\wedge .

In Corollary 2.3 necessary and sufficient conditions were given in order that λ be the transform of a bounded multiplier. However, for specific algebras the inequality in that corollary may be hard to check. The following theorem gives another necessary and sufficient condition that λ be the transform of a bounded multiplier.

THEOREM 3.2. *In order that the function λ on $\Delta(A)$ be the transform of a bounded multiplier it is necessary and sufficient that $\lambda e_\alpha^\wedge = g_\alpha^\wedge$ where $g_\alpha \in A$, and there exists $M > 0$ such that $\|g_\alpha\| \leq M$ for all α .*

Proof. The necessity is clear. Assume therefore, that $\lambda e_\alpha^\wedge = g_\alpha^\wedge$ where $g_\alpha \in A$ and $\|g_\alpha\| \leq M$ for all α . Define $T(x) = y$ where $y^\wedge = \lambda x^\wedge$ and $\mathcal{S}(T) = \{x \in A \mid \lambda x^\wedge \in A^\wedge\}$. It is easy to see that $\mathcal{S}(T)$ is an ideal and, since $\{e_\alpha\}$ is contained in $\mathcal{S}(T)$, it is dense. It follows directly from the definition that T is a multiplier. Now let $f \in \mathcal{S}(T)$. Then $T(e_\alpha)f = T(e_\alpha f) = T(f)e_\alpha$ and $T(f)e_\alpha \rightarrow T(f)$. Therefore $\|T(f)\| \leq \sup_\alpha \|T(e_\alpha)f\| \leq M\|f\|$.

REFERENCES

1. F. Birtel, *On a commutative extension of a Banach algebra*, Proc. Amer. Math. Soc., **13** (1962), p 818.
2. R. Larsen, *An Introduction to the Theory of Multipliers*, Grundlehren der mathematischen Wissenschaften, Springer-Verlag, 1969.
3. ———, *The Multiplier Problem*, Lecture Notes in Mathematics, Springer-Verlag, 1969.

4. J. A. Wood, *A necessary and sufficient condition that a function on the maximal ideal space of a Banach algebra be a multiplier*, Proc. Amer. Math. Soc., **66**, number 1, (1977), 39.

Received October 3, 1978 and in revised form June 5, 1979.

VIRGINIA COMMONWEALTH UNIVERSITY, RICHMOND, VA 23284

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$84.00 a year (6 Vols., 12 issues). Special rate: \$42.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1979 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Charles A. Akemann and Steve Wright, <i>Compact and weakly compact derivations of C^*-algebras</i>	253
Dwight Richard Bean, Andrzej Ehrenfeucht and George Frank McNulty, <i>Avoidable patterns in strings of symbols</i>	261
Richard Clark Brown, <i>Notes on generalized boundary value problems in Banach spaces. I. Adjoint and extension theory</i>	295
Kenneth Alexander Brown and John William Lawrence, <i>Injective hulls of group rings</i>	323
Jacob Burbea, <i>The Schwarzian derivative and the Poincaré metric</i>	345
Stefan Andrus Burr, <i>On the completeness of sequences of perturbed polynomial values</i>	355
Peter H. Chang, <i>On the characterizations of the breakdown points of quasilinear wave equations</i>	361
Joseph Nicholas Fadyn, <i>The projectivity of $\text{Ext}(T, A)$ as a module over $E(T)$</i>	383
Donald Eugene Maurer, <i>Arithmetic properties of the idèle discriminant</i>	393
Stuart Rankin, Clive Reis and Gabriel Thierrin, <i>Right subdirectly irreducible semigroups</i>	403
David Lee Rector, <i>Homotopy theory of rigid profinite spaces. I</i>	413
Raymond Moos Redheffer and Wolfgang V. Walter, <i>Comparison theorems for parabolic functional inequalities</i>	447
H. M. (Hari Mohan) Srivastava, <i>Some generalizations of Carlitz's theorem</i>	471
James Alan Wood, <i>Unbounded multipliers on commutative Banach algebras</i>	479
T. Yoshimoto, <i>Vector-valued ergodic theorems for operators satisfying norm conditions</i>	485
Jerry Searcy and B. Andreas Troesch, <i>Correction to: "A cyclic inequality and a related eigenvalue problem"</i>	501
Leslie Wilson, <i>Corrections to: "Nonopenness of the set of Thom-Boardman maps"</i>	501