

# Pacific Journal of Mathematics

## **THE SECOND LIE ALGEBRA COHOMOLOGY GROUP AND WEYL MODULES**

JOHN BRENDAN SULLIVAN

## THE SECOND LIE ALGEBRA COHOMOLOGY GROUP AND WEYL MODULES

JOHN BRENDAN SULLIVAN

*Dedicated to Gerhard Hochschild on the occasion of his 65th birthday*

**The Lie algebra 1-cohomology of classical Lie algebras in characteristic  $p$  is nonzero at some modules. In fact, Hochschild showed that the restricted 1-cohomology is nonzero. Here I will systematically produce Weyl modules at which the (unrestricted) 2-cohomology is nonzero. This will provide examples of nonsplit abelian extensions of Lie algebras by Weyl modules in characteristic  $p$ , where the Lie algebras are the reductions modulo  $p$  of integral Chevalley forms of complex semisimple Lie algebras.**

The method requires passages among cohomology groups for Lie algebras defined over the integers, rationals, and the prime field of characteristic  $p$ . In proving Theorem 1, we show that the 2-cohomology of an integral Lie algebra has nonzero  $p$ -torsion at a module whenever the reduced (modulo  $p$ ) algebra has nonzero 1-cohomology at the reduced module and the extension of the integral Lie algebra to the rationals has zero for its first and second cohomology at the extension of the module by the rationals. The tensor product of the integral cohomology group with the prime field is a nonzero piece of the cohomology group of the reduced Lie algebra. This will give the examples that we seek provided that we produce modules which satisfy the conditions given above for the groups of the reduced and rational algebras. The second condition is satisfied in the classical case since the rational Lie algebra will be semisimple. As for the first condition, in [4] I gave irreducible modules in characteristic  $p$  at which the 1-cohomology is nonzero. I will show that the cohomology is also nonzero at a Weyl module associated to the irreducible module. The module that we seek for the integral Lie algebra is one that reduces modulo  $p$  to the Weyl module. I have carried out this enterprise of showing that the cohomology is nonzero at the Weyl module only for the Lie algebra of the special linear group although I believe that it can be carried out generally.

In §3, I take a particular one cocycle  $g$  (constructed in [4]) and, working with an integral cochain which reduces to  $g$ , apply the coboundary operator to obtain explicitly a  $p$ -torsion element of the integral cohomology group.

I wish to thank Professor Hochschild for his kindness and generosity over the years.

NOTATION. Let  $L$  be a Lie algebra over the ring  $Z$  of integers, let  $V$  be a  $Z$ -module, and let  $F$  be any commutative ring with unit.

$V_F = V \otimes_Z F$ .  $F_p$ , the field with  $p$  element  $V_p = V_{F_p}$ .

$L_F = L \otimes_Z F$ , an  $F$ -Lie algebra.

If  $V$  is an  $L$ -module, then  $V_F$  is an  $L_F$ -module.

$V_F^{L_F} = \{v \in V_F \mid kv = 0 \text{ for all } k \in L_F\}$ , the space of invariants.

$C^i(L_F, V_F) =$  space of  $i$ -cochains for the Lie algebra cohomology of  $L_F$  with values in  $V_F$   
 $= \text{Hom}_F(\hat{i}L_F, V_F)$ , where  $\hat{i}L_F$  is the  $i$ th exterior power of  $L_F$ .

$Z^i(L_F, V_F) =$  space of  $i$ -cocycles.

$B^i(L_F, V_F) =$  space of  $i$ -coboundaries.

$H^i(L_F, V_F) =$   $i$ th cohomology group.

$\delta^i =$   $i$ th coboundary operator.

Let  $\text{sl}_{n+1, F}$  be the Lie algebra of  $n + 1$  by  $n + 1$  trace 0 matrices with entries in  $F$ . Let  $E_{ij}$  be the trace zero matrix whose only nonzero entry is a 1 in the  $ij$ th position. The  $\{E_{ij}\}$  belong to the standard Chevalley basis for  $\text{sl}_{n+1, F}$ .

$Y_1 = E_{21}$ .

$h_F =$  space of diagonal matrices in  $\text{sl}_{n+1, F}$ .

$b_F =$  space of upper triangular matrices in  $\text{sl}_{n+1, F}$ .

$b =$  space of upper triangular matrices in  $\text{sl}_{n+1, Z}$ .

$\alpha_i$  is the element in the dual space of  $h_F$  whose value at the matrix with diagonal entries  $t_j$  is  $t_i - t_{i+1}$ .

1. Take a Lie algebra  $L$  over the ring  $Z$  of integers which is a free  $Z$ -module of finite rank. Let  $V$  be any  $L$ -module which is a free  $Z$ -module of finite rank. Let  $Q$  be the rational numbers.

Suppose that  $L_Q$  is a semisimple Lie algebra and that the space of  $L_Q$ -invariants in  $V_Q$  and the space of  $L_p$ -invariants in  $V_p$  have the same dimension, say  $m$ . Then we have the following theorem.

**THEOREM.** *If  $H^1(L_p, V_p)$  is nonzero, then  $H^2(L_p, V_p)$  is nonzero.*

*Proof.* The proof is a comparison of the ranks and dimensions of various spaces coming from cochain complexes. The canonical map  $C^2(L, V) \otimes_Z F_p \rightarrow C^2(L_p, V_p)$  is a linear isomorphism since  $L$  and  $V$  are free  $Z$ -modules. Furthermore, the image of  $Z^2(L, V) \otimes_Z F_p$  lies in  $Z^2(L_p, V_p)$ . Hence, the dimension of  $Z^2(L_p, V_p)$  is at least as large as the rank of the free  $Z$ -module  $Z^2(L, V)$ , which equals the dimension of  $Z^2(L_Q, V_Q)$ .

The spaces  $Z^i(L_Q, V_Q)$  and  $B^i(L_Q, V_Q)$  coincide for  $i = 1, 2$  since  $H^i(L_Q, V_Q) = (0)$  for  $i = 1, 2$  for a semisimple Lie algebra [2]. Therefore,  $\dim(Z^2(L_Q, V_Q))$  equals  $\dim(C^1(L_Q, V_Q)) - \dim(V_Q) + m$ .

The dimension of  $B^2(L_p, V_p)$ , which equals  $\dim(C^1(L_p, V_p)) - \dim(Z^1(L_p, V_p))$ , is less than  $\dim(C^1(L_p, V_p)) - \dim(B^1(L_p, V_p))$  since  $H^1(L_p, V_p)$  is nonzero by hypothesis. Therefore, the dimension of  $B^2(L_p, V_p)$  is less than  $\dim(C^1(L_p, V_p)) - \dim(V_p) + m$ , which equals  $\dim(C^1(L_q, V_q)) - \dim(V_q) + m$ . By the conclusions of the previous two paragraphs, the dimension of  $Z^2(L_p, V_p)$  exceeds the dimension of  $B^2(L_p, V_p)$  and  $H^2(L_p, V_p)$  is nonzero.

2. We will turn our attention specifically to the Lie algebra  $\mathfrak{sl}_{n+1, F_p}$  of the special linear group  $SL_{n+1, F_p}$  over the prime field  $F_p$ . Let  $\lambda_i$  be the character on the diagonal subgroup of  $SL_{n+1, F_p}$  whose value at the diagonal matrix with diagonal entries  $\{t_j\}_j$  is  $t_1 \cdots t_i$ . Let  $W$  be the irreducible  $SL_{n+1, F_p}$ -module whose highest weight is  $(p-2)\lambda_1 + \lambda_2$ .  $W$  will also stand for the space  $W$  with the differential  $\mathfrak{sl}_{n+1, F_p}$ -module structure. For any  $SL_{n+1, F_p}$ -module  $W'$ , we will give the weight of a line stable under the diagonal subgroup by a character on the diagonal subgroup, even though it may be the differential  $\mathfrak{sl}_{n+1, F_p}$ -module structure of  $W'$  that is under consideration.

In [4], we showed that the one-cohomology of  $\mathfrak{sl}_{n+1, F_p}$  at  $W$  is nonzero, except possibly for  $\mathfrak{sl}_{3, F_3}$ . Below, we associate to  $W$  a Weyl module  $V_p$  and show that  $\mathfrak{sl}_{n+1, F_p}$  has nonzero two-cohomology at  $V_p$ , except possibly when  $p = n + 1$ . Let  $V_q$  be the irreducible  $SL_{n+1, q}$ -module whose highest weight is  $(p-2)\lambda_1 + \lambda_2$ . We construct a  $Z$ -form  $V$  for  $V_q$  which is stable under the Kostant  $Z$ -form  $U_Z$  for the universal enveloping algebra of  $\mathfrak{sl}_{n+1, q}$  (relative to the standard basis for  $\mathfrak{sl}_{n+1, q}$ ) and whose reduction modulo  $p$ ,  $V_p$ , contains  $W$  as a submodule for its  $SL_{n+1, F_p}$ -module structure.

The standard  $Z$ -form of  $V_q$  is the  $U_Z$ -submodule that is generated from a highest weight vector  $[1, A]$ . The  $Z$ -form  $V$  is the variant that is produced as follows. Take a lowest weight vector  $v^*$  from the  $SL_{n+1, q}$ -module  $V_q^*$  that is contragradient to  $V_q$ . The weight of  $v^*$ , reduced modulo  $p$ , equals the lowest weight of  $W^*$ . The  $U_Z$ -module  $U_Z v^*$  is a  $Z$ -form for  $V_q^*$  for which the  $SL_{n+1, F_p}$ -module  $U_Z v^* \otimes_Z F_p$  contains a unique maximal proper submodule. The quotient module is isomorphic to  $W^*$  since the lowest weight of the quotient module is the weight of  $v^*$ , reduced modulo  $p$ . The  $U_Z$ -module  $V = \text{Hom}(U_Z v^*, Z)$  is a  $Z$ -form for  $V_q$  and the  $SL_{n+1, F_p}$ -module  $V_p = V \otimes_Z F_p$ , which is contragradient to  $U_Z v^* \otimes_Z F_p$ , has a unique irreducible submodule. This submodule is isomorphic to  $W$ .

The following lemma and its use in proving the proposition below were suggested by John Ballard. The proof has been consigned to the last section.

LEMMA. *If  $p \neq n + 1$ , then  $V_p$  contains no weight  $p\lambda$  where  $\lambda$*

is dominant.

**PROPOSITION.** *If  $p \neq n + 1$ , then  $H^1(\mathfrak{sl}_{n+1, F_p}, V_p)$  is nonzero.*

*Proof.* From the long exact sequence of cohomology associated with the exact sequence  $0 \rightarrow W \rightarrow V_p \rightarrow V_p/W \rightarrow 0$ , take the exact sequence  $(V_p/W)^{\mathfrak{sl}_{n+1, F_p}} \rightarrow H^1(\mathfrak{sl}_{n+1, F_p}, W) \rightarrow H^1(\mathfrak{sl}_{n+1, F_p}, V_p)$ . It will be enough to show that the first term is (0) since  $H^1(\mathfrak{sl}_{n+1, F_p}, W)$  is nonzero.  $(V_p/W)^{\mathfrak{sl}_{n+1, F_p}}$  is an  $\mathrm{SL}_{n+1, F_p}$ -submodule of  $V_p/W$ , which is trivial as an  $\mathfrak{sl}_{n+1, F_p}$ -module. A highest weight vector of this submodule would be dominant weight of the form  $p\lambda$ , which the lemma excludes from  $V_p$ . Therefore,  $V_p/W$  has (0) as its space of invariants.

**THEOREM.** *If  $p \neq n + 1$ , then  $H^2(\mathfrak{sl}_{n+1, F_p}, -)$  is nonzero at the Weyl module  $V_p$ .*

*Proof.* In the theorem of section one, the hypothesis that  $H^1(\mathfrak{sl}_{n+1, F_p}, V_p)$  be nonzero is satisfied by the proposition above. The hypothesis that the spaces of invariants of  $V_Q$  and  $V_p$  have the same dimension is satisfied. In fact,  $V_p^{\mathfrak{sl}_{n+1, F_p}}$  is zero, by the argument used in proving the proposition. Therefore,  $V_Q^{\mathfrak{sl}_{n+1, F_p}}$  is zero as well. We may conclude that  $H^2(\mathfrak{sl}_{n+1, F_p}, V_p)$  is nonzero.

3. We will show the origin of one nonzero cohomology class in  $H^2(\mathfrak{sl}_{n+1, F_p}, V_p)$ .

Denote  $\mathfrak{sl}_{n+1, Z}$  by  $L$ , denote the weight  $(p - 2)\lambda_1 + \lambda_2$  by  $\beta$ , and let  $W$  and  $V$  be the modules given in §2. In [4], we constructed a noncobounding one cocycle  $g$  for  $L_p$  with values in  $W$  with the properties that  $g$  is zero on  $b_p$  and  $g(Y_1)$  is a highest weight vector in  $W$ . The reduction modulo  $p$  of the unique line of weight  $\beta$  in  $V$  in the unique line of weight  $\beta$  in  $V_p$ . Choose any cochain  $g_z$  in  $C^1(L, V)$  for which  $g_z \otimes_Z F_p = g$ ,  $g_z(b) = (0)$ , and  $g_z(Y_1)$  is a vector of weight  $\beta$ . We will show that the 2-cochain  $f = ((1/p)\delta^1(g_z)) \otimes_Z F_p$  represents a nonzero cohomology class for  $L_p$  with values in  $V_p$ . Since  $\delta^1 g = 0$ ,  $\delta^1 g_z$  is divisible by  $p$  in  $C^2(L, V)$  and we may form the cochain  $(1/p)(\delta^1(g_z))$  in  $C^2(L, V)$ . This is a cocycle since  $\delta^2((1/p)(\delta^1(g_z))) = (1/p)(\delta^2 \delta^1 g_z) = 0$  and it maps  $b \wedge b$  to 0 since  $g_z(b) = (0)$ . Therefore,  $f = ((1/p)(\delta^1(g_z)) \otimes_Z F_p)$  is a two cocycle for  $L_p$  with values in  $V_p$ , which maps  $b_p \wedge b_p$  to 0. We show that  $f$  is not a coboundary.

Suppose that  $f$  were the coboundary  $\delta^1 u$  for some cochain  $u: L_p \rightarrow V_p$ . The restriction of  $u$  to  $b_p$  is a cocycle since  $f(b_p \wedge b_p) = (0)$ . By the lemma of §3.1 of [4], the canonical image of  $H^1_b(b_p, V_p)$  in  $H^1(h_p, V_p)$  is contained in the image of  $H^1(h_p, V_p^b)$  in  $H^1(h_p, V_p)$ ;

here, since  $V_p^{b_p} = (0)$ ,  $(0)$  is the image of  $H^1(b_p, V_p)$  in  $H^1(h_p, V_p)$ . Therefore, there is an element  $v$  of  $V_p$  such that  $u - \delta^0 v$  is zero on  $h_p$ . Now use the symbol  $u$  for the 'normalized' cochain  $u - \delta^0 v$ . We show that  $f$  cannot equal  $\delta^1 u$  by showing that  $f(Y_1 \wedge H_1)$  has a nonzero component of weight  $\beta$ , while  $\delta^1 u(Y_1 \wedge H_1)$  has zero for its component of weight  $\beta$ .  $\delta^1 u(Y_1 \wedge H_1) = (-2 - H_1)u(Y_1)$  has zero as its component of weight  $\beta$  since  $-2 - H_1$  annihilates vectors of weight  $\beta$ . By the construction of  $g_z$ ,  $g_z(Y_1)$  has as its component of weight  $\beta$  a vector  $x$  which is nonzero modulo  $p$ . Write  $g_z(Y_1) = x + y$ . Then  $(1/p)\delta^1 g_z(Y_1 \wedge H_1) = (1/p)(-2 - H_1)g_z(Y_1) = x + (1/p)(-2 - H_1)y$ , reduced modulo  $p$ , has a nonzero component of weight  $\beta$ .

Thus, we see that  $(1/p)\delta^1 g_z$  represents a cohomology class in  $H^2(\mathfrak{sl}_{n+1, z}, V)$  whose reduction modulo  $p$  gives a nonzero element of  $H^2(\mathfrak{sl}_{n+1, F_p}, V_p)$ . We wish to remark that the cohomology class of  $f$  does not depend on the choice of lifting of  $g$  to an integral cochain. In fact, if  $g'_z$  is another cochain such that  $g'_z \otimes_{\mathbb{Z}} F_p = g$ , then  $(1/p)\delta^1 g_z - (1/p)\delta^1 g'_z$  is the two coboundary  $\delta^1((1/p)(g_z - g'_z))$ , and  $(1/p)\delta^1 g_z$  and  $(1/p)\delta^1 g'_z$  represent the same class in  $H^2(\mathfrak{sl}_{n+1, z}, V)$ .

It would be good to know if  $H^2(\mathfrak{sl}_{n+1, F_p}, W)$  is also nonzero. In particular, does the cohomology class of the element  $f$  lie in the canonical image of  $H^2(\mathfrak{sl}_{n+1, l}, W)$  in  $H^2(\mathfrak{sl}_{n+1, F_p}, V_p)$ ?

4. Here we give a proof of the lemma of §2. The weights of the  $SL_{n+1, q}$ -module  $V_q$  have the form  $\beta - \sum a_i \alpha_i$ , where  $\beta = (p - 2)\lambda_1 + \lambda_2$  and the  $a_i$  are integers in the range  $0 \leq a_1, a_n \leq p - 1$ , and  $0 \leq a_i \leq p$  for  $i \neq 1, n$ , c.f. [4, 3.3]. The weights of the  $SL_{n+1, F_p}$ -module  $V_p$  inherit the same property. The roots  $\alpha_i$  are expressed in terms of the fundamental dominant weights  $\lambda_i$  as follows:  $\alpha_1 = 2\lambda_1 - \lambda_2$ ,  $\alpha_n = 2\lambda_n - \lambda_{n-1}$ , and  $\alpha_i = 2\lambda_i - \lambda_{i-1} - \lambda_{i+1}$  for  $i \neq 1, n$ .

Suppose that  $p\lambda = \beta - \sum a_i \alpha_i$  is a weight of  $V_p$ , where  $\lambda$  is a dominant weight  $\sum b_i \lambda_i$ ,  $b_i \geq 0$ . Let  $j$  be the highest index such that  $a_j$  is nonzero. We will show that  $j$  must be  $n$  and that  $p = n + 1$ . If  $n > j > 2$ , then the coefficient of  $\lambda_{j+1}$  in  $\beta - \sum a_i \alpha_i = p\lambda$  is  $pb_{j+1} = a_j$ , and  $a_j = p$  since  $0 < a_j \leq p$ . The coefficient of  $\lambda_j$  is  $pb_j = -2a_j + a_{j-1} = -2p + a_{j-1}$ . This is not possible since  $b_j$  is nonnegative and  $a_{j-1} \leq p$ . A similar argument shows that it is not possible to have  $n > j$  when  $j = 1$  or  $2$ . Hence,  $j$  must be  $n$ .

The coefficient of  $\lambda_n$  in  $p\lambda = \beta - \sum a_i \alpha_i$  is  $pb_n = -2a_n + a_{n-1}$ . Since  $b_n$  is nonnegative,  $a_n$  is positive, and  $a_{n-1} \leq p$ ,  $a_{n-1}$  must equal  $2a_n$ . More generally, we will check that the formula  $a_i = (n + 1 - i)a_n$  holds for any  $i \geq 2$ . Suppose that it holds for all  $i \geq k$ , where  $k$  is an index in the range  $n - 1 > k > 2$ . The coefficient of  $\lambda_k$  is  $pb_k = a_{k-1} + a_{k+1} - 2a_k = a_{k-1} - (n + 2 - k)a_n$ . As above for  $k = n - 1$ ,

we conclude that  $a_{k-1} = (n + 2 - k)a_n$ . By the formula, the coefficient of  $\lambda_2$  is  $pb_2 = 1 + a_1 + a_3 - 2a_2 = 1 + a_1 - na_n$  and so,  $a_1 = na_n - 1$ . The coefficient of  $\lambda_1$  is  $pb_1 = p - 2 + a_2 - 2a_1 = p - (n + 1)a_n$ , and so,  $p = (n + 1)a_n$ . We may conclude that  $p = n + 1$ ,  $a_n = 1$ ,  $\beta = (n - 1)\alpha_1 + \sum_{i=2}^n (n + 1 - i)\alpha_i$ , and that only in characteristic  $p = n + 1$  can  $V_p$  have a dominant weight of the form  $p\lambda$ .

#### REFERENCES

1. A. Borel et al., *Seminar on Algebraic Groups and Related Finite Groups*, Lecture Notes in Mathematics, Springer-Verlag, Heidelberg, 1970, Vol. 131.
2. C. Chevalley and S. Eilenberg, *Cohomology theory of Lie groups and Lie algebras*, TAMS, **63** (1948), 85-124.
3. G. Hochschild, *Representations of restricted Lie algebras of characteristic  $p$* , Proc. Amer. Math. Soc., **5** (1954), 603-605.
4. J. Sullivan, *Lie algebra cohomology at irreducible modules*, Illinois J. Math., **23** (1979), 363-373.

Received June 19, 1978. Supported in part by a National Science Foundation Grant.

UNIVERSITY OF WASHINGTON  
SEATTLE, WA 98195

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California  
Los Angeles, California 90024

HUGO ROSSI

University of Utah  
Salt Lake City, UT 84112

C. C. MOORE AND ANDREW OGG

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. FINN AND J. MILGRAM

Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON



Gert Einar Torsten Almkvist, <i>Invariants, mostly old ones</i> .....	1
Hyman Bass, <i>Groups of integral representation type</i> .....	15
A. Białynicki-Birula, <i>On action of <math>SL(2)</math> on complete algebraic varieties</i> .....	53
Frederick Paul Greenleaf and Martin Allen Moskowitz, <i>Groups of automorphisms of Lie groups: density properties, bounded orbits, and homogeneous spaces of finite volume</i> .....	59
Raymond Taylor Hoobler, <i>A cohomological interpretation of Brauer groups of rings</i> .....	89
Irving Kaplansky, <i>Superalgebras</i> .....	93
Jerrold Lewis Kleinstein and Alex I. Rosenberg, <i>Succinct and representational Witt rings</i> .....	99
E. R. Kolchin, <i>On universal extensions of differential fields</i> .....	139
Andy R. Magid, <i>Analytic subgroups of affine algebraic groups. II</i> .....	145
Calvin Cooper Moore, <i>The Mautner phenomenon for general unitary representations</i> .....	155
George Daniel Mostow, <i>On a remarkable class of polyhedra in complex hyperbolic space</i> .....	171
Brian Lee Peterson, <i>Extensions of pro-affine algebraic groups. II</i> .....	277
John Henry Reineohl, <i>Lie algebras and affine algebraic groups</i> .....	287
Maxwell Alexander Rosenlicht, <i>Differential valuations</i> .....	301
John Brendan Sullivan, <i>The second Lie algebra cohomology group and Weyl modules</i> .....	321
Moss Eisenberg Sweedler, <i>Right derivations and right differential operators</i> .....	327
Bostwick Frampton Wyman, <i>Time varying linear discrete-time systems. II. Duality</i> .....	361