

# Pacific Journal of Mathematics

## **DIRECT LIMIT GROUPS AND THE KEESLING-MARDEŠIĆ SHAPE FIBRATION**

KENNETH R. GOODEARL AND THOMAS BENNY RUSHING

# DIRECT LIMIT GROUPS AND THE KEESLING-MARDEŠIĆ SHAPE FIBRATION

K. R. GOODEARL AND T. B. RUSHING

We show that the Keesling-Mardešić shape fibration has an uncountable number of fibers of different shape type. This is done by showing that an uncountable number of nonisomorphic groups can arise as direct limits of direct limit sequences having all groups  $Z \oplus Z$  and all bonding homomorphisms given by one of the two matrices  $\begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$  or  $\begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix}$ .

**A. Introduction.** The notion of shape fibration has been developed by Mardešić and Rushing in [4, 5, 6]. In [4] the following question was raised: Let  $p: E \rightarrow B$  be a shape fibration and let  $x, y \in B$  be points belonging to the same component. Do the fibers  $p^{-1}(x)$  and  $p^{-1}(y)$  have the same shape? In that paper, this was shown to be the case if  $x$  and  $y$  belong to the same path component, and in [5, Corollary 1], this was shown to be the case if  $x$  and  $y$  belong to a subcontinuum of  $B$  of trivial shape. In [3], Keesling and Mardešić gave an ingenious example of a shape fibration with connected base space and showed that it has two fibers of different shape. We show that their example, in fact, has an uncountable number of fibers of different shape.

Let an inverse limit sequence of spaces be given where each space is  $T^2 = S^1 \times S^1$  and each bonding map is given by one of the two matrices  $\begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$  or  $\begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix}$ . Here the given matrices each induce a continuous homomorphism on  $\mathbf{R}^2$  which in turn defines a continuous homomorphism on  $T^2$  via the covering map  $e \times e: \mathbf{R}^2 \rightarrow T^2$  where  $e(t) = e^{2\pi it}$ . Let  $X$  be the inverse limit space of such a sequence. Then the discussion in [3, §§2, 3] and [3, Lemma 1] implies that the Keesling-Mardešić fibration has a fiber homeomorphic to  $X$ .

Consider the direct limit sequence of groups where each group is  $Z \oplus Z$  and each bonding map is given by the transpose of the corresponding matrix in the inverse sequence above. Then by [3, §5], the first Čech cohomology group  $\check{H}^1(X, Z)$  of  $X$  is isomorphic to the direct limit of this sequence. Since Čech cohomology is an invariant of shape type, it suffices to show that there are an uncountable number of isomorphism classes of such direct limit groups.

Thus the groups we are interested in arise as direct limits of sequences

$$G_0 \xrightarrow{f_0} G_1 \xrightarrow{f_1} G_2 \xrightarrow{f_2} \dots$$

of abelian groups and group homomorphisms, where each  $G_n = \mathbf{Z} \oplus \mathbf{Z}$  and each  $f_n$  is given by one of two matrices  $\begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$  or  $\begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix}$ . (We are viewing the elements of  $\mathbf{Z} \oplus \mathbf{Z}$  as column vectors, so that  $f_n$  acts as left multiplication by the chosen matrix.) Our task is to show that among the groups  $G$  appearing as direct limits of the above form, there are uncountably many isomorphism classes. This result is folklore among abelian group theorists. An effort is made here to give a presentation that is as accessible as possible.

For convenience in labelling these groups and comparing them, we shall use a few computations within the field of 2-adic numbers. For the reader's convenience, we review the basics of this field and its construction in the following section. Details may be found, for example, in [1, Chapter I] or [2, Chapter V, §5]. We would like to thank Richard S. Pierce for bringing these 2-adic methods to our attention.

**B. 2-Adic numbers.** For any rational number  $x$ , the *2-adic valuation of  $x$* , denoted  $\nu_2(x)$ , is defined as follows. If  $x = 0$ , then  $\nu_2(x) = \infty$ . If  $x \neq 0$ , then  $x$  may be uniquely written in the form  $x = 2^n a/b$ , where  $n, a, b$  are integers,  $a$  and  $b$  are odd, and  $b \neq 0$ ; in this case,  $\nu_2(x) = n$ . Clearly  $\nu_2(xy) = \nu_2(x) + \nu_2(y)$ , and it may be checked that  $\nu_2(x + y) \geq \min\{\nu_2(x), \nu_2(y)\}$ .

The 2-adic valuation  $\nu_2(-)$  is used to define the *2-adic absolute value*  $|\cdot|_2$  on  $\mathbf{Q}$ , by the formula

$$|x|_2 = 2^{-\nu_2(x)}.$$

(Some authors refer to  $|\cdot|_2$ , rather than  $\nu_2(-)$ , as the 2-adic valuation on  $\mathbf{Q}$ .) The basic properties of  $\nu_2(-)$  translate into the following basic properties of  $|\cdot|_2$ :

(i)  $|x|_2$  is a nonnegative real number, and  $|x|_2 = 0$  if and only if  $x = 0$ .

(ii)  $|xy|_2 = |x|_2 |y|_2$ .

(iii)  $|x + y|_2 \leq \max\{|x|_2, |y|_2\} \leq |x|_2 + |y|_2$ .

It is clear from these properties that the rule  $\delta_2(x, y) = |x - y|_2$  defines a metric on  $\mathbf{Q}$ , called the *2-adic metric*.

We shall use  $\mathbf{Q}_2^*$  to denote the completion of  $\mathbf{Q}$  with respect to the metric  $\delta_2$ . Since addition and multiplication in  $\mathbf{Q}$  are uniformly continuous with respect to  $\delta_2$ , they induce addition and multiplication operations in  $\mathbf{Q}_2^*$ . Then  $\mathbf{Q}_2^*$  becomes a field, known as the *field of 2-adic numbers*. The completion of  $\mathbf{Z}$  with respect to  $\delta_2$  is then a subring  $\mathbf{Z}_2^*$  of  $\mathbf{Q}_2^*$ , known as the *ring of 2-adic integers*, and it may be checked that  $\mathbf{Q} \cap \mathbf{Z}_2^* = \mathbf{Z}$ .

The elements of  $Z_2^*$  may be explicitly represented as (limits of) series. Given any sequence  $\varepsilon_0, \varepsilon_1, \varepsilon_2, \dots$  of zeros and ones, the series  $\sum_{k=0}^\infty \varepsilon_k 2^k$  converges in  $Z_2^*$ ; conversely, every element of  $Z_2^*$  may be uniquely represented as such a series. In particular, it follows that  $Z_2^*$  is uncountable.

**C. Change of perspective.** Given a group  $G$  obtained as a direct limit as in §A, we wish to present  $G$  in a form more suitable for computation. Each of the maps  $f_k$  is given by a matrix  $\begin{pmatrix} 1 & \varepsilon_k \\ 0 & 2 \end{pmatrix}$ , where  $\varepsilon_k = 0$  or  $1$ . Set  $\alpha_n = \sum_{k=0}^n \varepsilon_k 2^k$  for all  $n = 0, 1, 2, \dots$ , and let  $\alpha$  be the 2-adic integer  $\sum_{k=0}^\infty \varepsilon_k 2^k$ .

Let  $V$  be a 2-dimensional vector space over  $\mathbb{Q}$ , with basis  $\{x_1, x_2\}$ . Set  $z_0 = x_2$ , set

$$(1) \quad z_n = (x_2 - \alpha_{n-1}x_1)/2^n$$

for  $n = 1, 2, \dots$ , and for all  $n$  let  $A_{\alpha,n}$  be the subgroup of  $V$  generated by  $x_1$  and  $z_n$ . Observing that

$$(2) \quad z_n = \varepsilon_n x_1 + 2z_{n+1},$$

we see that  $A_{\alpha,n} \subseteq A_{\alpha,n+1}$ . Thus  $A_\alpha = \bigcup_{n=0}^\infty A_{\alpha,n}$  is a subgroup of  $V$ , and we claim that  $A_\alpha \cong G$ .

Since  $A_{\alpha,n}$  is a free abelian group with basis  $\{x_1, z_n\}$ , there is a group isomorphism  $g_n: G_n \rightarrow A_{\alpha,n}$  given by the rule  $g_n \begin{pmatrix} a \\ b \end{pmatrix} = ax_1 + bz_n$ . Note, using (2), that the following diagram commutes:

$$\begin{array}{ccccccc} G_0 & \xrightarrow{f_0} & G_1 & \xrightarrow{f_1} & G_2 & \xrightarrow{f_2} & \dots \\ g_0 \downarrow & & g_1 \downarrow & & g_2 \downarrow & & \\ A_{\alpha,0} & \xrightarrow{\cong} & A_{\alpha,1} & \xrightarrow{\cong} & A_{\alpha,2} & \xrightarrow{\cong} & . \end{array}$$

As the direct limits of the top and bottom rows are  $G$  and  $A_\alpha$ , we obtain  $G \cong A_\alpha$ , as claimed.

There is a projection  $V \rightarrow \mathbb{Q}$  mapping  $x_1 \mapsto 0$  and  $x_2 \mapsto 1$ . The image of  $A_\alpha$  under this projection is the group of all rational numbers of the form  $a/2^n$  (where  $a, n \in \mathbb{Z}$  and  $n \geq 0$ ), which is not finitely generated. Therefore  $A_\alpha$  is not finitely generated.

**D. Isomorphic groups.** The question now is, for what values of  $\alpha$  are the groups  $A_\alpha$  isomorphic? Consider another group  $A_\beta$ , where  $\beta = \sum_{k=0}^\infty \delta_k 2^k$  in  $Z_2^*$ , and each  $\delta_k = 0$  or  $1$ . Set  $\beta_n = \sum_{k=0}^n \delta_k 2^k$  for all  $n$ . Set  $w_0 = x_2$ , set

$$(3) \quad w_n = (x_2 - \beta_{n-1}x_1)/2^n$$

for  $n = 1, 2, \dots$ , and for all  $n$  let  $A_{\beta, n}$  be the subgroup of  $V$  generated by  $x_1$  and  $w_n$ . Then

$$(4) \quad w_n = \delta_n x_1 + 2w_{n+1}$$

for all  $n$ , so that  $A_{\beta, n} \subseteq A_{\beta, n+1}$ , and we set  $A_\beta = \bigcup_{n=0}^{\infty} A_{\beta, n}$ .

We claim that if  $A_\alpha \cong A_\beta$ , then

$$(5) \quad (\alpha r_{12} - r_{22})\beta = r_{21} - \alpha r_{11}$$

for some integers  $r_{ij}$ .

Thus assume that there is a group isomorphism  $A_\alpha \rightarrow A_\beta$ . Observing that we may identify  $A_\alpha \otimes_{\mathbb{Z}} \mathbb{Q}$  and  $A_\beta \otimes_{\mathbb{Z}} \mathbb{Q}$  with  $V$ , we see that the group isomorphism  $A_\alpha \rightarrow A_\beta$  induces a vector space automorphism  $g: V \rightarrow V$  such that  $g(A_\alpha) = A_\beta$ . There exist integers  $r_{11}, r_{12}, r_{21}, r_{22}, s$  (with  $s \neq 0$ ) such that

$$(6) \quad g(x_i) = (r_{i1}/s)x_1 + (r_{i2}/s)x_2,$$

and we shall prove (5) using these  $r_{ij}$ . Note, from combining (6) and (1), that  $g(z_0) = (r_{21}/s)x_1 + (r_{22}/s)x_2$  and

$$(7) \quad g(z_n) = [(r_{21} - \alpha_{n-1}r_{11})/2^n s]x_1 + [(r_{22} - \alpha_{n-1}r_{12})/2^n s]x_2$$

for  $n = 1, 2, \dots$ .

**E. The computation.** As  $g(A_\alpha) = A_\beta$ , there is a nonnegative integer  $t$  such that  $g(x_1) \in A_{\beta, t}$ . For all  $n = 0, 1, 2, \dots$ , let  $k(n)$  be the least integer such that  $k(n) \geq t$  and  $g(z_n) \in A_{\beta, k(n)}$ . In view of (2), we see that  $g(z_n)$  lies in  $A_{\beta, k(n+1)}$ , whence  $k(n) \leq k(n+1)$ . Thus  $k(0) \leq k(1) \leq \dots$  is an increasing sequence of nonnegative integers, and we claim this sequence is unbounded.

If there is an integer  $k$  with all  $k(n) \leq k$ , then  $g(x_1) \in A_{\beta, k}$  and all  $g(z_n) \in A_{\beta, k}$ , whence  $g(A_\alpha) \subseteq A_{\beta, k}$ . Since  $A_{\beta, k}$  is finitely generated, this would imply that  $A_\alpha$  is finitely generated, which is false. Thus the  $k(n)$  are unbounded, as claimed.

In particular, there exists a positive integer  $N$  such that  $k(n) > t$  for all  $n \geq N$ .

As  $g(z_n) \in A_{\beta, k(n)}$ , we must have

$$(8) \quad g(z_n) = a_n x_1 + b_n w_{k(n)}$$

for some  $a_n, b_n \in \mathbb{Z}$ . If  $k(n) > t$  and  $b_n = 2c$  for some  $c \in \mathbb{Z}$ , then we see using (4) that

$$(9) \quad g(z_n) = a_n x_1 + c(w_{k(n)-1} - \delta_{k(n)-1} x_1)$$

and so  $g(z_n) \in A_{\beta, k(n)-1}$ , which contradicts the minimality of  $k(n)$ . Thus

$b_n$  must be odd whenever  $k(n) > t$ . In particular,  $b_n$  is odd for all  $n \geq N$ .

Comparing (7) and (8), with the help of (3), we obtain

$$(10) \quad (2^{k(n)}a_n - \beta_{k(n)-1}b_n)/2^{k(n)} = (r_{21} - \alpha_{n-1}r_{11})/2^n s$$

$$(11) \quad b_n/2^{k(n)} = (r_{22} - \alpha_{n-1}r_{12})/2^n s$$

for all  $n \geq 1$ . Cross-multiplying (10) and (11) yields

$$(12) \quad (r_{21} - \alpha_{n-1}r_{11})b_n = (r_{22} - \alpha_{n-2}r_{12})(2^{k(n)}a_n - \beta_{k(n)-1}b_n).$$

When  $n \geq N$ , we have  $b_n$  odd and so can divide by it, yielding

$$(13) \quad r_{21} - \alpha_{n-1}r_{11} = (r_{22} - \alpha_{n-1}r_{12})[2^{k(n)}(a_n/b_n) - \beta_{k(n)-1}].$$

We intend to take the limit of (13) in  $\mathbb{Q}_2^*$  as  $n \rightarrow \infty$ .

As  $b_n$  is odd,  $\nu_2(b_n) = 0$  and so  $|b_n|_2 = 1$ . As  $a_n$  is an integer,  $\nu_2(a_n) \geq 0$  and so  $|a_n|_2 \leq 1$ . Consequently,

$$(14) \quad |2^{k(n)}(a_n/b_n)|_2 \leq |2^{k(n)}|_2 = 2^{-k(n)}.$$

Since the  $k(n)$  form an unbounded increasing sequence,  $2^{-k(n)} \rightarrow 0$ , hence  $2^{k(n)}(a_n/b_n) \rightarrow 0$  in  $\mathbb{Q}_2^*$ .

Thus we can now compute the limit of (13) as  $n \rightarrow \infty$ , obtaining

$$(14) \quad r_{21} - \alpha r_{11} = (r_{22} - \alpha r_{12})(-\beta),$$

which is equivalent to (5).

F. Uncountably many groups. Put an equivalence relation  $\sim$  on  $\mathbb{Z}_2^*$ , defined by  $\alpha \sim \beta$  if and only if  $A_\alpha \cong A_\beta$ . If  $\alpha \notin \mathbb{Z}$  and  $\alpha \sim \beta$ , then (5) shows that

$$(15) \quad \beta = (r_{21} - \alpha r_{11})/(r_{22} - \alpha r_{12})$$

for some integers  $r_{ij}$ , hence there are only countably many possibilities for  $\beta$ . Thus if  $\alpha \notin \mathbb{Z}$ , then the equivalence class of  $\alpha$  is countable. For  $\alpha \in \mathbb{Z}$ , either the equivalence class of  $\alpha$  is contained in  $\mathbb{Z}$ , or it coincides with the equivalence class of some  $\beta \notin \mathbb{Z}$ ; in either case, the equivalence class of  $\alpha$  is again countable.

Thus all the equivalence classes with respect to  $\sim$  are countable. As  $\mathbb{Z}_2^*$  is uncountable, there must be uncountably many equivalence classes. Therefore there are uncountably many isomorphism classes of the groups  $A_\alpha$ .

### REFERENCES

1. N. Koblitz, *p-adic Numbers, p-adic Analysis, and Zeta-Functions*, New York, Springer-Verlag, 1977. (Graduate Texts in Math., 58.)

2. N. Jacobson, *Lectures in Abstract Algebra*, Vol. III, Princeton, Van Nostrand, 1964.
3. James Keesling and Sibe Mardešić, *A shape fibration with fibers of different shape*, Pacific J. Math., (to appear).
4. Sibe Mardešić and T. B. Rushing, *Shape fibrations I*, Gen. Top. and its Appl., **9** (1978), 193-215.
5. ———, *Shape fibrations II*, Rocky Mountain J. Math., **9** (2), (1979), 283-298.
6. ———, *n-shape fibrations*, Topology Proceedings, **3** (1978), 429-459.

Received March 30, 1979. Both authors partially supported by National Science Foundation grants.

UNIVERSITY OF UTAH  
SALT LAKE CITY, UT 84112

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California  
Los Angeles, CA 90024

HUGO ROSSI

University of Utah  
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University  
Stanford, CA 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$84.00 a year (6 Vols., 12 issues). Special rate: \$42.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).

8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1980 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan



Graham Donald Allen, David Alan Legg and Joseph Dinneen Ward, <i>Hermitian liftings in Orlicz sequence spaces</i> .....	379
George Bachman and Alan Sultan, <i>On regular extensions of measures</i> .....	389
Bruce Alan Barnes, <i>Representations Naimark-related to *-representations; a correction: "When is a representation of a Banach *-algebra Naimark-related to a *-representation?"</i> .....	397
Earl Robert Berkson, <i>One-parameter semigroups of isometries into <math>H^p</math></i> .....	403
M. Brodmann, <i>Piecewise catenarian and going between rings</i> .....	415
Joe Peter Buhler, <i>A note on tamely ramified polynomials</i> .....	421
William Lee Bynum, <i>Normal structure coefficients for Banach spaces</i> .....	427
Lung O. Chung, <i>Biharmonic and polyharmonic principal functions</i> .....	437
Vladimir Drobot and S. McDonald, <i>Approximation properties of polynomials with bounded integer coefficients</i> .....	447
Giora Dula and Elyahu Katz, <i>Recursion formulas for the homology of <math>\Omega(X \vee Y)</math></i> .....	451
John A. Ernest, <i>The computation of the generalized spectrum of certain Toeplitz operators</i> .....	463
Kenneth R. Goodearl and Thomas Benny Rushing, <i>Direct limit groups and the Keesling-Mardešić shape fibration</i> .....	471
Raymond Heitmann and Stephen Joseph McAdam, <i>Good chains with bad contractions</i> .....	477
Patricia Jones and Steve Chong Hong Ligh, <i>Finite hereditary near-ring-semigroups</i> .....	491
Yoshikazu Katayama, <i>Isomorphisms of the Fourier algebras in crossed products</i> .....	505
Meir Katchalski and Andrew Chiang-Fung Liu, <i>Symmetric twins and common transversals</i> .....	513
Mohammad Ahmad Khan, <i>Chain conditions on subgroups of LCA groups</i> .....	517
Helmut Kröger, <i>Padé approximants on Banach space operator equations</i> .....	535
Gabriel Michael Miller Obi, <i>An algebraic extension of the Lax-Milgram theorem</i> .....	543
S. G. Pandit, <i>Differential systems with impulsive perturbations</i> .....	553
Richard Pell, <i>Support point functions and the Loewner variation</i> .....	561
J. Hyam Rubinstein, <i>Dehn's lemma and handle decompositions of some 4-manifolds</i> .....	565
James Eugene Shirey, <i>On the theorem of Helley concerning finite-dimensional subspaces of a dual space</i> .....	571
Oved Shisha, <i>Tchebycheff systems and best partial bases</i> .....	579
Michel Smith, <i>Large indecomposable continua with only one composant</i> .....	593
Stephen Tefteller, <i>Existence of eigenvalues for second-order differential systems</i> .....	601