

Pacific Journal of Mathematics

**FREDHOLM THEORY OF PARTIAL DIFFERENTIAL
EQUATIONS ON COMPLETE RIEMANNIAN MANIFOLDS**

ROBERT COLMAN MCOWEN

FREDHOLM THEORY OF PARTIAL DIFFERENTIAL EQUATIONS ON COMPLETE RIEMANNIAN MANIFOLDS

ROBERT C. MCOWEN

This paper studies necessary and sufficient conditions for differential operators to be Fredholm on the Sobolev spaces of a complete (not necessarily compact) Riemannian manifold Ω . The conditions are formulated algebraically in terms of the nonvanishing of the operator's principal symbol on Ω (ellipticity) and its "total symbol" at infinity of Ω . The operators considered arise by taking sums of products of vector fields, all of whose covariant derivatives vanish at infinity; and the study involves C^* -algebra techniques. The required technical restrictions on the curvature and topology of Ω near infinity are much weaker than those in earlier joint work with H. O. Cordes.

0. Introduction. Let Ω be an n -dimensional paracompact C^∞ -manifold with complete Riemannian metric $ds^2 = g_{ij}dx^i dx^j$ and surface measure $d\mu = \sqrt{g} dx$ where $g = \det(g_{ij})$. Let $\Delta = g^{ij}\nabla_i\nabla_j$, the Beltrami-Laplace operator on Ω , where $(g^{ij}) = (g_{ij})^{-1}$ and ∇ denotes covariant differentiation with respect to the Riemannian connection. Then $A = (1 - \Delta)^{-1/2}$ is a positive-definite operator in $\mathcal{L}(\mathfrak{H})$, the bounded operators over the Hilbert space $\mathfrak{H} = L^2(\Omega, d\mu)$. Define the N th-Sobolev space $\mathfrak{H}_N \subset \mathfrak{H}$ for $N = 1, 2, \dots$ by requiring $A^N: \mathfrak{H} \rightarrow \mathfrak{H}_N$ to be an isometric isomorphism. It was shown in [3] that $C_0^\infty(\Omega)$ is dense in each \mathfrak{H}_N .

Now suppose we are given a differential operator L on Ω , of order N , such that we obtain a bounded map $L: \mathfrak{H}_N \rightarrow \mathfrak{H}$. We may ask the question when is L Fredholm (i.e., when are $\ker L$ and $\text{coker } L$ finite-dimensional subspaces of \mathfrak{H}_N and \mathfrak{H} respectively)? If Ω is compact, Seeley [13] showed that L is Fredholm if and only if L is elliptic (i.e., the principal symbol of L never vanishes on the cosphere bundle $S^*\Omega$). For the case $\Omega = \mathbf{R}^n$, on the other hand, ellipticity is not sufficient to imply L is Fredholm; Cordes and Herman [4] derived necessary and sufficient Fredholm criteria in terms of the "total symbol" of L .

In [4] the techniques involved considering operators LA^N as generators of a C^* -algebra $\mathfrak{A} \subset \mathcal{L}(\mathfrak{H})$ which is commutative modulo the compact ideal, \mathfrak{K} . The symbol of LA^N is then defined as the continuous function σ_{LA^N} on the maximal ideal space \mathcal{M} of $\mathfrak{A}/\mathfrak{K}$ provided by the Gel'fand theory. Thus LA^N is Fredholm if and only

if σ_{LAN} is never zero on M , and algebraic criteria are obtained by giving a precise description of M and σ_{LAN} .

In this paper we shall attempt to generalize the results of [4] and [13] to noncompact Riemannian manifolds Ω . We shall use techniques similar to those in [4], generating a C^* -algebra \mathfrak{A} which is commutative modulo \mathcal{K} , and then attempting to describe the maximal ideal space M and functions σ_{LAN} . Of course a precise description of M can only be made for specified Ω ; but our main result, Theorem 4.1, yields a sufficient condition for $L: \mathfrak{S}_N \rightarrow \mathfrak{S}$ to be Fredholm, namely that the "formal algebra symbol" $\hat{\sigma}_{LAN}$ never vanish on the space $\partial P^*\Omega$ (see §4 for definitions). Thus operators such as $(\lambda - \Delta)^N$ are Fredholm on the Sobolev spaces of Ω for $\lambda > 0$, whereas for $\lambda = 0$ we need to use weighted Sobolev spaces (see [2] and [11]). On the other hand, ellipticity is certainly a necessary condition for L to be Fredholm (c.f. §2).

This paper summarizes the principal results of the author's Ph.D. thesis, and generalizes earlier joint work with his advisor, H. O. Cordes, whose contributions to this research are also gratefully acknowledged.

1. The C^* -algebra for singular elliptic theory. We begin by describing the (global) differential operators that we will be considering. Suppose l_α is a C^∞ -contravariant tensor field of degree $\alpha \geq 0$ on Ω . Then in local coordinates l_α has components $l_\alpha^{i_1 \dots i_\alpha}$, and we may define a differential operator $l_\alpha \nabla^\alpha$ by

$$(1.1) \quad l_\alpha \nabla^\alpha u = l_\alpha^{i_1 \dots i_\alpha} \nabla_{i_1} \dots \nabla_{i_\alpha} u$$

for $u \in C_0^\infty(\Omega)$. (Note here and throughout this paper we employ the usual summation convention for tensor indices.) The differential operators we consider are just sums $L = \sum_{\alpha=0}^N l_\alpha \nabla^\alpha$.

For any tensor field T , we shall denote its length (a function over Ω) by $|T|$. In order to analyze the behavior of our operators "at infinity" let us make the following definition: if f is a (complex-valued) function on Ω , we shall write $\lim_{x \rightarrow \infty} f = M$ provided for any $\varepsilon > 0$ there is a compact set $K \subset \Omega$ such that $|f(x) - M| < \varepsilon$ for all $x \in \Omega \setminus K$. We shall consider the following conditions on the curvature tensor R .

Condition H. $|\nabla^k R|$ is bounded over Ω for all $k \geq 0$.

Condition H₀. In addition to Condition H, $\lim_{x \rightarrow \infty} |\nabla^k R| = 0$ for all $k \geq 1$.

Throughout this paper we shall be assuming Condition H with-

out further mentioning it.

We may now prove the following

THEOREM 1.1. *If $L = \sum_{\alpha=0}^N l_\alpha \nabla^\alpha$ has bounded coefficients (i.e., each $|l_\alpha|$ is bounded over Ω), then $LA^N \in \mathcal{L}(\mathfrak{S})$. If in addition L has coefficients vanishing at infinity (i.e., $\lim_{x \rightarrow \infty} |l_\alpha| = 0$ for each α), then $ALA^N \in \mathcal{H}(\mathfrak{S})$.*

Proof. It suffices to consider top-order terms only: $L = l_N \nabla^N$. Let $\psi \in C_0^\infty(\Omega)$. Condition H implies the following estimate for the L^2 -norms:

$$(1.2) \quad \|\nabla^N \psi\|^2 \leq C \left(\sum_{0 \leq k \leq (N/2)} \|\Delta^k \psi\|^2 + \sum_{0 \leq k \leq (N-1/2)} \|\nabla \Delta^k \psi\|^2 \right)$$

where C is a constant. (Note: if T is a tensor with length $|T|$, then $\|T\|^2 = \int_\Omega |T|^2 d\mu$. C. f. Aubin [1] for this notation, and Proposition 3 in [1] for the estimate (1.2).) So

$$(1.3) \quad \begin{aligned} \|L\psi\|^2 &\leq \left(\sup_{x \in \Omega} |l_N|^2 \right) \|\nabla^N \psi\|^2 \\ &\leq C_1 \left(\sum_{0 \leq k \leq (N/2)} \|\Delta^k \psi\|^2 + \sum_{0 \leq k \leq (N-1/2)} \|\nabla \Delta^k \psi\|^2 \right) \\ &\leq C_2 \left(\sum_{0 \leq k \leq (N/2)} \|(1 - \Delta)^k \psi\|^2 + \sum_{0 \leq k \leq (N-1/2)} \|\nabla(1 - \Delta)^k \psi\|^2 \right) \\ &\leq C_3 \|(1 - \Delta)^{N/2} \psi\|^2 \end{aligned}$$

where the C_i are constants and for the last inequality we have used the estimates (for $\phi \in C_0^\infty(\Omega)$)

$$(1.4) \quad \begin{aligned} \|\phi\|^2 &\leq \|(1 - \Delta)^{1/2} \phi\|^2 \leq \|(1 - \Delta)\phi\|^2 \\ \|\nabla \phi\|^2 &= \|(1 - \Delta)^{1/2} \phi\|^2 - \|\phi\|^2 \leq \|(1 - \Delta)^{1/2} \phi\|^2. \end{aligned}$$

Since $A^{-N}C_0^\infty(\Omega)$ is dense in \mathfrak{S} (c.f. Cordes [3]), estimate (1.3) proves $LA^N \in \mathcal{L}(\mathfrak{S})$. Now suppose $\lim_{x \rightarrow \infty} |l_N| = 0$ and for $\varepsilon > 0$ choose $K \subset \Omega$, compact, such that $|l_N| < \varepsilon$ for $x \in \Omega \setminus K$. Let $\phi \in C_0^\infty(\Omega)$ with $\text{supp } \phi \subset K$ and define $L_\varepsilon = \phi L$. Then $AL_\varepsilon A^N = A\phi \cdot LA^N \in \mathcal{H}(\mathfrak{S})$, since $LA^N \in \mathcal{L}(\mathfrak{S})$ and $A\phi \in \mathcal{H}(\mathfrak{S})$ by Theorem 3.1 in [5]. Since $AL_\varepsilon A^N \rightarrow ALA^N$ in norm as $\varepsilon \rightarrow 0$, we also have $ALA^N \in \mathcal{H}(\mathfrak{S})$.

In order to do Fredholm theory, we must restrict our attention to the following classes of operators

$$T^N = \left\{ L = \sum_{\alpha=0}^N l_\alpha \nabla^\alpha : \text{each } |l_\alpha| \text{ is bounded over } \Omega \text{ and satisfies} \right. \\ \left. \lim_{x \rightarrow \infty} |\nabla^k l_\alpha| = 0 \text{ for all } k \geq 1 \right\}.$$

Let $T_0^N \subset T^N$ be those operators for which $\lim_{x \rightarrow \infty} |l_\alpha| = 0$ for each α . We shall denote the function algebra T^0 by A , and the collection of vector fields $L = l_1 \nabla^1 \in T^1$ by D . Let L^N be the subcollection of T^N generated by taking sums of products of functions in A and vector fields in D . Finally, let $A_0 = T_0^0$, $D_0 = D \cap T_0^1$, and $L_0^N = L^N \cap T_0^N$. Note that the T^N and L^N form graded algebras with respect to ordinary operator product, and that D is a Lie algebra under commutator product with D_0 as an ideal. Also note that all classes defined are A -modules.

For $L = \sum_{\alpha=0}^N l_\alpha \nabla^\alpha \in T^N$, we define the formal adjoint by $L' = \sum_{\alpha=0}^N (-1)^\alpha \nabla^\alpha \bar{l}_\alpha$ where the operator $\nabla^\alpha \bar{l}_\alpha$ is defined analogously to (1.1):

$$(1.5) \quad \nabla^\alpha \bar{l}_\alpha u = \nabla_{i_1} \cdots \nabla_{i_\alpha} (\overline{l_\alpha^{i_1 \cdots i_\alpha}} u)$$

for $u \in C_0^\infty(\Omega)$. Using the Leibniz rule for covariant derivatives, note that $L' = (-1)^N \bar{L} +$ lower order terms $\in T^N$. Also note that $L \in L^N$ (resp. T_0^N) implies $L' \in L^N$ (resp. T_0^N). The formal adjoint has the property that for $u, v \in C_0^\infty(\Omega)$ we have $\langle Lu, v \rangle = \langle u, L'v \rangle$ (where $\langle \cdot, \cdot \rangle$ denotes the L^2 inner product). Thus, using self-adjointness of A , we have $\langle LA^N u, v \rangle = \langle u, A^N L'v \rangle$ for $v \in C_0^\infty(\Omega)$ and $u \in A^{-N} C_0^\infty(\Omega)$. Since both $C_0^\infty(\Omega)$ and $A^{-N} C_0^\infty(\Omega)$ are dense in \mathfrak{S} we obtain the following from Theorem 1.1.

COROLLARY 1.2. *If $L \in T^N$, then $LA^N, A^N L \in \mathcal{L}(\mathfrak{S})$ and $(LA^N)^* = A^N L'$ (where $*$ denotes the Hilbert space adjoint).*

We next begin investigating commutators. Let $L^N + T_0^N$ denote the algebraic sum of those two classes of operators.

LEMMA 1.3. *If $L \in L^{N_1} + T_0^{N_1}$ and $M \in L^{N_2} + T_0^{N_2}$ then $[L, M] \in T_0^{N_1+N_2-1}$ and $[L, A] \in T_0^{N_1+1}$.*

Proof. If $L \in T_0^{N_1}$ or $M \in T_0^{N_2}$, then $[L, M] \in T_0^{N_1+N_2-1}$ may easily be seen by collecting top-order terms. Hence we may assume $L \in L^{N_1}$ and $M \in L^{N_2}$. The proof of the first statement then follows using $[D_1, D_2] \in D_0$ and $[L, DM] = [L, D]M + D[L, M]$ for $D, D_1, D_2 \in D$. To prove the second statement we may assume $L \in L^{N_1}$, and has top terms only: $L = D_1 \cdots D_{N_1}$ where $D_j \in D$. Then $[L, A] = [D_1 \cdots D_{N_1}, A] = D_1 \cdots D_{N_1-1} [D_{N_1}, A] + [D_1 \cdots D_{N_1-1}, A] D_{N_1}$ so the second

statement follows provided we can show $[D, A] \in T_0^2$ for every $D \in D$. But letting $D = b^i \nabla_i$, a computation in local coordinates shows

$$\begin{aligned}
 [D, A] &= [b^i \nabla_i, g^{jk} \nabla_j \nabla_k] \\
 &\equiv b^i g^{jk} \nabla_i \nabla_j \nabla_k - b^i g^{jk} \nabla_j \nabla_k \nabla_i \pmod{T_0^2} \\
 &= b^i g^{jk} R_{kj}^i \nabla_l \\
 &= b^i g^{lp} R_{ipk}^k \nabla_l \\
 &= -g^{lp} (\nabla_k \nabla_p b^k - \nabla_p \nabla_k b^k) \nabla_l \\
 &\equiv 0 \pmod{T_0^2}.
 \end{aligned}
 \tag{1.6}$$

This lemma and Theorem 1.1 enable us to investigate commutators of certain bounded operators.

LEMMA 1.4. *If $L \in L^N + T_0^N$, then $[LA^N, A] = K_1 A = A K_2$ where $K_j \in \mathcal{K}(\mathfrak{S})$.*

Proof. If we let $R(\lambda) = (\lambda - A^2)^{-1}$ and $\Gamma = \{z \in \mathbb{C}: |z - 1| = 1\}$ (with positive orientation), then

$$[LA^N, A] = \frac{1}{2\pi i} \int_{\Gamma} \sqrt{\lambda} [LA^N, R(\lambda)] d\lambda
 \tag{1-7}$$

and

$$\begin{aligned}
 [LA^N, R(\lambda)] &= R(\lambda) [LA^{N-1}, A^2] R(\lambda) A \\
 &= R(\lambda) A^2 [L, A] A^{N+1} R(\lambda) A \\
 &= A R(\lambda) K R(\lambda) A
 \end{aligned}
 \tag{1.8}$$

where $K = A[L, A]A^{N+1} \in \mathcal{K}$ by Theorem 1.1 and Lemma 1.3. Let $K_1(\lambda) = A R(\lambda) K R(\lambda)$ and $K_2(\lambda) = R(\lambda) K R(\lambda) A$, so $K_j(\lambda)$ defines a norm continuous map from $\Gamma \setminus \{0\}$ to \mathcal{K} . Using (4.1) of [5] with $s = 1$ shows that $K_j(\lambda) = O(|\lambda|^{-3/4})$. So $K_j = (2\pi i)^{-1} \int_{\Gamma} \sqrt{\lambda} K_j(\lambda) d\lambda \in \mathcal{K}$ for $j = 1, 2$ and the lemma is proved.

The above lemma enables us to prove the following useful result.

PROPOSITION 1.5. *If $L \in L^N + T_0^N$ and P and Q are nonnegative integers such that $P + Q = N$, then*

(i) $A^P L A^Q \in \mathcal{L}(\mathfrak{S})$ and (ii) $A^P L A^Q - L A^N \in \mathcal{K}(\mathfrak{S})$. *In addition, if $L \in T_0^N$ then (iii) $A^{P+1} L A^Q \in \mathcal{K}(\mathfrak{S})$.*

Proof. We perform an induction on P . The case $P = 0$ is just Theorem 1.1: for $N \geq 0$ and $L \in L^N + T_0^N$ we have (i) $L A^N \in \mathcal{L}(\mathfrak{S})$, (ii) $L A^N - L A^N \in \mathcal{K}$, and (in case $L \in T_0^N$) (iii) $A L A^N \in \mathcal{K}$. Now

assume by induction that the proposition is true for P . For $N \geq P$, let $L \in \mathbf{L}^N + \mathbf{T}_0^N$ and $Q = N - P$. Then

$$(1.9) \quad [A^P L A^Q, A] = (2\pi i)^{-1} \int_r \sqrt{\lambda} [A^P L A^Q, R(\lambda)] d\lambda$$

and

$$(1.10) \quad \begin{aligned} [A^P L A^Q, R(\lambda)] &= R(\lambda) A^P [L A^{Q-1}, A^2] R(\lambda) \\ &= A R(\lambda) A^{P+1} [L, A] A^{Q+1} R(\lambda) \\ &= A R(\lambda) K R(\lambda) A \end{aligned}$$

where Γ and $R(\lambda)$ are as in the preceding proof and $K = A^{P+1} [L, A] A^{Q+1}$. But by Lemma 1.3, $[L, A] \in \mathbf{T}_0^{N+1}$, so $K = A(A^P [L, A] A^{Q+1}) \in \mathcal{K}$ by the induction hypothesis (for P and $N+1$). So if we let $K_1(\lambda) = A R(\lambda) K R(\lambda)$ and $K_2 = R(\lambda) K R(\lambda) A$, then $K_j(\lambda) = O(|\lambda|^{-3/4})$ and $[A^P L A^Q, A] = K_1 A = A K_2$ where $K_j = (2\pi i)^{-1} \int_r K_j(\lambda) d\lambda \in \mathcal{K}$ for $j = 1, 2$. Since $\mathfrak{S}_1 = A\mathfrak{S}$ is dense in \mathfrak{S} , this implies

$$(1.11) \quad A^P L A^Q - A^{P+1} L A^{Q-1} \in \mathcal{K}.$$

Using the induction hypothesis, we get (i) for $P+1$. In fact the induction hypothesis also yields

$$(1.12) \quad A^P L A^Q - L A^N \in \mathcal{K}$$

which together with (1.11) yields (ii) for $P+1$. Finally, if $L \in \mathbf{T}_0^N$, multiplying (1.11) on the left by A and invoking the case P once more yields (iii) for $P+1$.

COROLLARY 1.6. *If $L \in \mathbf{L}^N + \mathbf{T}_0^N$, then $L A^N - A^N L \in \mathcal{K}$.*

Let \mathfrak{U}^0 denote the algebra of bounded operators on \mathfrak{S} which is finitely generated by $L A^N$ and $A^N L$ for $L \in \mathbf{L}^N + \mathbf{T}_0^N$. Note that \mathfrak{U}^0 contains A and multiplications by functions in \mathbf{A} . Let \mathfrak{U} be the norm closure of the collection $\mathfrak{U}^0 + \mathcal{K}$. We may now prove the following.

THEOREM 1.7. *\mathfrak{U} is a C^* -subalgebra of $\mathcal{L}(\mathfrak{S})$ with compact commutators.*

Proof. \mathfrak{U} is closed under adjoints by Corollary 1.2. Concerning commutators, let $L_1 \in \mathbf{L}^{N_1} + \mathbf{T}_0^{N_1}$ and $L_2 \in \mathbf{L}^{N_2} + \mathbf{T}_0^{N_2}$ and suppose $N_1 \geq N_2$. Then $L_2 \in \mathbf{L}^{N_1} + \mathbf{T}_0^{N_1}$ and

$$\begin{aligned} L_1 A^{N_1} L_2 A^{N_2} &\equiv L_1 L_2 A^{N_1} A^{N_2} \equiv A^{N_1} A^{N_2} L_1 L_2 \equiv A^{N_2} A^{N_1} L_2 L_1 \\ &\equiv A^{N_2} L_2 A^{N_1} L_1 \equiv L_2 A^{N_2} L_1 A^{N_1} \pmod{\mathcal{K}} \end{aligned}$$

using Lemma 1.3 and Corollary 1.6. By Corollary 1.6 this takes care of all commutators.

We shall call \mathfrak{A} the C^* -algebra for singular elliptic theory on Ω . The reason for this is as follows. By Theorem 1.7, $\mathfrak{A}/\mathfrak{K}$ is a commutative C^* -algebra; hence by the Gel'fand theory it is isometrically $*$ -isomorphic to $C(\mathbf{M})$, the algebra of continuous functions on the space \mathbf{M} of maximal ideals in $\mathfrak{A}/\mathfrak{K}$. For $A \in \mathfrak{A}$, denote the image of its coset $A + \mathfrak{K}$ in $C(\mathbf{M})$ by σ_A . Since $\mathfrak{A}/\mathfrak{K}$ is a C^* -algebra, invertibility of $A + \mathfrak{K}$ in $\mathfrak{L}(\mathfrak{H})/\mathfrak{K}$ is equivalent to its invertibility in $\mathfrak{A}/\mathfrak{K}$, and hence to the nonvanishing of σ_A on \mathbf{M} . Thus a further analysis of σ_A and \mathbf{M} is desired to provide an algebraic criterion for A to be Fredholm. We turn to this question in the remaining sections.

2. Maximal ideals over finite points. Let $\mathfrak{A}_m \subset \mathfrak{A}$ denote the algebra of multiplication operators obtained by closing A under the uniform norm. Since \mathfrak{A}_m is a closed $*$ -subalgebra of all bounded continuous functions on Ω , \mathfrak{A}_m is isometrically $*$ -isomorphic to $C(\bar{\Omega})$ where $\bar{\Omega}$ is some compactification of Ω . Although $\bar{\Omega}$ has no simple geometrical structure, there is a continuous surjection $p: \mathbf{M} \rightarrow \bar{\Omega}$ defined as the associated dual map (c.f. [12]) for the continuous injection $\mathfrak{A}_m \rightarrow \mathfrak{A}/\mathfrak{K}$. Thus $S = p^{-1}(\Omega)$ is the set of maximal ideals over finite points of Ω . We can completely describe this part of \mathbf{M} and σ_A . If l_α is a contravariant tensor field of degree α , we shall denote by $l_\alpha \xi^\alpha$ the function defined on cotangent vectors (x, ξ) by $l_\alpha^{i_1 \dots i_\alpha}(x) \xi_{i_1} \dots \xi_{i_\alpha}$.

THEOREM 2.1. *S is homeomorphic to $S^* \Omega$, the bundle of unit cospheres, under the map $m \rightarrow (x, \xi)$ such that*

$$(2.1) \quad \sigma_{L, L^N}(m) = l_N(x) \xi^N$$

where $L = \sum_{\alpha=0}^N (-i)^\alpha l_\alpha \nabla^\alpha L^N + T_0^N$. Thus uniform ellipticity is a necessary condition for $L: \mathfrak{S}_N \rightarrow \mathfrak{S}$ to be Fredholm. (Note that as special cases of (2.1) we have $\sigma(m) = 0$, and $\sigma_a(m) = a(x)$ for all $a \in A$.)

Proof. For $m \in S$ with $p(m) = x \in \Omega$, let x^1, \dots, x^n be local coordinates in a chart U which diagonalizes the metric at x . Let N be a compact neighborhood of x such that $N \subset U$, and let $\phi \in C_0^\infty(U)$ satisfying $0 \leq \phi \leq 1$ on U and $\phi = 1$ on N . Note that for each ϕ , $S_\nu = -i\phi(\partial/\partial x^\nu)A \in \mathfrak{A}$ and $S_\nu^* \equiv S_\nu \pmod{\mathfrak{K}}$. Hence $\xi_\nu = \sigma_{S_\nu}(m)$ is real for each ν and (x, ξ) where $\xi = \sum_{\nu=1}^n \xi_\nu dx^\nu$ defines an element of the cotangent space at x . Furthermore

$$\begin{aligned}
 \sum_{\nu, \mu} S_\nu^* g^{\nu\mu} S_\mu &= \sum_{\nu, \mu} \Lambda(\partial/\partial x^\nu) \phi g^{\nu\mu} \phi(\partial/\partial x^\mu) \Lambda \\
 (2.2) \qquad \qquad \qquad &\equiv -\Lambda \phi^2 \Lambda (\text{mod } \mathcal{H}) \\
 &\equiv -\phi^2 \Lambda \Lambda (\text{mod } \mathcal{H}) \\
 &\equiv \phi^2 (\text{mod } \mathcal{H}) .
 \end{aligned}$$

So

$$\begin{aligned}
 |\xi|^2 &= \sum \xi_\nu^2 = \sum_{\nu, \mu} \sigma_{S_\nu}(m) g^{\nu\mu}(x) \sigma_{S_\mu}(m) \\
 (2.3) \qquad \qquad \qquad &= \sigma_{(\sum_{\nu, \mu} S_\nu^* g^{\nu\mu} S_\mu)}(m) \\
 &= \phi^2(x) = 1 .
 \end{aligned}$$

Property (i) is immediate from the definition of the surjection $p: M \rightarrow \bar{\Omega}$ while (ii) follows from $\sigma_\Lambda(m) = \sigma_{\phi\Lambda}(m) = 0$ since $\phi(x) = 1$ and $\phi\Lambda \in \mathcal{H}$. The formula (2.1) holds for $N = 1$ and $L = D = -ib^\nu(\partial/\partial x^\nu) \in D$ since

$$\begin{aligned}
 \sigma_{D\Lambda}(m) &= \sigma_{\phi^2 D\Lambda}(m) \\
 &= \sigma_{(\sum_{\nu=1}^n (\phi b^\nu)(-i\phi \partial/\partial x^\nu)\Lambda)}(m) \\
 &= \sum_{\nu=1}^n b^\nu(x) \xi_\nu
 \end{aligned}$$

and this may be extended to higher-order tensors: for example $L = -l^{\nu\mu} \nabla_\nu \nabla_\mu \in L^2 + T_0^2$ implies

$$\phi^2 L \Lambda^2 \equiv \sum_{\nu, \mu} (-\phi l^{\nu\mu} \nabla_\nu)(\phi \nabla_\mu) \Lambda^2 \equiv \sum_{\nu, \mu} (-\phi l^{\nu\mu} \nabla_\nu \Lambda)(\phi \nabla_\mu \Lambda) (\text{mod } \mathcal{H})$$

so

$$\begin{aligned}
 \sigma_{L\Lambda^2}(m) &= \sigma_{\phi^2 L\Lambda^2}(m) \\
 (2.5) \qquad \qquad \qquad &= \sum_{\nu, \mu} (-il^{\nu\mu}(x) \xi_\nu)(i\xi_\mu) \\
 &= l^{\nu\mu}(x) \xi_\nu \xi_\mu .
 \end{aligned}$$

The map $m \rightarrow (x, \xi)$ is clearly 1-1. To show it is surjective onto $S^*\bar{\Omega}$, we use the following lemma (whose proof is immediate) to achieve local rotations of M .

LEMMA 2.2. *Let $\phi \in \mathfrak{A}_m$ be real-valued satisfying $0 \leq \phi \leq 1$ on Ω and $\phi = 1$ on a closed set $U \subset \bar{\Omega}$, and consider the C^* -algebra $\mathfrak{A}_\phi \subset \mathfrak{A}$ generated by \mathcal{H} , $a \in \mathbf{A}$, $\phi\Lambda$, and $\phi^N L \Lambda^N: L \in L^N + T_0^N$. Let M_ϕ be the maximal ideal space of $\mathfrak{A}_\phi/\mathcal{H}$, $\sigma^\phi: \mathfrak{A}_\phi \rightarrow C(M_\phi)$ the symbol homomorphism, $p_\phi: M_\phi \rightarrow \bar{\Omega}$ the associated dual map to the inclusion $\mathfrak{A}_m \subset \mathfrak{A}_\phi/\mathcal{H}$. Let $N_\phi = p_\phi^{-1}(U) \subset M_\phi$ whereas $N = p^{-1}(U) \subset M$. Defining $\rho(a + \mathcal{H}) = a + \mathcal{H}$ and $\rho(L\Lambda^N + \mathcal{H}) = \phi^N L \Lambda^N + \mathcal{H}$ yields a surjective algebra *-homomorphism $\rho: \mathfrak{A}/\mathcal{H} \rightarrow \mathfrak{A}_\phi/\mathcal{H}$ whose associated dual map*

provides an injection $j: M_\phi \rightarrow M$ such that

$$\begin{array}{ccc}
 M_\phi & \xrightarrow{j} & M \\
 & \searrow p_\phi & \swarrow p \\
 & & \bar{\Omega}
 \end{array}$$

commutes and

$$(2.7) \quad \sigma_A(j(n)) = \sigma_B^\phi(n)$$

for all $n \in M_\phi$, $A \in \mathfrak{A}$, and $B \in \rho(A + \mathcal{H})$. Furthermore, the restriction of j to N_ϕ provides a homeomorphism onto N , so we may consider $N_\phi \subset M$ and $N \subset M_\phi$.

Applying the lemma in our case, since ϕ has compact support in a coordinate chart, the algebra \mathfrak{A}_ϕ is generated by $a \in \mathfrak{A}$, $S_\nu, \nu = 1, \dots, n$, and \mathcal{H} . Choose any $\xi' \in S_x^* \Omega$ and let $R = (r^{\nu\mu})$ be an orthogonal matrix such that $\xi' = R\xi$. Defining $\tau(a + \mathcal{H}) = a + \mathcal{H}$ and $\tau(S_\nu + \mathcal{H}) = T_\nu + \mathcal{H}$ where $T_\nu = -i\phi \sum_{\mu=1}^n r^{\nu\mu} (\partial/\partial x^\mu) \Lambda$, we obtain a surjective *-isomorphism $\tau: \mathfrak{A}_\phi/\mathcal{H} \rightarrow \mathfrak{A}_\phi/\mathcal{H}$ whose associated dual map must be a (surjective) homeomorphism $h: M_\phi \rightarrow M_\phi$ satisfying $\sigma_A^\phi(h(n)) = \sigma_B^\phi(n)$ for all $A \in \mathfrak{A}_\phi$, $n \in M_\phi$, and $B \in \tau(A + \mathcal{H})$. In particular, h preserves fibers over U . Thus $m' = j(h(m)) \in M$ has the desired property that $m' \rightarrow (x, \xi)$ under the correspondence of the theorem since $\xi'_\nu = \sum_{\mu=1}^n r^{\nu\mu} \xi_\mu = \sigma_{S_\nu}^\phi(h(m)) = \sigma_{S_\nu}(j(h(m)))$. So we need only check continuity. But the topology of M is defined so that each σ_A is continuous, and since the functions (2.1) are continuous on $S^* \Omega$ (with respect to the usual cosphere bundle topology) and these functions separate points of $S^* \Omega$, we must have the map $m \rightarrow (x, \xi)$ continuous. In a similar manner we have the inverse mapping continuous.

Theorem 2.1 is analogous to the result for compact Ω proved by Seeley [13]. The proof is similar to that in [13], except regarding the surjectivity of the correspondence: we have used associated dual maps, ideas generalized to manifolds from Herman [8], instead of the Gohberg type estimates invoked by Seeley. In fact, it should be pointed out that the validity of Theorem 2.1 depends only on the generators of the algebra at finite points of Ω (an idea expressed more precisely in Lemma 2.2). Hence Theorem 2.1 would remain true if we were to allow generators of \mathfrak{A} which do not behave asymptotically like products of vector fields in D . However, to complete the analysis at hand, we must analyze the asymptotic behavior of D which we take up in the next section.

3. The asymptotic behavior of D . We are interested in finding a set of generators of D at "infinite points of $\bar{\Omega}$." Let $\partial\bar{\Omega} = \bar{\Omega} \setminus \Omega$, but recall that this set only has meaning as a collection of maximal ideals for the algebra \mathfrak{A}_m . However, if $D_1, D_2 \in D$ then their pointwise inner product under the metric $g_{i,j}$ defines a function $g(D_1, D_2)$ which, by the Leibniz rule for covariant derivatives, is in A . Thus $g(D_1, D_2)(x)$ for $x \in \partial\bar{\Omega}$ is defined.

DEFINITION. We shall call $D_1, \dots, D_\kappa \in D$ an *orthonormal basis* for D at $x \in \partial\bar{\Omega}$ if

$$(i) \quad g(D_i, D_j)(x) = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

(ii) for every $D \in D$ there exists constants c_1, \dots, c_κ such that the function $|D - \sum c_i D_i|^2 \in A$ vanishes at x .

Using a Gram-Schmidt type of procedure, it is not hard to verify that there exists an orthonormal basis for D at each $x \in \partial\bar{\Omega}$, and that each basis contains the same number of vector fields, $\kappa = d(x)$, the "dimension" of D at x . Note that $\kappa \leq n$.

We would like to be able to extend the notion of orthonormal bases to open sets in $\partial\bar{\Omega}$. In order to do so we need the following condition.

Condition C. The function $d: \partial\bar{\Omega} \rightarrow Z_+$ is continuous. I.e., the dimension of D is constant on connected components of $\partial\bar{\Omega}$.

We may now prove the following.

THEOREM 3.1. Under Condition C, for every $x_0 \in \partial\bar{\Omega}$ we can find an open set $U_0 \subset \bar{\Omega}$ and real vector fields $D_1, \dots, D_\kappa \in D$ ($\kappa = d(x_0)$) such that

$$(i) \quad x_0 \in U_0$$

$$(ii) \quad g(D_i, D_j)(x) = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases} \text{ for every } x \in U_0$$

(iii) for every $D \in D$ we may find functions $a_1, \dots, a_\kappa \in A$ such that $D = a_1 D_1 + a_2 D_2 + \dots + a_\kappa D_\kappa + D_0$ everywhere on U_0 , where $D_0 \in D$ satisfies $g(D_0, D_0)(x) = 0$ for all $x \in \partial\bar{\Omega} \cap U_0$.

Proof. Pick an orthonormal basis at $x_0: \tilde{D}_1, \dots, \tilde{D}_\kappa \in D$. Without loss of generality we may assume $\tilde{D}_1, \dots, \tilde{D}_\kappa$ are all real vector

fields (since otherwise we may orthogonalize the real and imaginary parts at x_0). Since $g(\tilde{D}_i, \tilde{D}_j) \in \mathcal{A}$ for all i, j , there exists an open set $U_0 \subset \bar{\Omega}$ containing x_0 such that $\partial\bar{\Omega} \cap U_0$ is connected and

$$(3.1) \quad g(\tilde{D}_i, \tilde{D}_j)(x) = \begin{cases} > 3/4 & \text{if } i = j \\ < 1/2\kappa & \text{if } i \neq j \end{cases}$$

for all $x \in U_0$. Let $\chi \in C^\infty(\mathbf{R})$ with all derivatives bounded, $|\chi(t)| \leq 1$ for all t , and

$$(3.2) \quad \chi(t) = \begin{cases} 1 & \text{if } t > 1/3 \\ 0 & \text{if } t < 1/4. \end{cases}$$

Let $\phi_1 = \chi(g(\tilde{D}_1, \tilde{D}_1))$. It is easy to see that $\phi_1(g(\tilde{D}_1, \tilde{D}_1))^{-1/2} \in \mathcal{A}$ so $D_1 = \phi_1(g(\tilde{D}_1, \tilde{D}_1))^{-1/2} \tilde{D}_1 \in \mathbf{D}$ and satisfies $g(D_1, D_1) = 1$ on U_0 . Inductively, define $D_j = \phi_j(g(B_j, B_j))^{-1/2} B_j$ where $\phi_j = \chi(g(B_j, B_j))$ and $B_j = \tilde{D}_j - g(D_1, \tilde{D}_j)D_1 - \dots - g(D_{j-1}, \tilde{D}_j)D_{j-1}$. A computation shows

$$(3.3) \quad \begin{aligned} g(B_j, B_j) &= g(\tilde{D}_j, \tilde{D}_j) - g(D_1, \tilde{D}_j)^2 - \dots - g(D_{j-1}, \tilde{D}_j)^2 \\ &> 3/4 - 1/3\kappa - 1/3\kappa - \dots - 1/3\kappa > 1/3 \end{aligned}$$

for $x \in U_0$, so $D_j \in \mathbf{D}$. Hence D_1, \dots, D_κ is orthogonal at every $x \in U_0$, i.e., we have (ii). Now let $a_j = g(D, D_j) \in \mathcal{A}$, and let $D_0 = D - \sum_{j=1}^\kappa a_j D_j$. Then

$$(3.4) \quad g(D_0, D_j) = g(D - \sum a_i D_i, D_j) = a_j - a_j = 0$$

implies D_0 is never a linear combination of D_1, \dots, D_κ on U_0 . Hence by Condition C and the connectedness of $\partial\bar{\Omega} \cap U_0$, we must have $g(D_0, D_0)(x) = 0$ for all $x \in \partial\bar{\Omega} \cap U_0$.

4. The formal algebra symbol and maximal ideals at infinity. For $L = \sum_{\alpha=0}^N l_\alpha \mathcal{F}^\alpha \in \mathbf{T}^N$, let us define the *formal symbol*

$$(4.1) \quad \tilde{\sigma}_{L \mathcal{A}^N}(x, \xi) = \left(\sum_{\alpha=0}^N l_\alpha(x) \xi^\alpha \right) (1 + |\xi|^2)^{-N/2} = \tilde{\sigma}_{\mathcal{A}^N L}(x, \xi)$$

which is a continuous function on the cotangent bundle $T^*\Omega$. (Recall from §2 that the notation $l_\alpha^{\xi^\alpha}$ means the function $l_\alpha^{i_1 \dots i_\alpha}(x) \xi_{i_1} \dots \xi_{i_\alpha}$.) Note that the boundedness of the coefficient tensors of L implies $\tilde{\sigma}_{L \mathcal{A}^N}$ is bounded on $T^*\Omega$. The formal symbols for operators $L \mathcal{A}^N$ and $\mathcal{A}^N L$ in \mathfrak{Q}^0 generate a C^* -subalgebra \mathcal{F} of the algebra of bounded continuous functions on $T^*\Omega$. Hence \mathcal{F} is isometrically $*$ -isomorphic to $C(P^*\Omega)$, where $P^*\Omega$ denotes some compactification of $T^*\Omega$. The injection $\mathfrak{Q}_m \rightarrow \mathcal{F}$ yields a continuous surjection $\pi: P^*\Omega \rightarrow \bar{\Omega}$. Unfortunately the map $\tilde{\sigma}: \mathfrak{Q}^0 \rightarrow C(P^*\Omega)$ is not an algebra homomorphism: for example $\tilde{\sigma}_L \tilde{\sigma}_{D_1 D_2 A^2} = (b^i c^j \xi_i \xi_j + b^i c^j, \xi_j) (1 + |\xi|^2)^{-3/2}$

but $\tilde{\sigma}_{AD_1}\tilde{\sigma}_{D_2A^2} = (b^i c^j \xi_i \xi_j)(1 + |\xi|^2)^{-3/2}$ where $D_1 = b^i \nabla_i$, $D_2 = c^j \nabla_j \in \mathbf{D}$. However, if we restrict our functions to the subset $\partial P^* \Omega = P^* \Omega \setminus T^* \Omega$, we obtain an algebra homomorphism $\dot{\sigma}: \mathfrak{A}^0 \rightarrow C(\partial P^* \Omega)$ and we call $\dot{\sigma}_A = \tilde{\sigma}_A|_{\partial P^* \Omega}$ the *formal algebra symbol* of $A \in \mathfrak{A}^0$.

It is evident that the cosphere bundle $S^* \Omega$ is homeomorphic to $\pi^{-1}(\Omega)$ by the map $(x, \xi) \rightarrow \lim_{r \rightarrow \infty} (x, r\xi) \in \partial P^* \Omega$. Theorem 2.1 may be interpreted as providing a continuous injection $\theta: \mathbf{S} \rightarrow \partial P^* \Omega$ such that

$$(4.2) \quad \dot{\sigma}_A(\theta(m)) = \sigma_A(m)$$

for $m \in \mathbf{S}$ and operators $A \in \mathfrak{A}^0$. The main result of this paper extends this results as follows.

THEOREM 4.1. *Under Condition C, there exists continuous injection $\theta: \mathbf{M} \rightarrow \partial P^* \Omega$ such that*

$$(4.3) \quad \begin{array}{ccc} \mathbf{M} & \xrightarrow{\theta} & \partial P^* \Omega \\ & \searrow p & \swarrow \pi \\ & \bar{\Omega} & \end{array}$$

is commutative, and (4.2) holds for all $m \in \mathbf{M}$ and $A \in \mathfrak{A}^0$. Thus a sufficient condition that $L: \mathfrak{S}_N \rightarrow \mathfrak{S}$ be Fredholm is that $\dot{\sigma}_{LA^N} \neq 0$ on $\partial P^* \Omega$.

Before extending θ to $\mathbf{M}_\infty = \mathbf{M} \setminus \mathbf{S}$, we first must analyze $C(\mathbf{M})$ a little more closely. Let \mathbf{M}_p denote the closure in \mathbf{M} of \mathbf{S} and $\mathbf{M}_s = \mathbf{M} \setminus \mathbf{M}_p$. We may now prove the following.

LEMMA 4.2. $\mathbf{M}_p = \{m \in \mathbf{M}: \sigma_A(m) = 0\}$.

Proof. By Theorem 2.1 it is clear that $\mathbf{M}_p \subset \{m \in \mathbf{M}: \sigma_A(m) = 0\}$. Let $\mathcal{F} = \{A \in \mathfrak{A}: \sigma_A = 0 \text{ on } \mathbf{M}_p\}$. Then $\mathcal{F} / \mathcal{K}$ is a C^* -algebra (without unit) which is isomorphic to the algebra of continuous functions on \mathbf{M}_s which vanish at infinity. Since \mathcal{A} is the only generator of \mathfrak{A}^0 in \mathcal{F} , \mathcal{F} must be generated by operators of the form AA and AA where $A \in \mathfrak{A}^0$. So if $\sigma_A(m) = 0$ for $m \in \mathbf{M}_s$, then $\sigma_B(m) = 0$ for all $B \in \mathcal{F}$ which is impossible.

LEMMA 4.3. *If $L \in \mathbf{T}_0^N$, then $\sigma_{LA^N} = 0$ on \mathbf{M}_∞ .*

Proof. If $m \in \mathbf{M}_\infty \cap \mathbf{M}_p$, let m_β be a net in \mathbf{S} converging to m . Then using Theorem 2.1 we conclude that $\sigma_{LA^N}(m_\beta) \rightarrow 0 = \sigma_{LA^N}(m)$.

If $m \in \mathbf{M}_s$, then $\sigma_A(m) = 0$ by Lemma 4.2. But $\sigma_{LA^N}(m)\sigma_A(m) = \sigma_{LA^{N+1}}(m) = 0$ since $LA^{N+1} \in \mathcal{K}$; so $\sigma_{LA^N}(m) = 0$.

Now let $m_0 \in \mathbf{M}_\infty$ with $x_0 = p(m_0) \in \partial\bar{\Omega}$. By the methods used in the proof of Theorem 3.1, it is possible to construct (in addition to U_0) a closed neighborhood $U \subset U_0$ of x_0 , and $\phi \in \mathfrak{A}_m$ satisfying $\phi = 1$ on U , $0 \leq \phi \leq 1$ on Ω , and $\text{supp } \phi \subset U_0$. Let $\mathfrak{A}_\phi, p_\phi, \mathbf{M}_\phi, N_\phi$, and N all be as described in Lemma 2.2. Furthermore, let $U_\infty = \partial\bar{\Omega} \cap U$ and $N_\infty = p^{-1}(U_\infty)$. By Lemma 2.2 we will also consider $N_\infty \subset \mathbf{M}_\phi$. Let $D_1, \dots, D_\kappa \in D(\kappa = d(x_0))$ be real vector fields as in Theorem 3.1, with expressions in local coordinates $D_\nu = b_\nu^j(\partial/\partial x^j)$ for $\nu = 1, \dots, \kappa$. Let $S_\nu = -i\phi D_\nu A \in \mathfrak{A}_\phi$ for $\nu = 1, \dots, \kappa$. We define a contravariant tensor h^{jk} by

$$(4.4) \quad g^{jk} = \sum_{\nu=1}^{\kappa} b_\nu^j b_\nu^k + h^{jk}.$$

If (x, ξ) is a complex covector with $x \in U_0$ and $|\xi|^2 = 1$, then

$$\sum_{\nu=1}^{\kappa} b_\nu^j \xi_j b_\nu^k \bar{\xi}_k \leq 1$$

since the D_ν form an orthonormal set on U_0 . Hence $h^{jk} \xi_j \bar{\xi}_k \geq 0$, i.e., h^{jk} is positive semidefinite. We shall assume henceforth that h^{jk} is not identically zero, i.e., that $\kappa < n$. (See Remark 4.6 below for the case $\kappa = n$.) From (4.4) we obtain

$$(4.5) \quad \nabla_j \phi^2 g^{jk} \nabla_k = \sum_{\nu=1}^{\kappa} \nabla_j \phi^2 b_\nu^j b_\nu^k \nabla_k + \nabla_j \phi^2 h^{jk} \nabla_k.$$

Since

$$(4.6) \quad \begin{aligned} A \nabla_j \phi^2 g^{jk} \nabla_k A &\equiv A \phi^2 A A \pmod{\mathcal{K}} \\ &\equiv \phi^2 A A A \pmod{\mathcal{K}} \\ &= \phi^2 A^2 - \phi^2 \end{aligned}$$

and

$$(4.7) \quad \begin{aligned} A(\nabla_j \phi^2 b_\nu^j)(b_\nu^k \nabla_k)A &= -A(\nabla_j i\phi b_\nu^j)(i\phi b_\nu^k \nabla_k)A \\ &= -S_\nu^* S_\nu \\ &= -S_\nu^2 \pmod{\mathcal{K}} \end{aligned}$$

we get after multiplying (4.5) on the left and right by A and rearranging:

$$(4.8) \quad \phi^2 \equiv \phi^2 A^2 - A H A + \sum_{\nu=1}^{\kappa} S_\nu^2 \pmod{\mathcal{K}}$$

where H denotes the operator $\nabla_j \phi^2 h^{jk} \nabla_k$. The positive semi-definite-

ness of the tensor h^{jk} implies

$$(4.9) \quad \langle \Lambda H \Lambda u, u \rangle = - \int_{\Omega} \phi^2 h^{jk} (\nabla_k \Lambda u) (\overline{\nabla_j \Lambda u}) d\mu \leq 0$$

for $u \in C_0^\infty(\Omega)$, so the operator $\Lambda H \Lambda$ is nonpositive. Hence $\phi^2 \Lambda^2 - \Lambda H \Lambda$ is nonnegative and we may define $S_0 = (\phi^2 \Lambda^2 - \Lambda H \Lambda)^{-1/2}$. From (4.8) we conclude $\Lambda H \Lambda, S_0 \in \mathfrak{A}_\phi$ and

$$(4.10) \quad \phi^2 \equiv \sum_{\nu=0}^{\kappa} S_\nu^2 \pmod{\mathcal{N}}.$$

If we let $S_+^{\zeta} = \{\zeta = (\zeta_0, \dots, \zeta_\kappa) \in \mathbf{R}^{\kappa+1}; \sum_{\nu=0}^{\kappa} \zeta_\nu^2 = 1 \text{ and } \zeta_0 \geq 0\}$, then we are now able to prove the following.

PROPOSITION 4.4. *There exists a continuous injection*

$$(4.11) \quad \omega_1: N_\infty \longrightarrow U_\infty \times [0, 1] \times S_+^{\zeta}$$

such that $\omega_1(n) = (x, r, \zeta)$ satisfies (i) $\sigma_a(n) = a(x)$, (ii) $\sigma_\Lambda(n) = r$, (iii) $\sigma_{S_\nu}(n) = \zeta_\nu$ ($\nu = 0, 1, \dots, \kappa$), and (iv) $\zeta_0 \geq r$.

Proof. Since N_∞ is a closed subset of M_ϕ , $\mathcal{I}_\phi = \{A \in \mathfrak{A}_\phi: \sigma_A^\phi = 0 \text{ on } N_\infty\}$ is a closed ideal of \mathfrak{A}_ϕ , and $\mathfrak{A}_\phi/\mathcal{I}_\phi \simeq C(N_\infty)$. Furthermore, using $N_\infty \subset M_\infty$ and Lemma 4.3, $\mathfrak{A}_\phi/\mathcal{I}_\phi$ is generated by $a + \mathcal{I}_\phi$ ($a \in A$), $\phi A + \mathcal{I}_\phi$, and $S_\nu + \mathcal{I}_\phi$ ($\nu = 1, \dots, \kappa$). Thus if we let \mathcal{B}_m be the C^* -algebra generated by \mathfrak{A}_m and \mathcal{I}_ϕ , then $\mathcal{B}_m/\mathcal{I}_\phi$ is a commutative C^* -algebra isometrically $*$ -isomorphic to $C(U_\infty)$. Also if we let \mathfrak{A}_ϕ^* be the C^* -algebra generated by $\phi A, S_\nu$ ($\nu = 0, 1, \dots, \kappa$), and \mathcal{I}_ϕ ; then $\mathfrak{A}_\phi^*/\mathcal{I}_\phi$ is a commutative C^* -algebra (with unit using (4.8) and the fact that $\phi \equiv 1 \pmod{\mathcal{I}_\phi}$). In fact, since $\mathfrak{A}_\phi^*/\mathcal{I}_\phi$ is finitely generated, it has maximal ideal space N_∞^* homeomorphic to the joint spectrum of its generators (c.f. [12]). If we let $\sigma^*: \mathfrak{A}_\phi^* \rightarrow C(N_\infty^*)$ denote the symbol homomorphism, then (4.9) implies $\sigma_{\Lambda H \Lambda}^* \leq 0$, and so by (4.8), $0 \leq \sigma_{\phi^2 \Lambda^2}^* \leq 1$. By (4.10) we also have $\sum_{\nu=0}^{\kappa} (\sigma_{S_\nu}^*)^2 = 1$. Thus there is an injection

$$(4.12) \quad N_\infty^* \longrightarrow [0, 1] \times S_+^{\zeta}$$

where $n \rightarrow (r, \zeta)$ satisfies $\sigma_{\phi A}^*(n) = r$ and $\sigma_{S_\nu}^*(n) = \zeta_\nu$. Since $\mathfrak{A}_\phi/\mathcal{I}_\phi$ is generated by $\mathcal{B}_m/\mathcal{I}_\phi$ and $\mathfrak{A}_\phi^*/\mathcal{I}_\phi$, we can use Herman's Lemma (c.f. [8], Theorem 1) to conclude the existence of a continuous injection $N_\infty \rightarrow U_\infty \times N_\infty^*$ and composition with (4.12) yields a continuous injection $\omega_\phi: N_\infty \rightarrow U_\infty \times [0, 1] \times S_+^{\zeta}$ satisfying $\sigma_a^\phi(n) = a(x)$, $\sigma_{\phi A}^\phi(n) = r$, and $\sigma_{S_\nu}^\phi(n) = \zeta_\nu$. When we view $N_\infty \subset M$ (i.e., compose ω_ϕ with the homeomorphism $p^{-1}(U) \simeq p^{-1}(U)$), we obtain the map ω_ϕ of (4.11) satisfying (i)-(iii). Condition (iv) is simply a consequence of $\Lambda^2 \leq \Lambda^2 - \Lambda H \Lambda$.

In a similar manner we can let $P^*U = \pi^{-1}(U)$ and $P^*U_\infty = P^*U \cap P^*\Omega_\infty = \pi^{-1}(U_\infty)$. Applying the above argument to the function algebra \mathcal{F}_ϕ generated by the formal symbols of the generators of \mathfrak{A}_ϕ yields

PROPOSITION 4.5. *There exists a continuous injection*

$$(4.13) \quad \omega_2: P^*U_\infty \longrightarrow U_\infty \times [0, 1] \times S_+^n$$

such that for $\omega_2(p) = (x, r, \zeta)$ we have (i) $\dot{\sigma}_a(p) = a(x)$, (ii) $\dot{\sigma}_A(p) = r$, (iii) $\dot{\sigma}_{S_\nu}(p) = \zeta_\nu$ ($\nu = 0, 1, \dots, \kappa$), and (iv) $\zeta_0 \geq r$. In fact, every (x, r, ζ) in $U_\infty \times [0, 1] \times S_+^n$ satisfying (iv) is in the range of ω_2 .

Proof. To prove the last statement, let $x_\beta \in \Omega$ denote a net converging to x . Let $\xi_\beta \in \pi^{-1}(x_\beta)$ satisfy

$$(4.14) \quad b_\nu^j(\xi_\beta)_j r = \zeta_\nu \quad (\nu = 1, \dots, \kappa).$$

But this implies by (4.4) that

$$\{|\hat{\xi}_\beta|^2 - h^{ij}(\xi_\beta)_i(\xi_\beta)_j\}r^2 = \sum_{\nu=1}^{\kappa} \zeta_\nu^2$$

or

$$(4.15) \quad |\hat{\xi}_\beta|^2 = \frac{1}{r^2} - \frac{(\zeta_0)^2}{r^2} + h^{ij}(\xi_\beta)_i(\xi_\beta)_j.$$

The last term in (4.15) is nonnegative, so by adding something to ξ_β from the kernel of the system (4.14), which is nontrivial since we are assuming $\kappa < n$, we can also require

$$(4.16) \quad |\hat{\xi}_\beta|^2 = \frac{1}{r^2} - 1$$

provided $\zeta_0 \geq r$. (Note: if $r = 0$, take $\xi_\beta = \lim_{r \rightarrow \infty} r\xi'_\beta \in \pi^{-1}(x_\beta)$ where ξ'_β solves $b_\nu^j(\xi'_\beta)_j = \zeta_\nu$.) Observe that for each β , $\bar{\sigma}_A(x_\beta, \xi_\beta) = r$ and $\bar{\sigma}_{S_\nu}(x_\beta, \xi_\beta) = \zeta_\nu$ ($\nu = 1, \dots, \kappa$); hence also $[\bar{\sigma}_{S_0}(x_\beta, \xi_\beta)]^2 = 1 - \sum_{\nu=1}^{\kappa} \zeta_\nu^2 = \zeta_0^2$ for every β . Thus (x_β, ξ_β) must converge to a point $p \in P^*\Omega_\infty$ satisfying $\omega_2(p) = (x, r, \zeta)$.

These propositions easily imply our main result.

Proof of Theorem 4.1. Note that $\omega_2^{-1} \circ \omega_1$ provides a continuous injection $\theta: N_\infty \rightarrow P^*\Omega_\infty$. It is clear that the value $\theta(n) \in \partial P^*\Omega$ does not depend on the choice of neighborhood U , so we obtain a continuous injection $\theta: M_\infty \rightarrow \partial P^*\Omega$. Letting $m \in M_\infty \cap M_p$ and $m_\beta \in S$ denote a net converging to m , it is easy to check that $\dot{\sigma}_A(\theta(m)) =$

$\lim_{\beta} \dot{\sigma}_A(\theta(m_{\beta}))$ for every $A \in \mathcal{U}^0$, so we obtain a continuous injection $\theta: M \rightarrow \partial P^* \Omega$ satisfying (4.2).

REMARK 4.6. It was assumed above that $\kappa < n$. Actually, the case $\kappa = n$ is easier, for then we obtain from

$$(4.4') \quad g^{jk} = \sum_{\nu=1}^n b_j^{\nu} b_{\nu}^k$$

that

$$(4.10') \quad \begin{aligned} \phi^2 &\equiv \phi^2 A^2 + \sum_{\nu=1}^n S_{\nu}^2(\text{mod } \mathcal{K}) \\ &\equiv \sum_{\nu=0}^n S_{\nu}^2(\text{mod } \mathcal{K}) \end{aligned}$$

after defining $S_0 = \phi A$. In place of Proposition 4.4, the same proof establishes a continuous injection

$$(4.11') \quad \omega_1: N_{\infty} \longrightarrow U_{\infty} \times S_+^n$$

satisfying $\omega_1(n) = (x, \zeta)$ with $\sigma_a(n) = a(x)$ and $\sigma_{s_{\nu}}(n) = \zeta_{\nu}$ ($\nu = 0, \dots, n$). Similarly we get a continuous injection

$$(4.13') \quad \omega_2: P^* U_{\infty} \longrightarrow U_{\infty} \times S_+^n$$

satisfying $\omega_2(n) = (x, \zeta)$ with $\dot{\sigma}_a(n) = a(x)$ and $\dot{\sigma}_{s_{\nu}}(n) = \zeta_{\nu}$ ($\nu = 0, \dots, n$). In fact, imitating the proof of Proposition 4.5 establishes that (4.13') is surjective, and the proof of Theorem 4.1 follows as above.

REFERENCES

1. T. Aubin, *Espaces de Sobolev sur les varietes riemanniennes*, preprint.
2. M. Cantor, *Spaces of functions with asymptotic conditions on R^n* , Indiana Univ. Math. J., **24**, No. 9, (1975), 897-902.
3. H. O. Cordes, *Self-adjointness of powers of elliptic operators on noncompact manifolds*, Math. Ann., **195** (1972), 257-272.
4. H. O. Cordes and E. Herman, *Gel'fand theory of pseudo-differential operators*, Amer. J. Math., **90** (1968), 681-717.
5. H. O. Cordes and R. C. McOwen, *The C^* -algebra of a singular elliptic problem on a noncompact riemannian manifold*, Math. Zeit., **153** (1977), 101-116.
6. ———, *Remarks on singular elliptic theory for complete riemannian manifolds*, **70**, No. 1, (1977), 133-141.
7. L. P. Eisenhart, *Riemannian Geometry*, Princeton University Press, 1926.
8. E. Herman, *The symbol of the algebra of singular integral operators*, J. Math. Mech., **15** (1966), 147-156.
9. R. Jackson, *Canonical operators and lower-order symbols*, Amer. Math. Soc. Memoirs, (1973).
10. S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry*, Vol. I, New-York-London, Interscience, 1963.
11. R. C. McOwen, *The behavior of the laplacian on weighted Sobolev spaces*, Comm. Pure and App. Math., (to appear).

12. C. E. Rickart, *Banach Algebras*, Princeton, Van Nostrand, 1960.
13. R. T. Seeley, *Integro-differential operators on vector bundles*, Trans. Amer. Math. Soc., **117** (1965), 167-204.

Received April 11, 1979.

NEW YORK UNIVERSITY
COURANT INSTITUTE
NEW YORK, NY 10012

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE AND ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Spiros Argyros, <i>A decomposition of complete Boolean algebras</i>	1
Gerald A. Beer, <i>The approximation of upper semicontinuous multifunctions by step multifunctions</i>	11
Ehrhard Behrends and Richard Evans, <i>Multiplicity theory for Boolean algebras of L^p-projections</i>	21
Man-Duen Choi, <i>The full C^*-algebra of the free group on two generators</i>	41
Jen-Chung Chuan, <i>Axioms for closed left ideals in a C^*-algebra</i>	49
Jo-Ann Deborah Cohen, <i>The strong approximation theorem and locally bounded topologies on $F(X)$</i>	59
Eugene Harrison Gover and Mark Bernard Ramras, <i>Increasing sequences of Betti numbers</i>	65
Morton Edward Harris, <i>Finite groups having an involution centralizer with a 2-component of type $\text{PSL}(3, 3)$</i>	69
Valéria Botelho de Magalhães Iório, <i>Hopf C^*-algebras and locally compact groups</i>	75
Roy Andrew Johnson, <i>Nearly Borel sets and product measures</i>	97
Lowell Edwin Jones, <i>Construction of Z_p-actions on manifolds</i>	111
Manuel Lerman and Robert Irving Soare, <i>d-simple sets, small sets, and degree classes</i>	135
Philip W. McCartney, <i>Neighborly bushes and the Radon-Nikodým property for Banach spaces</i>	157
Robert Colman McOwen, <i>Fredholm theory of partial differential equations on complete Riemannian manifolds</i>	169
Ernest A. Michael and Carl Preston Pixley, <i>A unified theorem on continuous selections</i>	187
Ernest A. Michael, <i>Continuous selections and finite-dimensional sets</i>	189
Vassili Nestoridis, <i>Inner functions: noninvariant connected components</i>	199
Bun Wong, <i>A maximum principle on Clifford torus and nonexistence of proper holomorphic map from the ball to polydisc</i>	211
Steve Wright, <i>Similarity orbits of approximately finite C^*-algebras</i>	223
Kenjiro Yanagi, <i>On some fixed point theorems for multivalued mappings</i>	233
Wieslaw Zelazko, <i>A characterization of LC-nonremovable ideals in commutative Banach algebras</i>	241