

Pacific Journal of Mathematics

**CONTINUOUS SELECTIONS AND FINITE-DIMENSIONAL
SETS**

ERNEST A. MICHAEL

CONTINUOUS SELECTIONS AND FINITE-DIMENSIONAL SETS

E. MICHAEL

Some known selection theorems are strengthened by weakening the hypotheses on a finite-dimensional subset of the domain.

1. **Introduction.** The following selection theorem was recently obtained in [7] by C. Pixley and the author.

THEOREM 1.1. *Let X be paracompact, Y a Banach space, $Z \subset X$ with $\dim_x Z \leq 0$, and $\phi: X \rightarrow \mathcal{F}(Y)$ l.s.c. with $\phi(x)$ convex for all $x \in X - Z$. Then ϕ has a selection.*

Let us quickly define our terms: We write $2^Y = \{S \subset Y: S \neq \emptyset\}$ and $\mathcal{F}(Y) = \{S \in 2^Y: S \text{ closed in } Y\}$. A map $\phi: X \rightarrow 2^Y$ is *lower semi-continuous*, or l.s.c., if $\{x \in X: \phi(x) \cap V \neq \emptyset\}$ is open in X for every open V in Y . A *selection* for a map $\phi: X \rightarrow 2^Y$ is a continuous $f: X \rightarrow Y$ such that $f(x) \in \phi(x)$ for every $x \in X$. Finally, if $Z \subset X$, then $\dim_x Z \leq n$ means that $\dim S \leq n$ for every subset S of Z which is closed in X .¹

The purpose of this paper is first to generalize Theorem 1.1 to the case where $\dim_x Z \leq n$, and then to prove two other theorems of this type in which different assumptions are made on $\phi(x)$ when $x \in X - Z$. Just like Theorem 1.1, all of these "hybrid" selection theorems reduce to known results when $Z = \emptyset$ or $Z = X$. Theorem 1.1, along with the special cases of Theorems 1.3 and 1.4 where $\dim Z = 0$, will be applied in [6].

In contrast to Theorem 1.1, the theorems to be proved in this paper all have both a local and a global version. To state these results succinctly, we introduce some more terminology: A map $\phi: X \rightarrow 2^Y$ has the *selection extension property*, or SEP, if, whenever $A \subset X$ is closed, every selection g for $\phi|_A$ extends to a selection for ϕ . If g only extends to a selection for $\phi|U$ for some neighborhood U of A in X , then we say that ϕ has the *selection neighborhood extension property*, or SNEP. As pointed out in [6], Theorem 1.1 can be strengthened to conclude that ϕ actually has the SEP.

Before stating Theorem 1.2, we recall the following concepts from [4, p. 565], where $n \geq -1$: A space S is C^n if every continuous image of an i -sphere ($i \leq n$) in S is contractible in S . A collec-

¹ Observe that, for normal X , $\dim_x Z \leq n$ is valid if either $\dim Z \leq n$ or $\dim X \leq n$.

tion \mathcal{S} of subsets of a metric space Y is *uniformly equi- LC^n* if to every $\varepsilon > 0$ corresponds a $\nu(\varepsilon) > 0$ such that, for every $S \in \mathcal{S}$, every continuous image of an i -sphere ($i \leq n$) in S of diameter $< \nu(\varepsilon)$ is contractible over a subset of S of diameter $< \varepsilon$. Since there is no (-1) -sphere, the above properties are always vacuously satisfied when $n = -1$, and hence Theorem 1.2 really generalizes Theorem 1.1.

THEOREM 1.2. *Let X be paracompact, Y a Banach space, $Z \subset X$ with $\dim_x Z \leq n + 1$, and $\phi: X \rightarrow \mathcal{F}(Y)$ l.s.c. with $\phi(x)$ convex for all $x \in X - Z$ and with $\{\phi(x): x \in Z\}$ uniformly equi- LC^n . Then ϕ has the SNEP. If, moreover, $\phi(x)$ is C^n for every $x \in Z$, then ϕ has the SEP.*

Theorem 1.2 reduces to [3, Theorem 3.2''] when $Z = \emptyset$, and it implies [4, Theorem 1.2] when $Z = X$ (more generally, when Z is open in X and $\phi(x)$ is a singleton for $x \in X - Z$).

In our next theorem, a metric space Y is called an AR (resp. ANR) if it is a retract (resp. neighborhood retract) of every metric space E containing it as a closed subset. A collection \mathcal{S} of subsets of a topological space Y is called *equi- LC^n in Y* if, for every $y \in Y$, every neighborhood V of y contains a neighborhood W of y such that, for any $S \in \mathcal{S}$, every continuous image of an i -sphere ($i \leq n$) in $W \cap S$ is contractible in $V \cap S$. It is easy to see that every uniformly equi- LC^n collection of subsets of a metric space Y is equi- LC^n in Y ; for a partial converse, see Lemma 6.1(b).

THEOREM 1.3. *Let X be paracompact, Y a complete metric ANR, $Z \subset X$ with $\dim_x Z \leq n + 1$, and $\phi: X \rightarrow \mathcal{F}(Y)$ l.s.c. with $\phi(x) = Y$ for $x \in X - Z$ and with $\{\phi(x): x \in Z\}$ equi- LC^n in Y . Then ϕ has the SNEP. If moreover, Y is an AR and $\phi(x)$ is C^n for every $x \in Z$, then ϕ has the SEP.*

Observe that, when $Z = \emptyset$, Theorem 1.3 reduces to the known result that every complete metric AR (resp. ANR) has the extension property (resp. neighborhood extension property) with respect to paracompact spaces (see, for instance, C. H. Dowker [1]).

Our next result reduces to [4, Theorem 1.2] when $Z = \emptyset$.

THEOREM 1.4. *Let X be paracompact, $A \subset X$ closed with $\dim_x(X - A) \leq n + 1$, $Z \subset X - A$ with $\dim_x Z \leq m + 1$ (where $m \leq n$), Y complete metric, and $\phi: X \rightarrow \mathcal{F}(Y)$ l.s.c. such that $\{\phi(x): x \in X - Z\}$ is equi- LC^n in Y and $\{\phi(x): x \in Z\}$ is equi- LC^m in Y .*

Then ϕ has the SNEP at A .² If, moreover, $\phi(x)$ is C^n for all $x \in X - Z$ and C^m for all $x \in Z$, then ϕ has the SEP at A .

The paper is arranged as follows. Two known lemmas on l.s.c. maps are stated in §2, some general results on selections are established in §3, and a result which is implicitly contained in [4] is recorded in §4. After these preliminaries, Theorems 1.2, 1.3 and 1.4 are then proved in §§5, 6 and 7, respectively.

2. Two lemmas on l.s.c. maps. The following two known lemmas will be applied in the sequel.

LEMMA 2.1 [3, Example 1.3]. *Let $\phi: X \rightarrow 2^Y$ be l.s.c., $A \subset X$ closed, and g a selection for $\phi|_A$. Define $\phi_g: X \rightarrow 2^Y$ by $\phi_g(x) = \phi(x)$ if $x \in X - A$ and $\phi_g(x) = \{g(x)\}$ if $x \in A$. Then ϕ_g is also l.s.c.*

Henceforth, we shall freely refer to the map ϕ_g defined in Lemma 2.1.

LEMMA 2.2 [4, Lemma 11.3]. *Let $\phi: X \rightarrow 2^Y$ be l.s.c. with Y a metric space, let $K \subset Y$ be compact, and let $\varepsilon > 0$. Then $\{x \in X: K \subset B_\varepsilon(\phi(x))\}$ ³ is open in Y .*

3. Two properties of set-valued maps. In this section we consider the following two properties of a map $\phi: X \rightarrow 2^Y$, where Y is a metric space.

(3.1) To every $\varepsilon > 0$ corresponds an $\alpha(\varepsilon) > 0$ with the following property: If $A \subset X$ is closed, g is a selection for $\phi|_A$, and $h: X \rightarrow Y$ is continuous with $d(h, \phi_g) < \alpha(\varepsilon)$, then g extends to a selection f for ϕ with $d(f, h) < \varepsilon$.⁴

(3.2) To every $\varepsilon > 0$ corresponds a $\beta(\varepsilon) > 0$ with the following property: If $A \subset X$ is closed, g is a selection for $\phi|_A$, $h: X \rightarrow Y$ is continuous with $d(h, \phi_g) < \beta(\varepsilon)$, and $\mu > 0$, then there exists a continuous $f: X \rightarrow Y$ with $d(f, h) < \varepsilon$ and $d(f, \phi_g) < \mu$.

Property (3.2) (often with $A = \emptyset$) is used—implicitly or explicitly—in the proofs of almost all general selection theorems known to the author, including those in this paper. It is clear that the slightly simpler property (3.1) (which is called the *selection approximation property*, or SAP, in [6, §5]) always implies (3.2); the

² i.e., ϕ satisfies the definition of the SNEP for this particular A .

³ $B_\varepsilon(S)$ denotes the ε -neighborhood of S .

⁴ Here d is the metric on Y , and $d(h, \phi_g) < \alpha$ denotes that $d(h(x), \phi_g(x)) < \alpha$ for all $x \in X$. Similarly for $d(f, h) < \varepsilon$, etc.

following result shows that, under mild restrictions, the two properties are actually equivalent.

PROPOSITION 3.3. *If Y is a complete metric space, and if $\phi(x)$ is closed in Y for all $x \in X$, then (3.2) implies (3.1). Moreover, one can take $\alpha(\varepsilon) = \beta(\varepsilon/2)$; in particular, if one can take $\beta(\infty) = \infty^5$ then one can also take $\alpha(\infty) = \infty$.*

Proof. Assume that ϕ satisfies (3.2) and let us show that it satisfies (3.1) with $\alpha(\varepsilon) = \beta(\varepsilon/2)$. Let $f_1 = h$. By induction, use (3.2) to construct continuous maps $f_n: X \rightarrow Y$ such that $d(f_n, \phi_a) < \beta(2^{-n}\varepsilon)$ and $d(f_{n+1}, f_n) < 2^{-(n+1)}\varepsilon$ for all $n \geq 1$. Let $f = \lim_n f_n$. This f satisfies (3.1) with $\alpha(\varepsilon) = \beta(\varepsilon/2)$.

PROPOSITION 3.4. *Suppose $\phi: X \rightarrow 2^Y$, with Y metric. Then:*

(a) *If ϕ satisfies (3.1) with $\alpha(\infty) = \infty$, then ϕ has the SEP.*

(b) *If X is collectionwise normal⁶, if Y is a Banach space, and if $\phi|_B$ satisfies (3.1) for every closed $B \subset X$ with $\alpha(\infty)$ independent of B , then ϕ has the SNEP.*

Proof. (a) Clear (see Footnote 5).

(b) Suppose $A \subset X$ is closed and g is a selection for $\phi|_A$. By a theorem of C. H. Dowker [1], g can be extended to a continuous $h: X \rightarrow Y$. Let $U = \{x \in X: d(h, \phi) < \alpha(\infty)\}$. Then U is open in X (since ϕ is l.s.c.) and $A \subset U$, so there is an open V in X such that $A \subset V \subset \bar{V} \subset U$. Then $d(h|_{\bar{V}}, \phi_a|_{\bar{V}}) < \alpha(\infty)$, so by assumption g extends to a selection f for $\phi|_{\bar{V}}$. That completes the proof.

4. A lemma about nerves of coverings. If \mathcal{U} is a covering of X , then we write $N(\mathcal{U})$ for the nerve of \mathcal{U} and $N^i(\mathcal{U})$ for the i -skeleton of $N(\mathcal{U})$. A function $u: N^i(\mathcal{U}) \rightarrow Y$ is called a *representation* if $u|_\sigma$ is continuous for every simplex σ of $N^i(\mathcal{U})$.

LEMMA 4.1. *Let Y be a Banach space and $\mathcal{S} \subset 2^Y$ uniformly equi-LCⁿ. Then to every $\varepsilon > 0$ corresponds a $\gamma(\varepsilon) > 0$ with the following property: If X is paracompact, $\phi: X \rightarrow \mathcal{S}$ l.s.c., $h: X \rightarrow Y$ continuous with $d(h, \phi) < \gamma(\varepsilon)$, and $\mu > 0$, then there exists a locally finite open cover \mathcal{U} of X and a representation $u: N^{n+1}(\mathcal{U}) \rightarrow Y$ such that, if σ is a simplex of $N^{n+1}(\mathcal{U})$ with vertices U_1, \dots, U_j , and if $x \in U_1 \cap \dots \cap U_j$, then*

⁵ That one can take $\alpha(\infty) = \infty$ simply means, in effect, that there exists a selection f for ϕ_a , with nothing of consequence being assumed or concluded about h . This useful shorthand, which was introduced in [4], will be used throughout this paper.

⁶ This property lies between normality and paracompactness.

$$u(\sigma) \subset B_{r'}(\phi(x)) \cap B_\varepsilon(h(x)).$$

If every $S \in \mathcal{S}$ is C^n , then one can take $\gamma(\infty) = \infty$.

Proof. This result is, in effect, a slight generalization of [4, Lemma 6.1]. In that lemma, it was assumed—in addition to our present hypotheses—that $\dim X \leq n + 1$, and it was concluded—in addition to our present conclusions—that the cover \mathcal{U} is of order $\leq n + 2$ (so that $N(\mathcal{U}) = N^{n+1}(\mathcal{U})$). The proof of [4, Lemma 6.1] remains valid for our present result.

5. **Proof of Theorem 1.2.** By Propositions 3.3 and 3.4, it will suffice to show that ϕ satisfies condition (3.2),⁷ and that one can take $\beta(\infty) = \infty$ in case $\phi(x)$ is C^n for every $x \in Z$. In fact, we will show that ϕ satisfies (3.2) with $\beta(\varepsilon) = \gamma(\varepsilon)$, where $\gamma(\varepsilon)$ is as in Lemma 4.1.

Let A, g and h be as in (3.2), with $\beta(\varepsilon) = \gamma(\varepsilon)$. Our hypotheses imply that $\{\phi_g(x) : x \in X\}$ is uniformly equi- LC^n , so we can pick an open cover \mathcal{U} of X and a representation $u : N^{n+1}(\mathcal{U}) \rightarrow Y$ as in Lemma 4.1 (applied to ϕ_g). Now recall that, by Dugundji's extension theorem, every convex subset of a Banach space has the extension property with respect to metric spaces. By inductively climbing up the i -skeletons of $N(\mathcal{U})$ with $i > n + 1$, we can therefore extend u to a representation $v : N(\mathcal{U}) \rightarrow Y$ such that $v(\sigma) \subset \text{conv } v(\dot{\sigma})$ for every simplex σ of $N(\mathcal{U})$ with $\dim \sigma > n + 1$. It follows that $v(\sigma) \subset \text{conv } u(\sigma \cap N^{n+1}(\mathcal{U}))$ for every simplex σ of $N(\mathcal{U})$.

From now on, the proof will closely follow the proof of Theorem 1.1 given in [7].

Let $\{V_U : U \in \mathcal{U}\}$ be an open cover of X such that $\bar{V}_U \subset U$ for all $U \in \mathcal{U}$. For each $x \in X$, let $\sigma(x)$ be the simplex of $N(\mathcal{U})$ spanned by those $U \in \mathcal{U}$ for which $x \in \bar{V}_U$. Let $S = X - Z$. For each $s \in S$, let $H_s = \bigcup \{\bar{V}_U : s \in \bar{V}_U\}$, and let

$$G_s = \{x \in X : v(\sigma_s) \subset [B_{r'}(\phi_g(x)) \cap B_\varepsilon(h(x))]\} - H_s.$$

Then G_s is open in X by Lemmas 2.1 and 2.2, while $s \in G_s$ by the properties of \mathcal{U} and u stated in Lemma 4.1, our construction of v , and the fact that $B_{r'}(\phi_g(s))$ and $B_\varepsilon(h(s))$ are convex. Also, by the last part of the definition of G_s , $\sigma_x \subset \sigma_s$ for every $x \in G_s$.

Let $G = \bigcup_{s \in S} G_s$, and let $E = X - G$. Then $E \subset Z$ and E is closed in X , so $\dim E \leq n + 1$. Now $\{V_U \cap E : U \in \mathcal{U}\}$ is a locally finite, relatively open cover of E , so it has a relatively open refinement $\{O_U : U \in \mathcal{U}\}$ of (indexed) order $\leq n + 2$ with $O_U \subset V_U$ for all

⁷ It is worth recording that this will also establish following result: If ϕ is as in Theorem 1.2, then ϕ has property (3.1).

$U \in \mathcal{U}$. Let $W_U = V_U \cap (O_U \cup G)$. Then $\{W_U: U \in \mathcal{U}\}$ is an open cover of X , $W_U \subset V_U \subset U$ for all $U \in \mathcal{U}$, and every $x \in E$ is in W_U for at most $n + 2$ elements $U \in \mathcal{U}$. Let $\xi: X \rightarrow N(\mathcal{U})$ be a canonical map obtained by using a partition of unity on X subordinated to $\{W_U: U \in \mathcal{U}\}$, and define $f: X \rightarrow Y$ by $f = v \circ \xi$. Let us show that $d(f, \phi_g) < \mu$ and $d(f, h) < \varepsilon$, as required by (3.2).

If $x \in E$, then $\xi(x) \in N^{n+1}(\mathcal{U})$, so

$$f(x) = v(\xi(x)) = u(\xi(x)) \in u(\sigma_x) \subset B_\mu(\phi_g(x)) \cap B_\varepsilon(h(x)),$$

where the last inclusion holds by our hypotheses on u . Now suppose $x \in G$. Then $x \in G_s$ for some $s \in S$. Hence $\sigma_x \subset \sigma_s$, so

$$f(x) = v(\xi(x)) \in v(\sigma_x) \subset v(\sigma_s) \subset B_\mu(\phi_g(x)) \cap B_\varepsilon(h(x)),$$

where the last inclusion holds by the definition of G_s . That completes the proof.

6. Proof of Theorem 1.3. We begin by recalling a definition from [5]. A metric space Y is a *uniform ANR* if to every $\varepsilon > 0$ corresponds a $\delta(\varepsilon) > 0$ with the following property: If Y is embedded isometrically as a closed subset in a metric space (E, d) , then there exists a retraction $r: B_{\delta(\varepsilon)}(Y) \rightarrow Y$ such that $d(z, r(z)) < \varepsilon$ whenever $z \in E$ and $d(z, Y) < \delta(\varepsilon)$. If one can take $\delta(\infty) = \infty$ (so that the domain of r is always E), then Y is called a *uniform AR*.

Before proving Theorem 1.3, we need the following lemma.

LEMMA 6.1. *Let (Y, d_0) be a metric ANR (resp. AR), and let $\mathcal{S} \subset 2^Y$ be equi- LC^n in Y . Then there exists a compatible metric $d \geq d_0$ on Y such that:*

- (a) (Y, d) is a uniform ANR (resp. uniform AR).
- (b) \mathcal{S} is uniformly equi- LC^n with respect to d .

Proof. The existence of a metric d satisfying (a) was proved in [5, Theorem 1.1]⁸, and the existence of a metric d satisfying (b) was proved in [4, Proposition 2.1]. Both results were proved with the aid of [2, Theorem 1], and that theorem can similarly be applied to obtain a metric d satisfying (a) and (b) simultaneously. We omit the details.

Having established Lemma 6.1, we now proceed to the proof of Theorem 1.3. We begin by remetrizing Y with a metric d as in Lemma 6.1, with $\mathcal{S} = \{\phi(x): x \in Z\}$; since the original metric d_0 on Y complete, so is the metric $d \geq d_0$. Next, we embed (Y, d) isometrically in a Banach space (E, d) . Let $\delta(\varepsilon)$ and $r: B_{\delta(\varepsilon)}(Y) \rightarrow Y$ be as in

⁸ For AR's, it had previously been proved by H. Toruńczyk in [8, Proposition 2.2].

the above definition of a uniform ANR. We shall regard ϕ as a map from X to $\mathcal{F}(E)$.

By Propositions 3.3 and 3.4, it will suffice to show that ϕ satisfies condition (3.2),⁹ and that one can take $\beta(\infty) = \infty$ in case Y is an AR and $\phi(x)$ is C^n for every $x \in Z$. In fact, we will show that ϕ satisfies (3.2) with $\beta(\varepsilon) = \gamma(\frac{1}{2}\delta(\varepsilon/2))$, where γ is as in Lemma 4.1 and δ is as in the above definition of a uniform ANR.

We proceed as in the proof of Theorem 1.2, with a few modifications. Let A , g and h be as in (3.2), with $\beta(\varepsilon) = \gamma(\frac{1}{2}\delta(\varepsilon/2))$. Our hypotheses imply that $\{\phi_g(x): x \in X\}$ is uniformly equi- LC^n , so we can choose an open cover \mathcal{U} of X and a representation $u: N^{n+1}(\mathcal{U}) \rightarrow E$ as in Lemma 4.1 (applied to ϕ_g), but with ε replaced by $\varepsilon' = \frac{1}{2}\delta(\varepsilon/2)$ and with μ replaced by $\mu' = \delta(\mu/2)$. In terms of this ε' , the above definition of $\beta(\varepsilon)$ becomes $\beta(\varepsilon) = \gamma(\varepsilon')$, so that $d(h, \phi_g) < \gamma(\varepsilon')$. Moreover, our choice of ε' and μ' yields the following two assertions, whose easy verification is left to the reader. (Recall that $r: B_{\delta(\infty)}(Y) \rightarrow Y$ is our retraction.)

(1) If $x \in X$, $e \in E$, and $e \in B_{\varepsilon'}(h(x))$, then $e \in B_{\delta(\varepsilon/2)}(Y)$ and $r(e) \in B_{\varepsilon}(h(x))$.

(2) If $x \in A$, $e \in E$ and $e \in B_{\mu'}(g(x))$, then $r(e) \in B_{\mu}(g(x))$.

Now extend the representation $u: N^{n+1}(\mathcal{U}) \rightarrow E$ to a representation $v: N(\mathcal{U}) \rightarrow E$ precisely as in the proof of Theorem 1.2. The properties of \mathcal{U} , u and v imply that, if σ is a simplex of $N(\mathcal{U})$ and if x is in the intersection of its vertices, then $v(\sigma) \subset B_{\varepsilon'}(h(x))$, so $v(\sigma) \subset B_{\delta(\infty)}(Y)$ by (1). It follows that $v(N(\mathcal{U}))$ is contained in the domain of r , and we can define $w = r \circ v$. Our proof now continues just like the proof of Theorem 1.2, except that v is replaced by w in the definitions of G_s and f . Everything goes through as before, except that it requires a bit more care to check that $s \in G_s$ and that $f(x) \in B_{\varepsilon}(h(x)) \cap B_{\mu}(\phi_g(x))$ when $x \in E$.

(a) $s \in G_s$: We must check that $w(\sigma_s) \subset B_{\varepsilon}(h(s))$ and that $w(\sigma_s) \subset B_{\mu}(\phi_g(s))$. For the first inclusion, observe that $v(\sigma_s) \subset B_{\varepsilon'}(h(s))$ by the previous paragraph, so $w(\sigma_s) \subset B_{\varepsilon}(h(s))$ by (1). For the second inclusion, we distinguish two cases: If $s \notin A$, then $\phi_g(s) = Y$, so our inclusion is clear because $w(\sigma_s) \subset Y$. If $s \in A$, then $\phi_g(s) = \{g(s)\}$; since $u(\sigma_s) \subset B_{\mu'}(\phi_g(s))$ by Lemma 4.1, we have $u(\sigma_s) \subset B_{\mu'}(g(s))$, hence $v(\sigma_s) \subset B_{\mu'}(g(s))$ by the construction of v , so (2) implies that $w(\sigma_s) \subset B_{\mu}(g(s)) = B_{\mu}(\phi_g(s))$.

(b) $f(x) \in B_{\varepsilon}(h(x)) \cap B_{\mu}(\phi_g(x))$ when $x \in E$: Just as in the proof of Theorem 1.2, we have $v(\xi(x)) \in B_{\varepsilon'}(h(x)) \cap B_{\mu'}(\phi_g(x))$ whenever $x \in E$; since $f(x) = r(v(\xi(x)))$, our assertion follows from (1) and (2).

That completes the proof.

⁹ It is worth recording that this will also establish the following result: If $\phi: X \rightarrow \mathcal{F}(Y)$ is as in Theorem 1.3, then Y can be re metrized so that ϕ has property (3.1).

REMARK. Since the above proof of Theorem 1.3 is fairly complicated, it may be worthwhile to outline an alternative proof for the important special case where $n = -1$: First of all, we now only need part (a) of Lemma 6.1. Next, let us show that (3.2) is satisfied with $\beta(\varepsilon) = \frac{1}{2}\delta(\varepsilon/2)$, where δ is as in the definition of a uniform ANR: With h and ϕ_g as in (3.2), define

$$\begin{aligned}\theta(x) &= \phi_g(x) \cap B_{\beta(\varepsilon)}(h(x)), \\ \psi(x) &= [\text{conv}(\theta(x)) \cap r^{-1}(B_{\mu/2}(\phi(x)))]^-.\end{aligned}$$

Standard facts about l.s.c. maps imply that ψ is l.s.c. Now $\psi(x) = [\text{conv}(\theta(x))]^-$ for all $x \in Z$, so ψ has a selection f' by Theorem 1.1.¹⁰ It is easy to check that $f'(x)$ is in $B_{\delta(\varepsilon/2)}(Y)$ —and thus in the domain of r —for all $x \in X$, so we can define $f = r \circ f'$. It is not hard to check that this f is an extension of g which satisfies all the requirements of (3.2).

7. Proof of Theorem 1.4. As observed in the introduction, this theorem reduces to [4, Theorem 1.2] when $Z = \emptyset$. The proof of Theorem 1.4 is very similar to that of the older result given in [4], so it will suffice to indicate the necessary modifications.

(1) Analogously to [4, Proposition 2.1], one proves that there exists a compatible complete metric d on Y which simultaneously makes $\{\phi(x): x \in X - Z\}$ uniformly LC^n and $\{\phi(x): x \in Z\}$ uniformly equi- LC^m . (Observe that $\{\phi(x): x \in X\}$ is therefore also uniformly equi- LC^m .)

(2) After the modification in (1) above, we can follow the proof of [4, Theorem 1.2] essentially without change until we get to the proof of [4, Lemma 7.3]. In that proof, the hypothesis that $\{\phi(x): x \in X\}$ is uniformly equi- LC^n was needed to construct the maps $u_{\sigma,x}$ (for $\dim \sigma = i + 1$) in the middle of p. 572 of [4]. Since $\{\phi(x): x \in X\}$ is uniformly equi- LC^m in our present situation, we can construct $u_{\sigma,x}$ as before when $i \leq m$. For $i > m$, however, $u_{\sigma,x}$ can be defined as before only if $x \in X - Z$, but not, in general, when $x \in Z$. To overcome this obstacle, it suffices to modify the proof of [4, Lemma 7.3] when $i > m$ by first of all shrinking the given open cover \mathcal{U} of X to an open cover $\mathcal{U}' = \{U': U \in \mathcal{U}\}$ with the following two properties:

(a) $U' \subset U$ for all $U \in \mathcal{U}$.

(b) For the cover \mathcal{U}' , the construction of suitable maps $u'_{\sigma,x}$ is possible for all $x \in X$. More precisely: If σ is a simplex of

¹⁰ This is where we need the assumption that $n = -1$. For $n > -1$, there is no reason to suppose that ψ satisfies the hypotheses of Theorem 1.2. That explains why, in our proof of the general case of Theorem 1.3, we were unable to apply Theorem 1.2 and therefore had to imitate its proof instead.

$N^{i+1}(\mathcal{U})$ with $\dim \sigma = i + 1$ and with vertices U_1, \dots, U_{i+2} , and if $x \in U_1' \cap \dots \cap U_{i+2}'$ then $u|_{\sigma}$ can be extended to a continuous $u'_{x,\sigma}: \sigma \rightarrow Y$ such that $u'_{x,\sigma}(\sigma) \subset B_{\mu}(\phi(x))$ and $\text{diam } u'_{x,\sigma}(\sigma) < \varepsilon/2$.

We construct \mathcal{U}' as follows. Let $S = X - Z$. If σ is a simplex of $N^{i+1}(\mathcal{U})$ with $\dim \sigma = i + 1$, and if $x \in S$ is in the intersection of the vertices of σ , then (since $x \notin Z$) we can define $u_{x,\sigma}$, $W_{x,\sigma}$ and W_x precisely as in the middle of p. 572 of [4]. Now pick an open cover $\{U^*: U \in \mathcal{U}\}$ of X such that $\bar{U}^* \subset U$ for all $U \in \mathcal{U}$, and for $s \in S$ let

$$G_s = W_s - \bigcup \{ \bar{U}^*: U \in \mathcal{U}, s \notin \bar{U}^* \}.$$

Clearly G_s is open in X and $s \in G_s$. Let $G = \bigcup_{s \in S} G_s$, and let $E = X - G$. Then $E \subset Z$ and E is closed in X , so $\dim E \leq m + 1$. Hence the relatively open cover $\{U^* \cap E: U \in \mathcal{U}\}$ of E has a relatively open refinement $\{O_U: U \in \mathcal{U}\}$ of (indexed) order $\leq m + 2$ with $O_U \subset U^*$ for all $U \in \mathcal{U}$. Now let $U' = U^* \cap (O_U \cup G)$ for all $U \in \mathcal{U}$, and let $\mathcal{U}' = \{U': U \in \mathcal{U}\}$. Clearly \mathcal{U}' is an open cover of X , so let us verify that it satisfies conditions (a) and (b).

That (a) is satisfied is clear, so it remains to check (b). Now if $x \in E$, then (b) is trivially satisfied, since then $x \in U'$ for at most $m + 2 \leq i + 1$ sets $U \in \mathcal{U}$. So suppose $x \in G$. Then we can choose $s \in S$ such that $x \in G_s$. Now if σ and U_1, \dots, U_{i+2} are as in (b), then $x \in U_1^* \cap \dots \cap U_{i+2}^*$, so the definition of G_s implies that

$$s \in (\bar{U}_1^* \cap \dots \cap \bar{U}_{i+2}^*) \subset (U_1 \cap \dots \cap U_{i+2}).$$

Thus s is in the intersection of the vertices of σ , so $u_{s,\sigma}$ and $W_{s,\sigma}$ are defined and $W_s \subset W_{s,\sigma}$. Hence $x \in G \subset W_s \subset W_{s,\sigma}$. We now let $u'_{x,\sigma} = u_{s,\sigma}$. This choice of $u'_{x,\sigma}$ satisfies (b) because $x \in W_{s,\sigma}$. That completes the proof.

REFERENCES

1. C. H. Dowker, *On a theorem of Hanner*, Ark. Mat., **2** (1952), 307-313.
2. J. Dugundji and E. Michael, *On local and uniformly local topological properties*, Proc. Amer. Math. Soc., **7** (1956), 304-307.
3. E. Michael, *Continuous selections I*, Ann. of Math., **63** (1956), 361-382.
4. ———, *Continuous selections II*, Ann. of Math., **64** (1956), 562-580.
5. ———, *Uniform AR's and ANR's*, Compositio Math., **39** (1979), 129-149.
6. ———, *Continuous selections and countable sets*, to appear in Fund. Math.
7. E. Michael and C. Pixley, *A unified theorem on continuous selections*, Pacific J. Math., **87** (1980), 187-188.
8. H. Toruńczyk, *Absolute retracts as factors of normed linear spaces*, Fund. Math., **86** (1974), 53-76.

Received May 21, 1979. Supported in part by a grant from the National Science Foundation at the University of Washington and in part by a grant from the Alexander von Humboldt Foundation at the University of Stuttgart.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE AND ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Spiros Argyros, <i>A decomposition of complete Boolean algebras</i>	1
Gerald A. Beer, <i>The approximation of upper semicontinuous multifunctions by step multifunctions</i>	11
Ehrhard Behrends and Richard Evans, <i>Multiplicity theory for Boolean algebras of L^p-projections</i>	21
Man-Duen Choi, <i>The full C^*-algebra of the free group on two generators</i>	41
Jen-Chung Chuan, <i>Axioms for closed left ideals in a C^*-algebra</i>	49
Jo-Ann Deborah Cohen, <i>The strong approximation theorem and locally bounded topologies on $F(X)$</i>	59
Eugene Harrison Gover and Mark Bernard Ramras, <i>Increasing sequences of Betti numbers</i>	65
Morton Edward Harris, <i>Finite groups having an involution centralizer with a 2-component of type $PSL(3, 3)$</i>	69
Valéria Botelho de Magalhães Iório, <i>Hopf C^*-algebras and locally compact groups</i>	75
Roy Andrew Johnson, <i>Nearly Borel sets and product measures</i>	97
Lowell Edwin Jones, <i>Construction of Z_p-actions on manifolds</i>	111
Manuel Lerman and Robert Irving Soare, <i>d-simple sets, small sets, and degree classes</i>	135
Philip W. McCartney, <i>Neighborly bushes and the Radon-Nikodým property for Banach spaces</i>	157
Robert Colman McOwen, <i>Fredholm theory of partial differential equations on complete Riemannian manifolds</i>	169
Ernest A. Michael and Carl Preston Pixley, <i>A unified theorem on continuous selections</i>	187
Ernest A. Michael, <i>Continuous selections and finite-dimensional sets</i>	189
Vassili Nestoridis, <i>Inner functions: noninvariant connected components</i>	199
Bun Wong, <i>A maximum principle on Clifford torus and nonexistence of proper holomorphic map from the ball to polydisc</i>	211
Steve Wright, <i>Similarity orbits of approximately finite C^*-algebras</i>	223
Kenjiro Yanagi, <i>On some fixed point theorems for multivalued mappings</i>	233
Wieslaw Zelazko, <i>A characterization of LC-nonremovable ideals in commutative Banach algebras</i>	241