

Pacific Journal of Mathematics

HEIGHT ESTIMATES FOR CAPILLARY SURFACES

DAVID SIEGEL

HEIGHT ESTIMATES FOR CAPILLARY SURFACES

DAVID SIEGEL

In this paper estimates are obtained for any scalar function $u(x)$ that satisfies the equation

$$(1a) \quad \operatorname{div}(Tu) = \kappa u \quad \text{in } \Omega$$

and the boundary condition

$$(1b) \quad Tu \cdot \nu = \cos \gamma \quad \text{on } \Sigma = \partial\Omega .$$

Here κ is a positive constant, Ω is an open domain in n -dimensional Euclidean space, ν is the exterior unit normal on Σ , and Tu is the vector operator

$$Tu = \frac{1}{\sqrt{1 + |\nabla u|^2}} \nabla u .$$

For $n = 2$, $u(x)$ can be interpreted physically as the height of a capillary surface above the undisturbed fluid level when a vertical cylindrical tube with section Ω is dipped into a large reservoir. The "capillarity constant" κ and the "contact angle" γ are determined physically; $\kappa = (\rho - \rho_0)\sigma^{-1}g$, where ρ is the density of the fluid, ρ_0 is the density of the gas, g is the acceleration due to gravity, and σ is the surface tension. If the tube is homogeneous γ is constant [4].

The operator $Nu = \operatorname{div}(Tu)$ is n times the mean curvature of the surface $x_{n+1} = u(x)$. Geometrically stated, a capillary surface has mean curvature proportional to its height above a horizontal reference plane and it meets a vertical cylinder in a prescribed angle.

We shall distinguish three types of domains: "interior", "exterior", and "general exterior" corresponding to Ω bounded, the complement of Ω bounded, and Σ unbounded, respectively. Existence, uniqueness, and regularity of solutions to problem (1) have been established under fairly general conditions on Ω and γ . However, for a general exterior domain uniqueness has not been established and for an exterior domain uniqueness has been established only under the condition $u = o(1)$ as $|x| \rightarrow \infty$ [see 6, 7, 8, 9].

For simplicity we shall only consider solutions to problem (1) in domains with piecewise smooth boundaries, with boundary condition (1b) holding on the smooth part Σ^* of Σ . In §2, we extend the existence theory for smooth bounded domains to the case of piecewise smooth bounded or unbounded domains.

We now state our main results. In what follows $\kappa = 1$, and γ is constant, $0 \leq \gamma \leq \pi/2$.

The simplest examples of interior, exterior, and general exterior domains are $B_R(0) = \{x: |x| < R\}$, $B_R^c(0) = \{x: |x| > R\}$ and $H = \{x: x_1 > 0\}$. The solutions to problem (1) in these domains are $v(r, R, \gamma)$, $w(r, R, \gamma)$, and $z(x_1, \gamma)$, with $r = |x|$. An explicit formula is known for the "one-dimensional" solution $z(x_1, \gamma)$. We have

$$(2) \quad z(x_1, \gamma) \sim C_1(\gamma)e^{-x_1} \quad \text{as } x_1 \longrightarrow \infty$$

where $C_1(\gamma)$ is an explicitly known constant.

Chapter II is devoted to the study of v and w . General estimates are given for v and w that improve upon those given by Finn [5]. For small R , v is shown to be close to a spherical cap. Laplace's asymptotic formula for the center height is proved correct. For large R , near the boundary v and w are shown to be close to a one-dimensional solution, i.e.,

$$\lim_{R \rightarrow \infty} v(R - l, R, \gamma) = \lim_{R \rightarrow \infty} w(R + l, R, \gamma) = z(l, \gamma).$$

Away from the boundary, estimates are also given, in particular

$$(3) \quad v(0, R, \gamma) \sim C(\gamma) \frac{R^{(n-1)/2}}{\exp R} \quad \text{as } R \longrightarrow \infty$$

$$(4) \quad w(r, R, \gamma) \sim C(R, \gamma) \frac{\exp(-r)}{r^{(n-1)/2}} \quad \text{as } r \longrightarrow \infty$$

where $C(\gamma)$ is an explicitly given constant and $C(R, \gamma)$ is determined asymptotically as $R \rightarrow \infty$. Monotonicity and continuity properties of $C(R, \gamma)$ are given. Estimates on the derivatives of v and w are also given.

In Chapter III, solutions to problem (1) are estimated in terms of v , w , and z . This is done by use of an appropriate comparison principle due to Concus and Finn [3].

General estimates are given that apply to any solution of equation (1a): $|u(x)| \leq v(0, d, 0)$, d is the distance from x to Σ ; $|u(x)| \leq w(r, R, 0)$ if $B_R^c(0) \subset \Omega$; and $|u(x)| \leq z(x_1, 0)$ if $H \subset \Omega$. Combining these general estimates with results (2)-(3)-(4), we see that solutions to equation (1a) decay exponentially away from the boundary of the domain. The derivatives of a solution decay at the same rate at which the solution decays.

The third general estimate improves on an estimate given by Gerhardt [8]. The first general estimate implies that for an exterior domain, problem (1) has a unique solution.

We now consider estimating solutions to problem (1). To improve upon the above general estimates or to study solutions in general exterior domains one needs to know when $B_R(y) \subset \Omega$, $r = |x - y|$,

implies that $u(x) \leq v(r, R, \gamma)$. An example shows that this is not always true. We prove that if $\Omega \cap B_{\bar{r}}(y)$ is convex, where $B_{\bar{r}}(y)$ is the maximal domain of existence of v , then the assertion is true. Alternatively, if Σ is smooth and bounded and $R \leq \mathcal{R}_1$, where \mathcal{R}_1 is the “interior rolling number” of Ω , then the assertion is also true. Analogously, if $B_{\bar{r}}(y) \subset \Omega$, $r = |x - y|$, if Σ is smooth bounded and strictly convex, and if $R \geq \mathcal{R}_2$, where \mathcal{R}_2 is the “exterior rolling number”, then $u(x) \leq w(r, R, \gamma)$. Again, an example shows that this inequality is not always true when $B_{\bar{r}}(y) \subset \Omega$.

Estimates for solutions to problem (1) in certain general exterior domains are given. We show that $z(x, \gamma)$ is the unique solution to problem (1) in H . Consider the two-dimensional domain $\mathcal{H}_\alpha = \{(x_1, x_2): x_1 > |x| \cos \alpha\}$, with $0 < \alpha < \pi$. For $0 < \gamma < \pi/2$ there exists a unique positive solution to problem (1) in \mathcal{H}_α . Furthermore, for $\alpha < \pi/2$: on Σ^* , $u(x) \geq z(0, \gamma)$ and $\lim_{|x| \rightarrow \infty} u(x) = z(0, \gamma)$. For $\alpha > \pi/2$: on Σ^* , $u(x) \leq z(0, \gamma)$ and $\lim_{|x| \rightarrow \infty} u(x) = z(0, \gamma)$. For $\alpha + \gamma \geq \pi/2$, $\alpha < \pi/2$:

$$\liminf_{x \rightarrow 0} u(x) \geq \sqrt{2} [1 - (1 - k^2)^{1/2}]^{1/2}$$

and

$$\limsup_{x \rightarrow 0} u(x) \leq 2\sqrt{2} [1 - (1 - k^2)^{1/2}]^{1/2}$$

where $k = \cos \gamma / \sin \alpha$. These inequalities show there is a “rise” at the corner. Some higher dimensional generalizations are given.

Chapter I contains preliminaries that are needed in the other chapters.

Chapter I Preliminaries

1. The comparison principle. Our basic tool will be the comparison principle (CP):

THEOREM 1. *Let $\Sigma = \Sigma^0 + \Sigma^\alpha + \Sigma^\beta$ be a decomposition of Σ , such that Σ^β is C^1 and Σ^0 can be covered from within Ω by a sequence of smooth surfaces $\{A\}$, each of which meets Σ in a set of zero ($n - 1$) dimensional measure, and such that $A \rightarrow \Sigma^0$ and the area of A tends to zero. Let $u, v \in C^2(\Omega)$ and suppose*

- (i) $Nu \geq \kappa u$ and $Nv \leq \kappa v$ on Ω .
- (ii) for any approach to Σ^α from within Ω $\limsup [u - v] \leq 0$.
- (iii) on Σ^β , $(Tu - Tv) \cdot \nu \leq 0$ almost everywhere as a limit from points of Ω .
- (iv) if Ω is unbounded, $u - v \leq o(1)$ as $|x| \rightarrow \infty$.

Then $u \leq v$ on Ω ; if equality holds at any point then $u \equiv v$ on Ω .

Proof (see [3]). We have added condition (iv), but precisely the same proof holds.

Note. In condition (iii), ν has been extended continuously into a neighborhood of Σ^β .

The following corollary was proved in [3] in a different way.

COROLLARY. *If $u(x)$ satisfies (1a) in Ω and $B_R(y) \subset \Omega$ then $|u(x)| < 2/\kappa R + R$ in $B_R(y)$.*

Proof. Let $v(x) = n/\kappa R + R - \sqrt{R^2 - |x - y|^2}$. Then $Nv = n/R = \kappa \min v(x) \leq \kappa v$ and $\lim Tv \cdot \nu = 1$. Thus $\limsup (Tu - Tv) \cdot \nu \leq 0$. By CP $u(x) < v(x) < \sup v(x) = n/\kappa R + R$ on $B_R(y)$. Replace v by $-v$ to get the other inequality.

When Ω is contained in a symmetric domain we have the following comparison principle (CPS):

THEOREM 2. *Let Ω be bounded and Σ piecewise smooth. Suppose $u(x)$ is a solution to problem (1) with boundary data $\gamma(\sigma)$, $0 \leq \gamma(\sigma) \leq \gamma_0$, $0 \leq \gamma_0 \leq \pi/2$, on Σ^* .*

(i) *If $\Omega \subset B_R(0)$ then $v < u$ or $v \equiv u$ on Ω .*

(ii) *If $\Omega \subset B_R^c(0)$ then $w < u$ on Ω .*

(iii) *If $\Omega \subset H$ then $z(x_1) < u(x)$ on Ω .*

Here $v = v(r, R, \gamma_0)$, $w = w(r, R, \gamma_0)$, and $z(x_1) = z(x_1, \gamma_0)$.

Proof. The proofs for parts (i) and (ii) are in [5]. We present the proof for (iii) which is very similar. We introduce the angle $\psi_1(x_1)$ between the curve and the positive x_1 direction

$$\sin \psi_1(x_1) = \frac{z_{x_1}}{\sqrt{1 + z_{x_1}^2}}.$$

It is known that $\sin \psi_1(x_1)$ is negative and increasing in x_1 (see §5). On Σ^* , $Tu \cdot \nu = \cos \gamma \geq \cos \gamma_0 = -\sin \psi_1(0) \geq -\sin \psi_1(x_1) \geq Tz \cdot \nu$. The last inequality is true because $|Tz| = -\sin \psi_1(x_1)$. Apply CP to obtain $z(x_1) < u(x)$ on Ω .

We shall need the following consequence of the boundary point form of Hopf's maximum principle [11].

LEMMA 1. *Let $u \in C^1(\bar{B}_R) \cap C^2(B_R)$ and $v \in C^2(\bar{B}_R)$. Suppose u and*

v satisfy condition (i) of Theorem 1, $u < v$ on B_R and $u(x_0) = v(x_0)$ at $x_0 \in \partial B_R$. Then $(\partial u / \partial \nu)(x_0) \neq (\partial v / \partial \nu)(x_0)$.

Proof. Let $w = u - v$, then w satisfies

$$(5) \quad Lw = \sum_{i,j} a_{ij} w_{x_i x_j} + \sum_k b_k w_{x_k} \geq \kappa w$$

where

$$a_{ij} = a_{ij}(\nabla u), \quad Nu = \sum a_{ij}(\nabla u) u_{x_i x_j}$$

$$b_k = \sum_{i,j} v_{x_i x_j} \int_0^1 \frac{\partial a_{ij}}{\partial p_k} [\nabla v + t(\nabla u - \nabla v)] dt .$$

Under the hypotheses L is uniformly elliptic with bounded coefficients on B_R . Since $w < 0$ on B_R and $w(x_0) = 0$, Hopf's theorem gives $(\partial w / \partial \nu)(x_0) \neq 0$.

REMARK. The smoothness properties of u and v can be interchanged in Lemma 1.

2. Existence theorems. Gerhardt [9] proves the following result:

THEOREM 3. *If Ω is bounded, $\Sigma \in C^{2,\lambda}$, $0 < \gamma(\sigma) < \pi$, $\gamma(\sigma) \in C^{1,\lambda}$, then there exists a unique solution $u(x)$ to problem (1) with $u(x) \in C^{2,\alpha}(\bar{\Omega})$.*

Simon and Spruck [16] prove a similar theorem under the condition $\Sigma \in C^4$. They give a local estimate of the gradient up to the boundary. With this estimate we prove:

THEOREM 4. *Let Ω be a domain with a piecewise smooth boundary Σ . Let $\Sigma = \Sigma^0 + \Sigma^*$, where Σ^* is open in Σ and $\Sigma^* \in C^4$. Suppose that on Σ^* , $\gamma(\sigma) \in C^{1,\alpha}$ and $0 < \gamma(\sigma) < \pi$. Then there exists a solution $u(x)$ to (1), with boundary condition (1b) holding on Σ^* .*

Proof.

Step 1. We construct approximating domains Ω_i with C^4 boundaries Σ_i . For Ω bounded we require:

- (a) if $x \in \Sigma$ and $\text{dist}(x, \Sigma^0) > \varepsilon_i$ then $x \in \Sigma_i$
- (b) $\text{dist}(\Sigma, \Sigma_i) < \varepsilon_i$

with $\lim_{i \rightarrow \infty} \varepsilon_i = 0$. For Ω unbounded we require $\Omega_{R_i} = \Omega \cap B_{R_i}(0)$ to be piecewise C^4 , with $\lim_{i \rightarrow \infty} R_i = \infty$, and that conditions (a) and (b) hold with Σ replaced by $\partial \Omega_{R_i}$.

Extend $\gamma(\sigma)$ to all of Σ_i so that $0 < \gamma(\sigma) < \pi$ and $\gamma(\sigma) \in C^{1,\alpha}$. Let u_i be the solution to problem (1) with this data, $u_i \in C^2(\bar{\Omega})$.

Step 2. We now obtain a local Hölder estimate of the gradient of u_i up to Σ^* , independent of i .

Take $x_0 \in \Sigma^*$, choose $\delta > 0$ so that $\bar{B}_{4\delta}(x_0) \cap \bar{\Omega} \subset \Omega \cup \Sigma^*$, choose N so that $i > N$ implies $\bar{B}_{4\delta} \cap \Sigma^* \subset \Sigma_i$.

We estimate $|u_i(x)| \leq M$ on $\bar{B}_{3\delta}(x_0) \cap \bar{\Omega}$. By the corollary to Theorem 1 we can choose $M = n(\kappa R)^{-1} + R$ if each point of $\bar{B}_{3\delta} \cap \bar{\Omega}$ is contained in a closed ball of radius R lying in $\bar{B}_{4\delta} \cap \bar{\Omega}$. One can choose $R = \min[\delta, (\bar{k})^{-1}]$ if $\bar{k} > 0$, $R = \delta$ otherwise, where $\bar{k} = \max k(\sigma)$ over $\bar{B}_{4\delta} \cap \Sigma^*$, $k(\sigma) =$ the maximum principal curvature of Σ at σ with respect to the interior normal.

By Theorem 3 of [16]: $|\nabla u_i|_{0, \bar{\Omega} \cap B_{2\delta}(x_0)} < \Gamma_1$.

By the argument on pp. 467-8 of [13]: $|u_i|_{1, \alpha, \bar{\Omega} \cap B_\delta(x_0)} < \Gamma_2$. Here Γ_1 and Γ_2 are independent of i .

Step 3. We now obtain interior estimates on the derivatives of u_i independent of i .

Take $x_0 \in \Omega$, choose $\delta > 0$ so that $\bar{B}_{4\delta}(x_0) \subset \Omega$, choose N so that $i > N$ implies $B_{4\delta} \subset \Omega_i$.

We estimate $|u_i| \leq n(4\kappa\delta)^{-1} + 4\delta$ on $\bar{B}_{4\delta}$.

The gradient can be estimated on $B_{3\delta}$ since

$$|\nabla u_i(y)| \leq C_1 \exp\{C_2 \sup [u_i - u_i(y)]/\delta\}$$

where C_1 and C_2 depend on n and $\delta \sup |u_i|$ (see [10]). This gives $|\nabla u_i|_{0, B_{3\delta}(x_0)} < \Gamma_3$.

The interior Hölder estimate for divergence structure equations [11, p. 265] yields $|\nabla u_i|_{\beta, B_{2\delta}(x_0)} < \Gamma_4$.

Finally, we apply the interior Schauder estimate, since we can treat the equation as a linear, uniformly elliptic equation with C^β coefficients. This gives $|u_i|_{2, \beta, B_\delta(x_0)} < \Gamma_5$. Here Γ_3 , Γ_4 , and Γ_5 are independent of i .

Step 4. Because of the boundary and interior estimates we can choose a subsequence of $\{u_i\}$ that converges in C^2 on every compact set $k \subset \Omega$, and in C^1 on every compact set $k \subset \Omega \cup \Sigma^*$. The limit function $u(x)$ will belong to $C^2(\Omega) \cap C^1(\Omega \cup \Sigma^*)$ and will be a solution to problem (1).

REMARK. If Ω is bounded the solution to problem (1) is unique. This is an immediate consequence of CP.

3. Two normalizations. From now on we shall take κ to be

one. This is no loss in generality because of the following lemma:

LEMMA 2. *Let $u(x)$ satisfy $Nu = u$ in Ω and $Tu \cdot \nu = \cos \gamma$ on Σ . Define $v(x) = (1/\sqrt{\kappa})u(\sqrt{\kappa}x)$. Let Ω' , Σ' , and ν' be the images of Ω , Σ , and ν under the transformation $x \rightarrow (1/\sqrt{\kappa})x$. Then $Nv = \kappa v$ in Ω' and $Tv \cdot \nu' = \cos \gamma$ on Σ' .*

Proof. From the expression

$$Nu = \frac{\Delta u(1 + |\nabla u|^2) - \sum u_{x_i} u_{x_j} u_{x_i x_j}}{(1 + |\nabla u|^2)^{3/2}}$$

we see that $Nv(x) = \sqrt{\kappa}Nu(\sqrt{\kappa}x) = \sqrt{\kappa}u(\sqrt{\kappa}x) = \kappa v(x)$. Also, $Tv(x/\sqrt{\kappa}) = Tu(x)$ in Ω and $\nu'(x/\sqrt{\kappa}) = \nu(x)$ on Σ . Thus, $Tv \cdot \nu' = \cos \gamma$ on Σ' .

If γ is constant we shall assume $0 \leq \gamma \leq \pi/2$. This is no loss in generality because if $u(x)$ satisfies $Nu = u$ in Ω and $Tu \cdot \nu = \cos \gamma$ on Σ , then $v(x) = -u(x)$ satisfies $Nv = v$ in Ω and $Tv \cdot \nu = \cos(\pi - \gamma)$ on Σ .

4. Solutions to the linearized equation. We examine the solutions to the linearized equation $\Delta u = u$ that depend only on $r = |x|$; u must satisfy

$$(6) \quad u_{rr} + \frac{n-1}{r} u_r = u.$$

The general solution is $u(r) = AI(r; n) + BK(r; n)$; A and B are arbitrary constants, $I(r; n) = r^{-m}I_m(r)$, $K(r; n) = r^{-m}K_m(r)$, $m = (n-2)/2$, I_m is a modified Bessel function of the first kind, and K_m is a MacDonald's function.

To be explicit (see [14]):

$$(7) \quad I_m(r) = \sum_{k=0}^{\infty} \frac{(r/2)^{m+2k}}{\Gamma(k+1)\Gamma(k+m+1)}$$

$$(8) \quad K_m(r) = \int_0^{\infty} e^{-r \cosh u} \cosh(mu) du.$$

The following recurrence relations hold:

$$(9) \quad (r^{-m}I_m)_r = r^{-m}I_{m+1}$$

$$(10) \quad (r^{-m}K_m)_r = -r^{-m}K_{m+1}.$$

From equations (7) through (10) we conclude:

$$(11) \quad \begin{aligned} I_{rr} > 0 \text{ and } I > 0 \text{ for } r \geq 0; \quad I_r > 0 \text{ for } r > 0; \\ I_r(0) = 0 \quad \text{and} \quad I(0) = [2^{(n-2)/2} \Gamma(n/2)]^{-1} \end{aligned}$$

$$(12) \quad \begin{aligned} K_{rr} > 0, \quad K_r < 0, \quad \text{and} \quad K > 0 \quad \text{for } r > 0; \\ \lim_{r \rightarrow \infty} K(r) = \lim_{r \rightarrow \infty} K_r(r) = 0. \end{aligned}$$

We shall employ I and K to estimate $v(r, R, \gamma)$ and $w(r, R, \gamma)$. In fact, AI and AK , for any positive constant A , satisfy a super solution condition:

LEMMA 3. For any constant A , $A > 0$, $N(AI) \leq AI$, and $N(AK) \leq AK$.

Proof. For a function $u = u(r)$

$$(13) \quad Nu = \frac{1}{r^{n-1}} \left(\frac{r^{n-1} u_r}{\sqrt{1 + u_r^2}} \right)_r.$$

We use the properties listed under (11) and (12):

$$\begin{aligned} N(AI) &= \frac{AI_{rr}}{[1 + (AI_r)^2]^{3/2}} + \frac{(n-1)}{r} \frac{AI_r}{[1 + (AI_r)^2]^{1/2}} \\ &\leq AI_{rr} + \frac{(n-1)}{r} AI_r = AI \\ N(AK) &= \frac{AK_{rr} + \frac{(n-1)}{r} (AK_r)[1 + (AK_r)^2]}{[1 + (AK_r)^2]^{3/2}} \\ &= \frac{AK + \frac{(n-1)}{r} (AK_r)^3}{[1 + (AK_r)^2]^{3/2}} < AK. \end{aligned}$$

The asymptotic behavior of I_m and K_m has been studied (see [14]). As a consequence of these estimates we obtain

$$(14) \quad I(r) = \frac{1}{\sqrt{2\pi}} \frac{e^r}{r^{(n-1)/2}} [1 + O(1/r)] \quad \text{as } r \longrightarrow \infty$$

$$(15) \quad K(r) = \sqrt{\frac{\pi}{2}} \frac{e^{-r}}{r^{(n-1)/2}} [1 + O(1/r)] \quad \text{as } r \longrightarrow \infty$$

$$(16) \quad I_r(r) = \frac{1}{\sqrt{2\pi}} \frac{e^r}{r^{(n-1)/2}} [1 + O(1/r)] \quad \text{as } r \longrightarrow \infty$$

$$(17) \quad K_r(r) = -\sqrt{\frac{\pi}{2}} \frac{e^{-r}}{r^{(n-1)/2}} [1 + O(1/r)] \quad \text{as } r \longrightarrow \infty.$$

We now give explicit estimates on I_r/I and K_r/K . For any solution u of equation (6), $v = u_r/u$ satisfies a nonlinear first order differential equation

$$(18) \quad v_r = 1 - \frac{(n-1)}{r} v - v^2.$$

Let

$$\begin{aligned} \mathcal{S} &= \left\{ (r, v): r > 0 \text{ and } 1 - \frac{(n-1)}{r} v - v^2 > 0 \right\} \\ &= \{ (r, v): r > 0 \text{ and } v_1(r) < v < v_2(r) \} \end{aligned}$$

where

$$v_1(r) = -\frac{(n-1)}{2r} - \left[1 + \left(\frac{n-1}{2r} \right)^2 \right]^{1/2}$$

and

$$v_2(r) = -\frac{(n-1)}{2r} + \left[1 + \left(\frac{n-1}{2r} \right)^2 \right]^{1/2}.$$

With this notation, we have:

LEMMA 4. For $r > 0$:

- (i) I_r/I and K_r/K are strictly increasing in r .
- (ii) $0 < I_r/I < v_2(r)$.
- (iii) $v_1(r) < K_r/K < -1$.

Proof. Part (i) is equivalent to saying that $(r, I_r/I)$ and $(r, K_r/K)$ are contained in \mathcal{S} , thus we need only prove parts (ii) and (iii).

We note that $v_1(r)$ and $v_2(r)$ are increasing functions of r .

Let $v = I_r/I$, then $v(0) = 0$ and $v(r) > 0$ for $r > 0$. Suppose $v(r_1) \geq v_2(r_1)$, then $v(r) \geq v_2(r_1) > 0$ for $r < r_1$, contradicting that $v(0) = 0$. This proves part (ii).

Let $w = K_r/K$, then $w < 0$. Since $K_r < 0$, we have $K_{rr} = K - ((n-1)/r)K_r > K$ and $K_r K_{rr} < K_r K$. Integrating from r to ∞ gives $K_r^2 > K^2$ and hence $K_r/K < -1$.

Suppose $w(r_1) \leq v_1(r_1)$, then for $r > r_1$, $w(r) < v_1(r_1) = -a$, with $a > 1$. Integrating from r_1 to r gives $K(r) < K(r_1) \exp a(r_1 - r)$ for $r > r_1$. This contradicts equation (15) as $r \rightarrow \infty$. The proof of part (iii) is complete.

REMARK. The upper bound on I_r/I and the lower bound on K_r/K are asymptotically exact to order $1/r$ since

$$\frac{I_r}{I} = 1 - \frac{(n-1)}{2r} + O\left(\frac{1}{r^2}\right) \quad \text{as } r \longrightarrow \infty$$

and

$$\frac{K_r}{K} = -1 - \frac{(n-1)}{2r} + O\left(\frac{1}{r^2}\right) \quad \text{as } r \longrightarrow \infty .$$

5. The one-dimensional solution. There is only one explicitly known solution to problem (1). We denote by $z(x; \gamma)$ the unique solution to

$$(19a) \quad \left(\frac{z_x}{\sqrt{1+z_x^2}}\right)_x = z \quad \text{for } 0 < x < \infty$$

$$(19b) \quad z_x(0+) = -\cot \gamma$$

$$(19c) \quad \lim_{x \rightarrow \infty} z(x) = 0 .$$

Here, z can be interpreted physically as the height of a capillary surface on one side of an infinite vertical plate.

An explicit solution with x as a function of z is given by

$$(20) \quad \begin{aligned} x &= -2 \sqrt{1 - \frac{z^2}{4}} - \ln \frac{z}{2} + \ln \left(1 + \sqrt{1 - \frac{z^2}{4}}\right) + D \\ D &= \sqrt{2(1 + \sin \gamma)} + \ln \sqrt{\frac{1 - \sin \gamma}{2}} - \ln \left(1 + \sqrt{\frac{1 + \sin \gamma}{2}}\right) . \end{aligned}$$

THEOREM 5. For $0 \leq \gamma < \pi/2$:

(i) There is a unique solution z to problem (19) given by (20), $z(x) \in C^2(0, \infty) \cap C[0, \infty)$, with $z > 0$ and $z_x < 0$.

(ii) $z = C_1(\gamma)e^{-x}[1 + O(e^{-2x})]$ as $x \rightarrow \infty$ with $C_1(\gamma) = 4 \exp(D - 2)$.

(iii) $\sin \psi_1(x) = -C_1(\gamma)e^{-x}[1 + O(e^{-2x})]$ as $x \rightarrow \infty$ and $\cos \psi_1(x) = 1 + O(e^{-2x})$ as $x \rightarrow \infty$.

(iv) $z < z_0 e^{-x}$ for $x > 0$, $z_0 = z(0, \gamma) = \sqrt{2(1 - \sin \gamma)}$.

(v) $z(x) \in C^2[0, \infty)$ if $0 < \gamma < \pi/2$.

Here $\sin \psi_1(x) = z_x/\sqrt{1+z_x^2}$.

Proof.

Part (i). From equation (20) x is defined for $0 < z < z_0$ and

$$(21) \quad x_z = \frac{z^2/2 - 1}{z\sqrt{1 - z^2/4}} .$$

Thus $x_z < 0$ for $0 < z < z_0$. Inverting, z is defined for $0 < x < \infty$ and $z_x < 0$. From equation (21):

$$(22) \quad \sin \psi_1 = -z\sqrt{1 - z^2/4}$$

and

$$(23) \quad \cos \psi_1 = 1 - z^2/2 .$$

We can rewrite equation (19a) as

$$-(\cos \psi_1)_z = z .$$

Thus, by equation (23) we see that z satisfies equation (19a). By equation (22) we see that z satisfies equation (19b). That z satisfies condition (19c) comes from equation (20).

Part (ii). We rewrite equation (20) as

$$(24) \quad \frac{e^x z}{C_1(\gamma)} = \frac{1}{2} (\sqrt{1 - z^2/4} + 1) \exp 2(1 - \sqrt{1 - z^2/4}) .$$

By Part (i) this quantity is $1 + o(1)$, thus $z = O(e^{-x})$; putting this back into equation (24) gives

$$\frac{e^x z}{C_1(\gamma)} = 1 + O(e^{-2x}) .$$

Part (iii). These estimates follow from Part (ii) and equations (22) and (23).

Part (iv). Let $v(x) = z_0 e^{-x}$, then

$$Nv = \frac{z_0 e^{-x}}{(1 + z_0^2 e^{-2x})^{3/2}} < z_0 e^{-x} = v$$

$v(0) = z(0)$, and $z - v = o(1)$ as $x \rightarrow \infty$. By CP $z < v$ for $x > 0$.

Part (v). For $0 < \gamma < \pi/2$, equation (21) shows that $z_x \in C[0, \infty)$ and by equation (19a) $z_{xx} = z(1 + z_x^2)^{3/2} \in C[0, \infty)$.

REMARK. If $\gamma = \pi/2$, $z(x, \pi/2) \equiv 0$ is the unique solution to problem (21). This is an immediate consequence of CP.

Chapter II Solutions in Symmetric Domains

6. General estimates—interior case. We obtain estimates for the function $v(r; R, \gamma)$ that satisfies

$$(25a) \quad \frac{1}{r^{n-1}} \left(\frac{r^{n-1} v_r}{\sqrt{1 + v_r^2}} \right)_r = v \quad \text{for } 0 < r < R$$

$$(25b) \quad v_r(0+) = 0$$

$$(25c) \quad v_r(R-) = \cot \gamma .$$

Johnson and Perko [12] prove by the method of successive approximations:

THEOREM 6. *Problem (25) has a unique solution $v(r, R, \gamma)$ which is continuous in (r, R, γ) for $0 \leq r \leq R$, $R > 0$, and $0 \leq \gamma \leq \pi/2$. If $0 \leq \gamma < \pi/2$ then $v > 0$, $v_r > 0$, and $v_{rr} > 0$ for $0 < r < R$. If $\gamma = \pi/2$ then $v \equiv 0$.*

Note. Johnson and Perko give the proof for $n = 2$, but the same proof holds for $n > 2$.

Introducing $\sin \psi(r) = v_r/\sqrt{1 + v_r^2}$ we have

$$(26a) \quad \frac{1}{r^{n-1}} (r^{n-1} \sin \psi)_r = v$$

$$(26b) \quad \sin \psi(0+) = 0$$

$$(26c) \quad \sin \psi(R-) = \cos \gamma .$$

Equation (26a) can be rewritten as

$$(27) \quad (\sin \psi)_r + \frac{(n-1)}{r} \sin \psi = v .$$

Partially inverting equation (27):

$$(28) \quad -(\cos \psi)_v + \frac{(n-1)}{r} \sin \psi = v .$$

We shall estimate $\sin \psi/r$ and then use equations (28) to estimate v .

LEMMA 5.

$$\int_{r_1}^{r_2} r^{n-1} v(r) dr = r_2^{n-1} \sin \psi(r_2) - r_1^{n-1} \sin \psi(r_1) .$$

Proof. Multiply equation (26a) by r^{n-1} and integrate.

LEMMA 6.

$$(i) \quad \frac{\sin \psi}{r} < \frac{v}{n} \quad \text{for } 0 < r \leq R$$

$$(ii) \quad \left(\frac{\sin \psi}{r} \right)_r > 0 \quad \text{for } 0 < r \leq R .$$

Proof.

Part (i). From Lemma 5 and Theorem 6:

$$r^{n-1} \sin \psi = \int_0^r s^{n-1} v(s) ds < \frac{r^n}{n} v(r) .$$

Part (ii).

$$\left(\frac{\sin \psi}{r} \right)_r = \frac{r(\sin \psi)_r - \sin \psi}{r^2} = \frac{-n \sin \psi + rv}{r^2} > 0 .$$

The last equality comes from equation (27); the inequality comes from Part (i).

LEMMA 7.

$$\frac{v(0)}{n} < \frac{\sin \psi}{r} < \min \left\{ \frac{v}{n}, \frac{\cos \gamma}{R} \right\}$$

for $0 < r < R$.

Proof. From Lemma 6 (ii)

$$\lim_{s \rightarrow 0+} \frac{\sin \psi(s)}{s} < \frac{\sin \psi(r)}{r} < \lim_{s \rightarrow R-} \frac{\sin \psi(s)}{s} .$$

We note that

$$\lim_{s \rightarrow 0+} \frac{\sin \psi(s)}{s} = \lim_{s \rightarrow 0+} [\sin \psi(s)]_s .$$

Thus, using equation (27):

$$n \cdot \lim_{s \rightarrow 0+} \frac{\sin \psi(s)}{s} = \lim_{s \rightarrow +0} v(s) = v(0) .$$

By condition (26b)

$$\lim_{s \rightarrow R-} \frac{\sin \psi(s)}{s} = \frac{\cos \gamma}{R} .$$

We have shown

$$\frac{v(0)}{n} < \frac{\sin \psi(r)}{r} < \frac{\cos \gamma}{R} .$$

Combining this inequality with Part (i) of Lemma 6 gives the stated inequality.

THEOREM 7. For $0 \leq \gamma < \pi/2$:

$$(i) \quad \max \left\{ \sqrt{2(1 - \sin \gamma)}, \frac{n \cos \gamma}{R} \right\} < v(R, R, \gamma) .$$

- (ii) $v(R, R, \gamma) < (n-1) \frac{\cos \gamma}{R} + \sqrt{2(1 - \sin \gamma) + \frac{\cos^2 \gamma}{R^2}}$
- (iii) $2(1 - \sin \gamma) + \frac{2(n-1)}{n} [v(R) - v(0)]v(0) < v^2(R) - v^2(0) < 2n(1 - \sin \gamma).$

Proof. Parts (i) and (iii) are due to Finn [5]. Integrate equation (28) from $v(0)$ to $v(R)$:

$$(29) \quad \frac{v^2(R) - v^2(0)}{2} = 1 - \sin \gamma + (n-1) \int_{v(0)}^{v(R)} \frac{\sin \psi}{r} dv.$$

Employ the estimates of Lemma 7:

$$(30) \quad \int_{v(0)}^{v(R)} \frac{\sin \psi}{r} dv > \frac{v(0)}{n} [v(R) - v(0)]$$

$$(31) \quad \int_{v(0)}^{v(R)} \frac{\sin \psi}{r} dv < \int_{v(0)}^{n \cos \gamma / R} \frac{v}{n} dv + \int_{n \cos \gamma / R}^{v(R)} \frac{\cos \gamma}{R} dv \\ = -\frac{n}{2} \left(\frac{\cos \gamma}{R} \right)^2 - \frac{v^2(0)}{2n} + v(R) \frac{\cos \gamma}{R}.$$

Part (i). Equation (29) gives $v(R) > \sqrt{2(1 - \sin \gamma)}$; *Part (i)* of Lemma 6 with $r = R$ gives $v(R) > n \cos \gamma / R$.

Part (ii). Combining equation (29) and inequality (31) gives

$$\frac{v^2(R)}{2} < 1 - \sin \gamma + \frac{1}{2n} v^2(0) - \frac{n(n-1) \cos^2 \gamma}{2R^2} \\ + v(R)(n-1) \frac{\cos \gamma}{R}.$$

Solving this quadratic inequality:

$$v(R) < (n-1) \frac{\cos \gamma}{R} + \left[2(1 - \sin \gamma) + (1-n) \frac{\cos^2 \gamma}{R^2} + \frac{v^2(0)}{n} \right]^{1/2}.$$

Estimating $v(0) < n(\cos \gamma / R)$ gives the stated inequality.

Part (iii). Combining equation (29) and inequality (30) gives the left-hand side of inequality (iii). The right-hand side is obtained by employing $\sin \psi / r < v/n$; this estimate yields

$$\int_{v(0)}^{v(R)} \frac{\sin \psi}{r} dv < \frac{v^2(R) - v^2(0)}{2n}.$$

Combining this with equation (29) gives the result.

Inequalities (i) and (ii) are sharp for small and large R , more precisely:

COROLLARY.

$$v(R, R, \gamma) = z(0, \gamma) + O(1/R) \quad \text{as } R \longrightarrow \infty$$

$$v(R, R, \gamma) = n \frac{\cos \gamma}{R} + O(R) \quad \text{as } R \longrightarrow 0 .$$

7. The narrow tube. A study is made of $v(r, R, \gamma)$ for small R . We show that the solution is close to a spherical cap.

First, we prove a technical lemma:

LEMMA 8. *Suppose $Nf(r) \geq Ng(r)$ for $0 \leq r \leq R$, then $f(r_2) - f(r_1) \geq g(r_2) - g(r_1)$ for $0 \leq r_1 \leq r_2 \leq R$.*

Proof.

$$\frac{1}{r^{n-1}} \left(\frac{r^{n-1} f_r}{\sqrt{1 + f_r^2}} \right)_r \geq \frac{1}{r^{n-1}} \left(\frac{r^{n-1} g_r}{\sqrt{1 + g_r^2}} \right)_r .$$

Multiply by r^{n-1} and integrating from 0 to r gives

$$\frac{r^{n-1} f_r}{\sqrt{1 + f_r^2}} \geq \frac{r^{n-1} g_r}{\sqrt{1 + g_r^2}} .$$

Simplifying: $f_r \geq g_r$. The conclusion follows upon integrating from r_1 to r_2 .

We take $0 \leq \gamma < \pi/2$ in what follows. Introduce

$$S_1(r) = R \tan \gamma - \sqrt{(R \sec \gamma)^2 - r^2} + n \frac{\cos \gamma}{R}$$

and

$$S_2(r) = S_1(r) + R \sec \gamma (1 - \sin \gamma) .$$

LEMMA 9. $S_1(r) < v(r) < S_2(r)$ for $0 \leq r < R$.

Proof. To apply CP, we check that

$$NS_1 = n \frac{\cos \gamma}{R} \geq S_1(r)$$

$$NS_2 = n \frac{\cos \gamma}{R} \leq S_2(r)$$

for $0 \leq r < R$, and $TS_1 \cdot \nu = TS_2 \cdot \nu = \cos \gamma$ for $r = R$.

Two special cases of Lemma 9 are

$$(32) \quad n \frac{\cos \gamma}{R} \leq v(R, R, \gamma) \leq R \sec \gamma(1 - \sin \gamma) + n \frac{\cos \gamma}{R}$$

$$(33) \quad n \frac{\cos \gamma}{R} > v(0, R, \gamma) > n \frac{\cos \gamma}{R} - R \sec \gamma(1 - \sin \gamma) .$$

Introduce $R_1 = R \sec \gamma[1 - (R^2/n) \sec^2 \gamma(1 - \sin \gamma)]^{-1}$ and $R_2 = R \sec \gamma[1 + (R^2/n) \sec^2 \gamma(1 - \sin \gamma)]^{-1}$.

THEOREM 8. $(0 \leq \gamma < \pi/2)$ $v(r, R, \gamma) > S_3(r) = H_2(R, \gamma) + R_1 - \sqrt{R_1^2 - r^2}$ and $v(r, R, \gamma) = S_3(r) + O(R^3)$ as $R \rightarrow 0$, where

$$H_2(R, \gamma) = n \frac{\cos \gamma}{R} - R_2 + \frac{n}{R^n} \int_0^R r^{n-1} \sqrt{R_2^2 - r^2} dr$$

for $\gamma \neq 0$, and

$$H_2(R, 0) = \frac{n}{R} \left(1 + \frac{R^2}{n}\right) - R \left(1 + \frac{R^2}{n}\right)^n - R_2 + \frac{n}{R_2^n} \int_0^{R_2} r^{n-1} \sqrt{R_2^2 - r^2} dr .$$

Proof.

Case $\gamma \neq 0$. Let $f_1(r) = R_1 - \sqrt{R_1^2 - r^2}$ and $f_2(r) = R_2 - \sqrt{R_2^2 - r^2}$. Clearly, $R_1 > R$ and $R_2 > R$ for $R \leq R_0$, with R_0 sufficiently small. By estimates (32) and (33)

$$Nf_1(r) = \frac{n}{R_1} < v(0) \leq v(r) = Nv$$

$$Nf_2(r) = \frac{n}{R_2} \geq v(R) \geq v(r) = Nv .$$

Hence, by Lemma 8

$$(34) \quad R_1 - \sqrt{R_1^2 - r^2} < v(r) - v(0) < R_2 - \sqrt{R_2^2 - r^2} .$$

Let

$$H_1(R, \gamma) = n \frac{\cos \gamma}{R} - R_1 + \frac{n}{R^n} \int_0^R r^{n-1} \sqrt{R_1^2 - r^2} dr .$$

Multiplying estimate (34) by r^{n-1} and integrating from 0 to R gives

$$(35) \quad H_2 < v(0) < H_1 .$$

Let

$$S_3(r) = H_2 + f_1(r)$$

and

$$S_4(r) = H_1 + f_2(r) .$$

By inequalities (34) and (35): $S_3(r) < v(r) < S_4(r)$. Now

$$S_4(r) - S_3(r) = R_2 - R_1 + \frac{n}{R^n} \int_0^R r^{n-1} (\sqrt{R_1^2 - r^2} - \sqrt{R_2^2 - r^2}) dr \\ + (\sqrt{R_1^2 - r^2} - \sqrt{R_2^2 - r^2}) .$$

Clearly $R_2 - R_1 = O(R^3)$ as $R \rightarrow 0$, and

$$\sqrt{R_1^2 - r^2} - \sqrt{R_2^2 - r^2} = \frac{(R_1 - R_2)(R_1 + R_2)}{\sqrt{R_1^2 - r^2} + \sqrt{R_2^2 - r^2}} = O(R^3)$$

as $R \rightarrow 0$. Hence $S_4(r) - S_3(r) = O(R^3)$ as $R \rightarrow 0$.

Case $\gamma = 0$. We must modify the above proof. We still have $v(0) < H_1(R, 0)$ and $R_1 - \sqrt{R_1^2 - r^2} < v(r) - v(0)$; however, $R_2 < R$ and $v(r) - v(0) < R_2 - \sqrt{R_2^2 - r^2}$ only for $0 \leq r < R_2$. We obtain

$$(36) \quad v(0) > \frac{n}{R^2} \sin \psi(R_2) - R_2 + \frac{n}{R_2^n} \int_0^{R_2} r^{n-1} \sqrt{R_2^2 - r^2} dr .$$

Here

$$R_1 = R \left(1 - \frac{R^2}{n}\right)^{-1} \quad \text{and} \quad R_2 = R \left(1 + \frac{R^2}{n}\right)^{-1} .$$

We estimate $\sin \psi(R_2)$; Lemma 5 gives

$$\sin \psi(R_2) = \left(\frac{R}{R_2}\right)^{n-1} - \frac{1}{R_2^{n-1}} \int_{R_2}^R r^{n-1} v(r) dr .$$

Thus

$$\sin \psi(R_2) > \left(\frac{R}{R_2}\right)^{n-1} - \left(\frac{n}{R} + R\right) \frac{R^n - R_2^n}{nR_2^{n-1}} \\ = 1 - \frac{R^2}{n} \left(1 + \frac{R^2}{n}\right)^{n-1} .$$

Putting this into estimate (36) gives $v(0) > H_2(R, 0)$. Thus $v(r) > S_3(r)$. Now

$$H_2(R, 0) - H_1(R, 0) = R_1 - R_2 + R \left[1 - \left(1 + \frac{R^2}{n}\right)^n\right] \\ + n \int_0^{R_2} r^{n-1} \left(\frac{\sqrt{R_2^2 - r^2}}{R_2^n} - \frac{\sqrt{R_1^2 - r^2}}{R^n}\right) dr$$

$$\begin{aligned}
 & + \frac{n}{R^n} \int_{R_2}^R r^{n-1} \sqrt{R_1^2 - r^2} dr \\
 & = O(R^3) \quad \text{as } R \longrightarrow 0 .
 \end{aligned}$$

We estimate $v(R, R, \gamma)$ by Part (iii) of Theorem 7:

$$(37) \quad v(R) - v(0) < \frac{2n}{v(R) + v(0)} < \frac{2n}{\frac{n}{R} + \frac{n}{R} - R} = R \left(1 - \frac{R^2}{2n}\right)^{-1}$$

$$\begin{aligned}
 (38) \quad v(R) - v(0) & > \left[\frac{v(R) + v(0)}{2} - \frac{(n-1)}{n} v(0) \right]^{-1} \\
 & > R \left[1 + R^2 \left(1 - \frac{1}{n}\right) \right]^{-1} .
 \end{aligned}$$

We have employed estimates (32) and (33) for $v(R)$ and $v(0)$. Estimates (37) and (38) show that $v(R) = v(0) + R + O(R^3)$ as $R \rightarrow 0$.

Since $Nv = v > NS_s$ for $0 \leq r < R$, Lemma 8 gives $v(R) - v(r) > S_s(R) - S_s(r)$. Thus $v(r) < v(R) - S_s(R) + S_s(r)$. We note that $S_s(R) - S_s(0) = R_1 - \sqrt{R_1^2 - R^2} = R + O(R^3)$ as $R \rightarrow 0$. Therefore

$$\begin{aligned}
 v(r) & < [v(0) + R] - [S_s(0) + R] + S_s(r) + O(R^3) \\
 & = S_s(r) + [v(0) - S_s(0)] + O(R^3) \\
 & < S_s(r) + O(R^3) \quad \text{as } R \longrightarrow 0 .
 \end{aligned}$$

We have used that $v(0) - S_s(0) = v(0) - H_2(R, 0) < H_1(R, 0) - H_2(R, 0) = O(R^3)$. Since we started with $S_s(r) < v(r)$ the proof is complete.

COROLLARY. For $n = 2$, $H_2(R, \gamma)$ can be calculated explicitly, yielding

$$(39) \quad v(0, R, \gamma) = 2 \frac{\cos \gamma}{R} - \frac{R}{\cos \gamma} \left(1 - \frac{2}{3} \frac{1 - \sin^3 \gamma}{\cos^2 \gamma} \right) + O(R^3)$$

as $R \rightarrow 0$.

REMARK. The formula (39) is known as Laplace's formula. Ferguson has given a formula differing from formula (39) [1]. We have settled the question.

8. The wide tube.

8A. Estimate near the boundary. We show that near the boundary the solution v approaches the one-dimensional solution as $R \rightarrow \infty$.

LEMMA 10. For $0 \leq \gamma < \pi/2$ and $0 \leq l \leq R$:

$$(40) \quad v^2(R - l) < v^2(R) + 2 \sin \gamma - 2 \cos \psi(R - l)$$

$$(41) \quad \sin \psi(R - l) < -\sin \psi_1(l) + \frac{l}{R} (n - 1) \cos \gamma .$$

Proof. Integrating equation (28) from $v(R - l)$ to $v(R)$ and estimating $\sin \psi > 0$ gives inequality (40).

By CPS, $z(x_1, \gamma) < v(|x - y|, R, \gamma)$ in $B_R(y)$ if $B_R(y) \subset H$. Choose $y = (R, 0, \dots, 0)$ and $x = (x_1, 0, \dots, 0)$. Thus

$$(42) \quad z(x_1, \gamma) < v(R - x_1, R, \gamma) \quad \text{for } 0 < x_1 \leq R .$$

Let $r = R - x_1$. Then $[\sin \psi_1(x_1)]_{x_1} = -[\sin \psi(x_1)]_r$. Therefore $(\sin \psi)_r + (n - 1)(\sin \psi/r) > -[\sin \psi_1(x_1)]_r$. Integrating from $R - l$ to R and estimating $\sin \psi/r < \cos \gamma/R$ gives inequality (41).

THEOREM 9. For $0 \leq \gamma < \pi/2$ and $l \geq 0$:

(i) $v(R - l, R, \gamma)$ is strictly decreasing in R .

(ii) $\lim_{R \rightarrow \infty} v(R - l, R, \gamma) = z(l, \gamma)$.

Proof.

Part (i). By CPS, $v(|x - y_2|, R_2, \gamma) < v(|x - y_1|, R_1, \gamma)$ in $B_{R_1}(y_1)$ if $B_{R_1}(y_1) \subset B_{R_2}(y_2)$. Choosing $y_i = (R_i, 0, \dots, 0)$ $i = 1$ or 2 , and $x = (l, 0, \dots, 0)$ gives $v(R_2 - l, R_2, \gamma) < v(R_1 - l, R_1, \gamma)$ for $R_1 < R_2$ and $0 < l \leq R_1$.

For $l = 0$ we have $v(R_2, R_2, \gamma) \leq v(R_1, R_1, \gamma)$. If $\gamma \neq 0$ Lemma 1 rules out equality.

A special argument is needed for the case $\gamma = 0$ and $l = 0$: Let $u_1(r) = v(r, R_1, 0)$ and $u_2(r) = v(r + R_2 - R_1, R_2, 0)$. Suppose $v(R_2, R_2, 0) = v(R_1, R_1, 0)$, then $u_1(R_1) = u_2(R_1)$. Also $u_1(r) > u_2(r)$ for $0 \leq r < R_1$. At R_1

$$(\sin \varphi_1)_r + \frac{(n - 1)}{R_1} = (\sin \varphi_2)_r + \frac{(n - 1)}{R_2}$$

since $Nu_1 = Nu_2$ at R_1 ; here φ_i ($i = 1$ or 2) is the angle between the curve given by $u_i(r)$ and the positive r direction. Thus, $(\sin \varphi_1)_r < (\sin \varphi_2)_r$ at R_1 . This inequality must hold for $R_1 - \varepsilon \leq r \leq R_1$, for some $\varepsilon > 0$. Integrating from r to R_1 gives $\sin \varphi_1(r) > \sin \varphi_2(r)$; this implies that $u_1(r) < u_2(r)$ for $R_1 - \varepsilon \leq r \leq R_1$ after another integration. This is a contradiction.

Part (ii). For R such that

$$(43) \quad R > \frac{(n + 1)l \cos \gamma}{1 + \sin \psi_1(l)}$$

the right-hand member of inequality (41) is < 1 . Combining inequalities (40) and (41) and noting by equation (23) that $2[\sin \gamma - \cos \psi_1(l)] = z^2(l) - z^2(0)$ gives

$$(44) \quad v^2(R - l) < v^2(R) - z^2(0) + z^2(l) + 2\eta(l, R) \\ \eta(l, R) = \cos \psi_1(l) - [\cos^2 \psi_1(l) + 2\varepsilon \sin \psi_1(l) - \varepsilon^2]^{1/2}$$

where $\varepsilon = (n - 1)(l/R) \cos \gamma$. Clearly, $\lim_{R \rightarrow \infty} \eta(l, R) = 0$. By the corollary to Theorem 7 $\lim_{R \rightarrow \infty} v^2(R) = z^2(0)$. Thus $\lim_{R \rightarrow \infty} v(R - l, R, \gamma) \leq z(l, \gamma)$. By inequality (42) $v(R - l, R, \gamma) \geq z(l, \gamma)$. Hence Part (ii) is proved.

8B. *Estimate of $v(r, R, \gamma)$ from above.*

THEOREM 10. *For $0 \leq \gamma < \pi/2$: $v(r, R, \gamma) < H(R, \gamma)I(r)[1 + O(1/R^{1/2})]$ for $0 \leq r \leq R - (1/4) \ln R$ as $R \rightarrow \infty$, where*

$$H(R, \gamma) = \sqrt{2\pi}C_1(\gamma) \frac{R^{(n-1)/2}}{\exp R} .$$

Proof. By CP and Lemma 3, $v(r, R) < (I(r)/I(R - l))v(R - l, R)$ for $0 \leq r < R - l$. By inequality (44)

$$v(R - l) < z(l)\{1 + [2z(l)]^{-2}[v^2(R) - z^2(0) + 2\eta(l, R)]\}^{1/2} .$$

Choose $l = (1/4) \ln R$, for R sufficiently large condition (43) shows that η is defined. By the estimates of Chapter I we find

$$I(R - l) = \frac{1}{\sqrt{2\pi}} \frac{\exp(R - l)}{R^{(n-1)/2}} \left[1 + O\left(\frac{\ln R}{R}\right) \right]$$

and

$$z(l) = C_1(\gamma) \exp(-l) \left[1 + O\left(\frac{1}{R^{1/2}}\right) \right] .$$

To estimate η , we note that

$$0 < \eta(l, R) < \varepsilon[-2 \sin \psi_1(l) + \varepsilon][\cos \psi_1(l)]^{-1} .$$

Thus

$$\eta(l, R) = O\left(\frac{\ln R}{R^{5/4}}\right) .$$

Also $v^2(R) - z^2(0) = O(1/R)$ by the corollary to Theorem 7. Combining estimates and noting that $z(l) = O(1/R^{1/4})$ gives the result.

8C. *Estimate of $v(r, R, \gamma)$ from below.*

THEOREM 11. *For $0 \leq \gamma < \pi/2$: $v(r, R, \gamma) > H(R, \gamma)I(r)[1 + O(\ln R/R)]$*

for $0 \leq r \leq R - (5/2) \ln R$ as $R \rightarrow \infty$.

Proof. By Theorem 9, $v(R - l, R, \gamma) > z(l, \gamma)$ for $0 \leq l \leq R$. We construct a subsolution $u(r)$ for $0 \leq r \leq R - l$ with $u(R - l) = z(l, \gamma)$, where l will be chosen later. Let

$$u(r) = h \left(1 - \frac{\varepsilon}{r^2 + a} \right) I(r) = hB(r)I(r)$$

where h, ε , and a are positive constants with $\varepsilon < a$.

$$u_r = h \left[B(r)I_r + \frac{2\varepsilon r}{(r^2 + a)^2} I \right].$$

We note: $u > 0$ and $u_r \geq 0$. The subsolution condition $Nu \geq u$ becomes

$$u_{rr} + \frac{(n-1)}{r} u_r + \frac{(n-1)}{r} u_r^3 \geq u(1 + u_r^2)^{3/2}.$$

We require the stronger inequalities

$$(45a) \quad u_{rr} + \frac{(n-1)}{r} u_r \geq u(1 + 2u_r^2)$$

$$(45b) \quad \max u_r < 1.$$

These are stronger because $(1 + x)^{3/2} \leq 1 + 2x$ for $0 \leq x \leq 1$. Condition (45a) reduces to

$$(46) \quad u_r^2 < \frac{\varepsilon AB^{-1}}{(r^2 + a)^2}$$

$$A(r) = 2r \frac{I_r}{I} + \frac{a - 3r^2}{r^2 + a} + (n - 1).$$

By Lemma 4 we can choose b so that

$$b \frac{I_r(b)}{I(b)} \geq \frac{5 - n}{2}.$$

Let $a = 3b^2$. With this choice for a , $A(r) > 1$ for all $r \geq 0$, as can be seen by checking the cases $r > b$ and $r < b$. Choose $h = z(l)[I(R - l)B(R - l)]^{-1}$. Let

$$c = \max_{r>0} \frac{2r}{(r^2 + a)^2}.$$

By Lemma 4 we can estimate $I_r < I$, thus

$$\begin{aligned}
 \max_{0 \leq r \leq R-l} u_r &< h[B(R-l) + \varepsilon c]I(R-l) \\
 &= z(l)[1 + \varepsilon cB^{-1}(R-l)] \\
 (47) \quad &< z(l) \left[1 + \varepsilon c \left(1 - \frac{\varepsilon}{a} \right)^{-1} \right] \\
 \min_{0 \leq r \leq R-l} \frac{\varepsilon AB^{-1}}{(r^2 + a)^2} &> \frac{\varepsilon B^{-1}(R)}{(R^2 + a)^2}.
 \end{aligned}$$

We can satisfy condition (46) by choosing l so that

$$(48) \quad 2e^{-2l} \left[1 + \varepsilon c \left(1 - \frac{\varepsilon}{a} \right)^{-1} \right]^2 < \frac{\varepsilon B^{-1}(R)}{(R^2 + a)^2}.$$

We have estimated $z(l) < \sqrt{2}e^{-l}$. Finally, choose $\varepsilon = \ln R/R$ and $l = (5/2) \ln R$, then $l > (1/2) \ln [2R(R^2 + a)^2/\ln R] + o(1)$ as $R \rightarrow \infty$. Thus condition (48) is satisfied for R sufficiently large. With estimate (47) we check that condition (45b) holds for R sufficiently large.

By CP, $u(r) < v(r)$ for $0 \leq r \leq R - l$. Thus

$$\begin{aligned}
 v(r) &> hB(r)I(r) \\
 &> \frac{z(l)}{I(R-l)} B(0)B^{-1}(R-l)I(r).
 \end{aligned}$$

By the estimates in Chapter I we find

$$\begin{aligned}
 z(l) &= C_1(\gamma)e^{-l} \left[1 + O\left(\frac{1}{R^{5/2}}\right) \right] \\
 I(R-l) &= \frac{1}{\sqrt{2\pi}} \frac{\exp(R-l)}{R^{(n-1)/2}} \left[1 + O\left(\frac{\ln R}{R}\right) \right].
 \end{aligned}$$

Combining estimates and noting that $B^{-1}(R-l) = 1 + O(\ln R/R)$ gives the result.

COROLLARY 1. $v(r, R, \gamma) = H(R, \gamma)I(r)[1 + O(1/R^{1/2})]$ for $0 \leq r \leq R - (5/2) \ln R$ as $R \rightarrow \infty$.

Proof. Combine Theorems 10 and 11.

We can estimate the first derivative as follows:

COROLLARY 2. For $0 \leq r \leq R - (5/2) \ln R$

$$\sin \psi(r) = H(R, \gamma)I_r(r) \left[1 + O\left(\frac{1}{R^{1/2}}\right) \right]$$

for $R \rightarrow \infty$.

Proof. By Lemma 5,

$$\sin \psi(r) = \frac{1}{r^{n-1}} \int_0^r s^{n-1} v(s) ds .$$

Estimating $v(r)$ with Theorem 10 gives

$$\sin \psi(r) < H(R, \gamma) I_r(r) \left[1 + O\left(\frac{1}{R^{1/2}}\right) \right]$$

for $0 \leq r \leq R - (1/4) \ln R$. Estimating $v(r)$ with Theorem 11 gives

$$\sin \psi(r) > H(R, \gamma) I_r(r) \left[1 + O\left(\frac{\ln R}{R}\right) \right]$$

for $0 \leq r \leq R - (5/2) \ln R$. Combining estimates gives the result.

REMARK. One expects the error in Corollaries 1 and 2 to be $O(\ln R/R)$. Perko [15] obtains this error estimate for $\gamma > 0$ by a completely different method. The case $\gamma = 0$ is important in view of the general estimates of Chapter III.

9. The exterior problem. We obtain estimates for the function $w(r, R, \gamma)$ which satisfies

$$(49a) \quad \frac{1}{r^{n-1}} \left(\frac{r^{n-1} w_r}{\sqrt{1 + w_r^2}} \right)_r = w \quad \text{for } r > R$$

$$(49b) \quad w_r(R+) = -\cot \gamma$$

$$(49c) \quad \lim_{r \rightarrow \infty} w(r) = 0 .$$

Johnson and Perko [12] prove by a “shooting argument”:

THEOREM 12. *Problem (49) has a unique solution $w(r, R, \gamma)$ which is continuous in (r, γ) for $r \geq R$ and $0 < \gamma \leq \pi/2$; $w > 0$, $w_r < 0$ if $0 \leq \gamma < \pi/2$; $w \equiv 0$ if $\gamma = \pi/2$; $w(r, R, 0) = \lim_{r \rightarrow 0+} w(r, R, \gamma)$; and $\lim_{r \rightarrow \infty} w_r = 0$.*

REMARK. The note after Theorem 6 applies here as well.

9A. Continuity with respect to (r, R, γ) . We fill in a gap in the continuous dependence properties.

LEMMA 11. *For $0 \leq \gamma < \pi/2$, $R_1 < R_2$ and $r > R_2$*

$$(50) \quad w(r + R_1 - R_2, R_1, \gamma) < w(r, R_2, \gamma) < \frac{R_2}{R_1} w\left(\frac{R_1}{R_2} r, R_1, \gamma\right) .$$

Proof. Let $v_i = w(r_i, R_i, \gamma)$ and $r_i = |x - y_i|$ for $i = 1$ or 2 . If

$B_{R_2}^c(y_2) \subset B_{R_1}^c(y_1)$ then the proof of Theorem 2 shows that $Tv_1 \cdot \nu \leq \cos \gamma = Tv_2 \cdot \nu$ on $\partial B_{R_2}^c$. Since $v_1 = o(1)$ and $v_2 = o(1)$ as $|x| \rightarrow \infty$, CP gives $v_1 < v_2$ on $B_{R_1}^c$. Choose $y_2 = 0$, $y_1 = (R_2 - R_1, 0, \dots, 0)$, and $x = (x_1, 0, \dots, 0)$; thus $w(x_1 + R_1 - R_2, R_1, \gamma) < w(x_1, R_2, \gamma)$ for $x_1 > R_2$.

Let

$$u = \frac{R_2}{R_1} w\left(\frac{R_1}{R_2} r, R_1, \gamma\right).$$

Then by Lemma 2

$$Nu = \left(\frac{R_1}{R_2}\right)^2 u < u$$

in $B_{R_2}^c(0)$ and $Tu \cdot \nu = \cos \gamma$ on $\partial B_{R_2}^c$. Thus by CP, $u > w$ in $B_{R_2}^c(0)$.

LEMMA 12. For $0 \leq \gamma \leq \pi/2$:

$$0 < \varepsilon < \sqrt{2}, \quad \sigma(\varepsilon) = \frac{\varepsilon^3}{3}.$$

If $|r_1 - r_2| < \sigma(\varepsilon)$ and $r_1, r_2 \geq R$ then $|w(r_2, R, \gamma) - w(r_1, R, \gamma)| < \varepsilon$.

Proof. See [5].

THEOREM 13. For $0 \leq \gamma \leq \pi/2$: $w(r, R, \gamma)$ is continuous in (r, R) independent of γ .

Proof. We show continuity at (r_1, R_1) .

For $R > R_1$: By inequality (50)

$$w(r, R_1, \gamma) < w(r, R, \gamma) < \frac{R}{R_1} w\left(\frac{R_1}{R} r, R_1, \gamma\right).$$

Suppose

$$(51) \quad \left. \begin{aligned} |r - r_1| &< \sigma(\varepsilon) \\ \left| \frac{R_1}{R} r - r_1 \right| &< \sigma(\varepsilon) \end{aligned} \right\}.$$

By Lemma 12

$$(52) \quad w(r_1, R_1, \gamma) - \varepsilon < w(r, R, \gamma) < \frac{R}{R_1} [w(r_1, R_1, \gamma) + \varepsilon].$$

For $R < R_1$: By inequality (50)

$$\frac{R}{R_1} w\left(\frac{R_1}{R} r, R_1, \gamma\right) < w(r, R, \gamma) < w(r, R_1, \gamma).$$

Supposing condition (51):

$$(53) \quad \frac{R}{R_1} [w(r_1, R_1, \gamma) - \varepsilon] < w(r, R, \gamma) < w(r_1, R_1, \gamma) + \varepsilon .$$

Inequalities (52) and (53) show continuity at (r_1, R_1) independent of γ .

COROLLARY. $w(r, R, \gamma)$ is continuous in (r, R, γ) for $r \geq R$, $R > 0$, and $0 \leq \gamma \leq \pi/2$.

Proof. We show continuity at (r_1, R_1, γ_1) .

Note:

$$\begin{aligned} |w(r, R, \gamma) - w(r_1, R_1, \gamma_1)| &\leq |w(r, R, \gamma) - w(r_1, R_1, \gamma)| \\ &\quad + |w(r_1, R_1, \gamma) - w(r_1, R_1, \gamma_1)| \\ &= I + II . \end{aligned}$$

For $|\gamma - \gamma_1| < \delta_1$: $II < \varepsilon$ by Theorem 12. For $|r - r_1| < \delta_2$ and $|R - R_1| < \delta_3$: $I < \varepsilon$ by Theorem 13. Hence $I + II < 2\varepsilon$.

9B. General estimates—exterior case. Introducing $\sin \varphi(r) = w_r/\sqrt{1 + w_r^2}$ we have

$$(54a) \quad \frac{1}{r^{n-1}} (r^{n-1} \sin \varphi)_r = w \quad \text{for } r > R$$

$$(54b) \quad \sin \varphi(R-) = -\cos \gamma .$$

Equation (54a) can be rewritten as

$$(55) \quad (\sin \varphi)_r + \frac{(n-1)}{r} \sin \varphi = w .$$

Partially inverting equation (55):

$$(56) \quad -(\cos \varphi)_w + \frac{(n-1)}{r} \sin \varphi = w .$$

LEMMA 13. For $0 \leq \gamma < \pi/2$ and $r > R$, $(\sin \varphi)_r > 0$ and

$$(57) \quad -\frac{\cos \gamma}{R} < \frac{\sin \varphi}{r} < 0 .$$

Proof. Since $\sin \varphi < 0$, $(\sin \varphi)_r = w - (n-1)/r \sin \varphi > 0$. Thus $-\cos \gamma < \sin \varphi(r) < 0$ for $r > R$ implying inequality (57).

THEOREM 14. For $0 \leq \gamma < \pi/2$:

(i) $w(R, R, \gamma) < \sqrt{2(1 - \sin \gamma)} .$

(ii) $w(R, R, \gamma) > -(n - 1)\frac{\cos \gamma}{R} + \sqrt{2(1 - \sin \gamma) + \left[\frac{(n - 1) \cos \gamma}{R}\right]^2} .$

Proof. Part (i) is due to Finn [5]. Integrate equation (56) from $w(r)$ to $w(R)$:

(58) $\frac{w^2(R) - w^2(r)}{2} = -\sin \gamma + \cos \varphi + (n - 1) \int_{w(r)}^{w(R)} \frac{\sin \varphi}{r} dw .$

Let $r \rightarrow \infty$:

$$w^2(R) = 2(1 - \sin \gamma) + 2(n - 1) \int_0^{w(R)} \frac{\sin \varphi}{r} dw .$$

Estimating with inequality (57) gives

$$w^2(R) < 2(1 - \sin \gamma)$$

$$w^2(R) > 2(1 - \sin \gamma) - 2(n - 1) \frac{\cos \gamma}{R} w(R) .$$

Solving this quadratic inequality gives part (ii).

COROLLARY. $w(R, R, \gamma) = \sqrt{2(1 - \sin \gamma)} + O(1/R)$ as $R \rightarrow \infty$.

9C. Estimate near the boundary.

LEMMA 14. For $0 \leq \gamma < \pi/2$:

(59) $w(r, R, \gamma) < z(r - R, \gamma) \quad \text{for } r > R$

(60) $\sin \varphi(R + l) < \sin \psi_1(l) + (n - 1) \frac{l \cos \gamma}{R} \quad \text{for } l > 0 .$

Proof. Let $u(x) = z(x_1 - R, \gamma)$. By the proof of Theorem 2: $Tu \cdot \nu \leq \cos \gamma = Tu \cdot \nu$ on $\{x: x_1 = R\}$. Since $w = o(1)$ and $u > 0$ we have $w - u \leq o(1)$ as $|x| \rightarrow \infty$. By CP,

$$w(r, R, \gamma) < z(x_1 - R, \gamma) \leq z(r - R, \gamma) \quad \text{for } r > R .$$

Let $x_1 = r - R$, then (cf. Lemma 10)

$$(\sin \varphi)_r + \frac{(n - 1)}{r} \sin \varphi < -[\sin \psi_1(x_1)]_r .$$

Integrating from R to $R + l$ and employing inequality (57) gives inequality (60).

THEOREM 15. For $0 \leq \gamma < \pi/2$ and $l \geq 0$:

- (i) $w(R + l, R, \gamma)$ is strictly increasing in R .
- (ii) $\lim_{R \rightarrow \infty} w(R + l, R, \gamma) = z(l, \gamma)$.

Proof.

Part (i). For $l > 0$ and $R_1 < R_2$, take $r = R_2 + l$ in Lemma 11: $w(R_1 + l, R_1, \gamma) < w(R_2 + l, R_2, \gamma)$. For $l = 0$ we have $w(R_1, R_1, \gamma) \leq w(R_2, R_2, \gamma)$. If $\gamma \neq 0$ Lemma 1 rules out equality. The case $l = 0$ and $\gamma = 0$ is handled as in the proof of Theorem 9.

Part (ii). From equation (58): $w^2(r) > w^2(R) + 2[\sin \gamma - \cos \varphi(r)]$. Let $r = R + l$ and combine this with inequality (60) noting that $2[\sin \gamma - \cos \psi_1(l)] = z^2(l) - z^2(0)$:

$$(61) \quad w^2(R + l) > w^2(R) - z^2(0) + z^2(l) + 2\eta(l, R)$$

where η is the function introduced in § 8. Now, by the corollary to Theorem 14 $\lim_{R \rightarrow \infty} w^2(R) = z^2(0)$. As before $\lim_{R \rightarrow \infty} \eta(l, R) = 0$, thus $\lim_{R \rightarrow \infty} w(R + l, R, \gamma) \geq z(l)$. By inequality (59): $w(R + l, R, \gamma) \leq z(l, \gamma)$. Thus $\lim_{R \rightarrow \infty} w(R + l, R, \gamma) \leq z(l, \gamma)$.

9D. Behavior at infinity. We study $w(r, R, \gamma)$ for r large.

THEOREM 16. For $0 \leq \gamma < \pi/2$:

- (i) $C(R, \gamma) = \lim_{r \rightarrow \infty} \frac{w(r, R, \gamma)}{K(r)}$ exists and $C(R, \gamma) > 0$.
- (ii) $w(r, R, \gamma) = C(R, \gamma)K(r)\{1 + O[K^2(r)]\}$ as $r \rightarrow \infty$.
- (iii) $C(R, \gamma) < \sqrt{2/\pi}C_1(\gamma)R^{(n-1)/2}e^R[1 + O(\ln R/R)]$
 $C(R, \gamma) > \sqrt{2/\pi}C_1(\gamma)R^{(n-1)/2}e^R[1 + O(1/R^{1/2})]$ as $R \rightarrow \infty$.

Proof.

Part (i). Since $K(r)$ is a supersolution (Lemma 3) CP gives

$$w(r) < \frac{w(R_1)}{K(R_1)} K(r) \quad \text{for } r > R_1.$$

Thus $w(r)/K(r)$ is monotone decreasing in r and positive. Hence $C(R, \gamma)$ exists and $C(R, \gamma) \geq 0$.

We now construct a subsolution. Let $u(r) = AK(r)[1 + aK^2(r)]$ for $r \geq R_1 \geq R$. Let $A = w(R_1, R, \gamma)\{K(R_1)[1 + aK^2(R_1)]\}^{-1}$. We will determine $a(R, R_1) > 0$ so that u satisfies the subsolution condition $Nu \geq u$ for $r \geq R_1$. By CP we will then have $u < v$ for $r > R_1$.

We calculate $u_r = AK_r(1 + 3aK^2)$ and note that $u_r < 0$. We estimate u_r by noting

$$\frac{w(R_1)}{K(R_1)} \leq \frac{w(R)}{K(R)} < \frac{\sqrt{2}}{K(R)} .$$

Thus

$$\begin{aligned} -u_r &< \sqrt{2} \left[-\frac{K_r(R)}{K(R)} \right] \frac{1 + 3aK^2(R_1)}{1 + aK^2(R_1)} \\ &< 3\sqrt{2} \left[\frac{(n-1)}{2R} + \sqrt{1 + \left(\frac{n-1}{2R}\right)^2} \right] = b(R) . \end{aligned}$$

The last inequality comes from Lemma 4. The subsolution condition becomes upon dividing by u :

$$(62) \quad 1 + 2a(K^2 + 3K_r^2)(1 + aK^2)^{-1} + \frac{(n-1)}{r} \frac{K_r}{K} \frac{(1 + 3aK^2)}{(1 + aK^2)} u_r^2 \geq (1 + u_r^2)^{3/2} .$$

We will require the stronger inequality obtained by replacing $(1 + u_r^2)^{3/2}$ with $1 + c(R)u_r^2$, with

$$c(R) = \frac{[b^2(R) + 1]^{3/2} - 1}{b^2(R)} .$$

We require

$$(63) \quad u_r^2 f < 2ag(1 + aK^2)^{-1}$$

where

$$\begin{aligned} f &= c + \frac{(n-1)}{r} \left(-\frac{K_r}{K} \right) \left(\frac{1 + 3aK^2}{1 + aK^2} \right) \\ g &= K^2 + 3K_r^2 . \end{aligned}$$

Rearranging condition (63);

$$(64) \quad 2a > A^2 K_r^2 f g^{-1} (1 + 3aK^2)^2 (1 + aK^2) .$$

As above,

$$A < \frac{\sqrt{2}}{K(R)} [1 + aK^2(R_1)]^{-1} .$$

Thus

$$A^2(1 + 3aK^2)(1 + aK^2) < \frac{6}{K^2(R)} .$$

Also

$$f < \bar{f} = c + \frac{n-1}{r} \left(-\frac{K_r}{K} \right) (3) .$$

Thus

$$\max_{r \geq R} K_r^2 f g^{-1} \leq \max_{r \geq R} K_r^2 \bar{f} g^{-1} = d(R) .$$

Choose $a = 3d(R)/[K^2(R) - 9d(R)K^2(R_1)]$ and condition (64) will be satisfied. We note that $b(R)$, $c(R)$ and $d(R)$ are decreasing in R .

Take R_1 sufficiently large, then $w(r) > u(r)$ for $r > R_1$ and thus

$$C(R, \gamma) = \lim_{r \rightarrow \infty} \frac{w(r)}{K(r)} \geq A > 0 .$$

Part (ii). From Part (i) $C(R, \gamma) < w(r)/K(r)$ or

$$(65) \quad C(R, \gamma)K(r) < w(r, R, \gamma) \quad \text{for } r > R .$$

Next, note

$$w(r) > \frac{w(R_1)}{K(R_1)} \frac{K(r)[1 + aK^2(r)]}{[1 + aK^2(R_1)]}$$

for $r \geq R_1 \geq R$. Divide by $K(r)$ and let $r \rightarrow \infty$:

$$(66) \quad C(R, \gamma) \geq \frac{w(R_1, R, \gamma)}{K(R_1)[1 + aK^2(R_1)]} .$$

Replacing R_1 by r gives

$$(67) \quad w(r, R, \gamma) < C(R, \gamma)K(r)[1 + a(R, r)K^2(r)] .$$

Combining estimates (65) and (67) gives Part (ii).

Part (iii). We find an *upper bound* as follows:

$$C(R, \gamma) < \frac{w(R_1, R, \gamma)}{K(R_1)} < \frac{z(l, \gamma)}{K(R_1)}$$

where $l = R_1 - R$. The second inequality comes from Lemma 14. Choose $l = (1/2) \ln R$, then by the estimates of Chapter I:

$$z(l) = C_1(\gamma)e^{-l} \left[1 + O\left(\frac{1}{R}\right) \right]$$

$$K(R_1) = \sqrt{\pi/2} \frac{\exp -(R + l)}{R^{(n-1)/2}} \left[1 + O\left(\frac{\ln R}{R}\right) \right] .$$

Thus

$$\frac{z(l)}{K(R_1)} = \sqrt{2/\pi} C_1(\gamma) e^{R} R^{(n-1)/2} \left[1 + O\left(\frac{\ln R}{R}\right) \right] .$$

We find a *lower bound* by combining inequalities (61) and (66) with $R_1 = R + l$:

$$C(R, \gamma) > \frac{z(l)}{K(R_1)} \frac{\{1 + z^{-2}(l)[w^2(R) - z^2(0) + 2\eta(l, R)]\}^{1/2}}{[1 + a(R, R_1)K^2(R_1)]} .$$

Choose $l = (1/4) \ln R$, then

$$z(l) = C_1(\gamma)e^{-l} \left[1 + O\left(\frac{1}{R^{1/2}}\right) \right] = O\left(\frac{1}{R^{1/4}}\right)$$

$$K^2(R_1)K^{-2}(R) = O\left(\frac{1}{R^{1/2}}\right)$$

$$\eta(l, R) = O\left(\frac{\ln R}{R^{5/4}}\right).$$

By the corollary to Theorem 14: $w^2(R) - z^2(0) = O(1/R)$. Noting that $d(R)$ is decreasing in R we have

$$C(R, \gamma) > \sqrt{2/\pi}C_1(\gamma)e^R R^{(n-1)/2} \left[1 + O\left(\frac{1}{R^{1/2}}\right) \right].$$

We can determine the asymptotic behavior of the derivatives of w :

COROLLARY.

- (i) $w_r = C(R, \gamma)K_r(r)\{1 + O[K^2(r)]\}$ as $r \rightarrow \infty$.
- (ii) $w_{rr} = C(R, \gamma)K_{rr}(r)\{1 + O[K^2(r)]\}$ as $r \rightarrow \infty$.

Proof.

Part (i). Integrate the equation $(r^{n-1} \sin \varphi)_r = r^{n-1}w$ from r to ∞ :

$$-\sin \varphi(r) = \frac{1}{r^{n-1}} \int_r^\infty s^{n-1}w(s)ds.$$

Estimate the integral with $w(s) \leq w(r)(K(s)/K(r))$ and

$$w(s) \geq \frac{w(r)K(s)[1 + aK^2(s)]}{K(r)[1 + aK^2(r)]} > \frac{w(r)K(s)}{K(r)[1 + aK^2(r)]}.$$

Note:

$$\int_r^\infty s^{n-1}K(s)ds = -r^{n-1}K_r(r).$$

Hence

$$-\frac{K_r}{K(1 + aK^2)} \cdot w \leq -\sin \varphi \leq -\frac{K_r}{K} w(r).$$

Thus $\sin \varphi = C(R, \gamma)K_r[1 + O(K^2)]$ as $r \rightarrow \infty$. This implies Part (i).

Part (ii):

$$\begin{aligned} \frac{w_{rr}}{(1 + w_r^2)^{3/2}} &= (\sin \varphi)_r = w - (n - 1) \frac{\sin \varphi}{r} \\ &= C(\gamma, R) \left[K - \frac{(n - 1)}{r} K_r \right] [1 + O(K^2)] \\ &= C(\gamma, R) K_{rr} [1 + O(K^2)] . \end{aligned}$$

We give some further properties of $C(R, \gamma)$:

THEOREM 17.

- (i) $C(R, \gamma)$ is continuous in (R, γ) for $R > 0$ and $0 \leq \gamma \leq \pi/2$.
- (ii) $C(R, \gamma)$ is strictly increasing in R for $0 \leq \gamma < \pi/2$; $C(R, \gamma)$ is strictly decreasing in γ .

Proof.

Part (i). We show continuity at (R_0, γ_0) . For $\gamma_0 < \pi/2$ we combine inequalities (65) and (66):

$$\begin{aligned} [1 + a(R)K^2(R_1)]^{-1} \frac{w(R_1, R, \gamma)}{w(R_1, R_0, \gamma_0)} &\leq \frac{C(R, \gamma)}{C(R_0, \gamma_0)} \\ &\leq \frac{w(R_1, R, \gamma)}{w(R_1, R_0, \gamma_0)} [1 + a(R_0)K^2(R_1)] . \end{aligned}$$

Require $|R - R_0| < R_0/2$, then choose R_1 so that $a(R_0)K^2(R_1) < \varepsilon$ and $a(R)K^2(R_1) < \varepsilon$. Here $a(R_0) = a(R_0, R_1)$ and $a(R) = a(R, R_1)$. By the corollary to Theorem 13 there are $\delta_1 > 0$ and $\delta_2 > 0$ so that

$$\left| \frac{w(R_1, R, \gamma)}{w(R_1, R_0, \gamma_0)} - 1 \right| < \varepsilon$$

for $|R - R_0| < \delta_1$ and $|\gamma - \gamma_0| < \delta_2$. Thus

$$\frac{1 - \varepsilon}{1 + \varepsilon} \leq \frac{C(R, \gamma)}{C(R_0, \gamma_0)} \leq (1 + \varepsilon)^2 .$$

For $\gamma_0 = \pi/2$ and $\gamma < \pi/2$:

$$0 < C(R, \gamma) < \frac{w(R, R, \gamma)}{K(R)} < \frac{\sqrt{2(1 - \sin \gamma)}}{K(R)} .$$

Require $|R - R_0| < R_0/2$. Clearly there is a $\delta > 0$ such that $0 < C(R, \gamma) < \varepsilon$ for $|\gamma - \pi/2| < \delta$. This shows continuity since $C(R_0, \pi/2) = 0$.

Part (ii). Let $R_1 < R_2$. By Lemma 11 $w(r, R_1, \gamma) < w(r, R_2, \gamma)$ for $r \geq R_2$. Dividing by $K(r)$ and taking $r \rightarrow \infty$ gives $C(R_1, \gamma) \leq C(R_2, \gamma)$. Likewise if $\gamma_1 < \gamma_2$, CP gives $w(r, R, \gamma_2) < w(r, R, \gamma_1)$ for $r > R$. Dividing by $K(r)$ and taking $r \rightarrow \infty$ gives $C(R, \gamma_2) \leq C(R, \gamma_1)$.

To complete the proof we need only show that if w_1 and w_2 are two solutions of equation (49a) with $w_1 \sim CK(r)$ as $r \rightarrow \infty$ and $w_2 \sim CK(r)$ as $r \rightarrow \infty$ then $w_1 \equiv w_2$. Let $u = w_1 - w_2$; without loss of generality we can assume $u \geq 0$, since by CP if $u(R) \geq 0$ then $u(r) \geq 0$ for $r \geq R$. Suppose $w_1 \not\equiv w_2$, then we must have $u(r) > 0$ for $r > R$. Note: $u(r) = o[K(r)]$ as $r \rightarrow \infty$. The equation satisfied by u is

$$(68) \quad (r^{n-1}u_r q)_r = r^{n-1}u$$

with

$$q(r) = \int_0^1 \{1 + [w_{2,r} + t(w_{2,r} - w_{1,r})]^2\}^{-3/2} dt.$$

By the corollary to Theorem 16:

$$q(r) = 1 + O[K^2(r)] \quad \text{and} \quad q_r(r) = O[K^2(r)]$$

as $r \rightarrow \infty$. Rewriting equation (68):

$$(69) \quad u_{rr} + u_r \left(\frac{n-1}{r} + \frac{q_r}{q} \right) = \frac{1}{q} u.$$

We construct a subsolution for equation (69). Let

$$U = \frac{Ae^{-r}}{r^{(n-1)/2}} \left(1 + \frac{a}{r} \right)$$

then

$$\begin{aligned} \Delta U &= U + \frac{Ae^{-r}}{r^{(n+3)/2}} \left\{ \frac{[8a - (n-1)(n-3)]}{4} - \frac{a(n+1)(n-5)}{4r} \right\} \\ U_r &= -\frac{Ae^{-r}}{r^{(n-1)/2}} \left[1 + \left(a + \frac{n-1}{2} \right) \frac{1}{r} + \frac{n+1}{2} \frac{a}{r^2} \right]. \end{aligned}$$

Choose a : $a > (n-1)(n-3)/4$. For R_1 sufficiently large

$$U_{rr} + U_r \left(\frac{n-1}{r} + \frac{q_r}{q} \right) > \frac{1}{q} U \quad \text{for } r \geq R_1$$

(independent of A because the equation is linear) and $1/q(r) > 0$ for $r \geq R_1$. Choose

$$A = (\exp R_1) R_1^{(n-1)/2} \left(1 + \frac{a}{R_1} \right)^{-1} u(R_1).$$

Let $V(r) = u(r) - U(r)$; V satisfies

$$V_{rr} + V_r \left(\frac{n-1}{r} + \frac{q_r}{q} \right) < \frac{1}{q} V \quad \text{for } r \geq R_1$$

and $V(R_1) = 0$. By the weak maximum principle [11] $\min_{R_1 \leq r \leq R_2} V(r) \geq \min [0, V(R_2)]$. Letting $R_2 \rightarrow \infty$ yields $V(r) \geq 0$ for $r \geq R_1$. Thus $u(r) \geq U(r)$ for $r \geq R_1$, but

$$u = o[K(r)] = o\left[\frac{e^{-r}}{r^{(n-1)/2}}\right]$$

a contradiction as $r \rightarrow \infty$.

Chapter III Solutions in Unsymmetrical Domains

10. General estimates, exponential decay. General estimates are given that apply to any solution of equation (1a):

THEOREM 18. *Suppose $u(x) \in C^2(\Omega)$ satisfies $Nu = u$ in Ω .*

(i) *If $B_R(y) \subset \Omega$ then $|u(x)| \leq v(|x - y|, R, 0)$ for $x \in B_R(y)$.*

(ii) *If $B_R^c(y) \subset \Omega$ then $|u(x)| \leq w(|x - y|, R, 0)$ for $x \in B_R^c(y)$.*

(iii) *If $H = \{x: \bar{n} \cdot x > b, |\bar{n}| = 1\} \subset \Omega$ then $|u(x)| \leq z(\bar{n} \cdot x - b, 0)$ for $x \in H$.*

Proof.

Part (i). This follows directly from CP (cf. the corollary to Theorem 1) since $\limsup (Tu - Tv) \cdot \nu \leq 0$ on $\partial B_R(y)$ implies $u(x) \leq v(x)$ in $B_R(y)$; for the same reason $-u(x) \leq v(x)$ in $B_R(y)$.

Part (ii). From Part (i)

$$(70) \quad |u(x)| \leq v(0, d, 0)$$

where $d = \text{dist}(x, \partial\Omega)$. By Theorem 10 $\lim_{d \rightarrow \infty} v(0, d, 0) = 0$; thus $|u(x)| = o(1)$ as $|x| \rightarrow \infty$. Also $w(|x - y|) = o(1)$ as $|x| \rightarrow \infty$, thus $w(|x - y|) - u(x) = o(1)$ as $|x| \rightarrow \infty$. On $\partial B_R^c(y)$, $\limsup (Tu - Tw) \cdot \nu \leq 0$; hence by CP, $u(x) \leq w(|x - y|)$ in $B_R^c(y)$. The same reasoning gives $-u(x) \leq w(|x - y|)$ in $B_R^c(y)$.

Part (iii). For $x \in H$, let $y = x + (R - \bar{n} \cdot x + b)\bar{n}$, then $B_R(y) \subset H$. By Part (i) $|u(x)| \leq v(|x - y|, R, 0) = v(R - \bar{n} \cdot x + b, R, 0)$. By Theorem 9 $\lim_{R \rightarrow \infty} v(R - \bar{n} \cdot x + b, R, 0) = z(\bar{n} \cdot x - b, 0)$. This gives Part (iii).

COROLLARY 1. *Let $d = \text{dist}(x, \partial\Omega)$, then*

$$|u(x)| \leq C \frac{d^{(n-1)/2}}{\exp d} \quad \text{for } d \geq d_0 > 0$$

where C depends on d_0 .

Proof. By inequality (70) $|u(x)| \leq v(0, d, 0)$; by Corollary 1 to Theorem 11 $v(0, d, 0) \sim H(d, 0)I(0)$ as $d \rightarrow \infty$, where

$$H(d, 0) = \sqrt{2\pi}C_1(0) \frac{d^{(n-1)/2}}{\exp d}.$$

Combining estimates gives the result.

COROLLARY 2. *If $B_{\tilde{r}}(0) \subset \Omega$ then*

$$|u(x)| \leq C \frac{\exp(-|x|)}{|x|^{(n-1)/2}} \quad \text{for } |x| \geq R$$

where C depends on R .

Proof. By Part (ii) $|u(x)| \leq w(|x|, R, 0)$; by Theorem 16 $w(|x|, R, 0) \sim C(R, 0)K(|x|)$ as $|x| \rightarrow \infty$. By § 4

$$K(|x|) \sim \sqrt{\pi/2} \frac{\exp(-|x|)}{|x|^{(n-1)/2}} \quad \text{as } |x| \rightarrow \infty.$$

Combining estimates gives the result.

COROLLARY 3. *If $H \subset \Omega$ then $|u(x)| \leq \sqrt{2} \exp(b - \bar{n} \cdot x)$ for $x \in H$.*

Proof. By Part (iii) $|u(x)| \leq z(\bar{n} \cdot x - b, 0)$ for $x \in H$. By Theorem 5 $z(\bar{n} \cdot x - b, 0) \leq \sqrt{2} \exp(b - \bar{n} \cdot x)$ for $x \in H$.

Corollaries 1, 2, and 3 give different exponential rates of decay for solutions to equation (1a) away from the boundary of the domain. The estimate of Corollary 3 is best possible because if $\Omega = H$ and $u(x) = z(\bar{n} \cdot x - b, 0)$ then $u = z(0, 0) = \sqrt{2}$ on ∂H . Gerhardt [8] gives an estimate of the same form with $\sqrt{2}$ replaced by $(n + 1)$.

COROLLARY 4. *If Ω is an exterior domain then problem (1) has a unique solution.*

Proof. Two solutions u_1 and u_2 to problem (1) must satisfy $u_1(x) = o(1)$ and $u_2(x) = o(1)$ as $|x| \rightarrow \infty$; thus $u_1(x) - u_2(x) = o(1)$ as $|x| \rightarrow \infty$. CP gives that $u_1 \equiv u_2$.

The derivatives of any solution to equation (1a) can be estimated as follows:

LEMMA 15. *Suppose $u(x) \in C^2(\Omega)$ and $Nu = u$ in Ω . If $B_{4\delta}(x_0) \subset \Omega$ then $|u|_{2,\alpha,B_\delta} < C|u|_{0,B_{4\delta}}$ where C depends only on δ and n .*

Proof. See Step 3 of the proof of Theorem 4.

Lemma 15 in conjunction with Corollaries 1, 2, and 3 shows that the derivatives of a solution to equation (1a) decay at the same rate at which the solution decays:

COROLLARY.

$$|\nabla u(x)| < C \frac{d^{(n-1)/2}}{\exp d} \quad \text{for } d \geq d_0, \quad d = \text{dist}(x, \partial\Omega)$$

where C depends on d_0 .

Proof. Combining Corollary 1 and Lemma 15 (with $8\delta = d_0$) gives

$$|\nabla u(x)| < \bar{C} \frac{(d + d_0/2)^{(n-1)/2}}{\exp(d - d_0/2)}$$

for $d \geq d_0$, where \bar{C} depends on d_0 . Thus the corollary holds with a different constant C .

REMARK. Similar estimates hold for higher derivatives, and if $B_R^c \subset \Omega$ or $H \subset \Omega$, the analogous estimates are based upon Corollaries 2 and 3. These estimates improve upon the derivative estimate given by Gerhardt [8].

11. **Special estimates.** We now put restrictions on $\gamma(\sigma)$ and Ω to obtain special estimates on solutions to problem (1). For simplicity we assume that Σ is piecewise smooth.

We first extend CPS of Chapter I.

THEOREM 19. *Suppose $Nu = u$ in Ω and $Tu \cdot \nu = \cos \gamma$ on Σ^* , with $0 \leq \gamma(\sigma) \leq \gamma_0 \leq \pi/2$. If $\Omega \subset B_R^c(y)$ and $u \geq o(1)$ as $|x| \rightarrow \infty$, then $w(r, R, \gamma_0) \leq u(x)$ where $r = |x - y|$.*

Proof. By the proof of Theorem 2, $Tw \cdot \nu \leq \cos \gamma_0 \leq Tu \cdot \nu$ on Σ^* . By hypothesis $w - u \leq o(1)$ as $|x| \rightarrow \infty$. Thus CP gives the conclusion.

COROLLARY. *If Ω is an exterior domain, $0 \in \Omega^c$, and u is a solution to problem (1) with $0 \leq \gamma(\sigma) \leq \gamma_0 < \pi/2$, then*

$$C_1 \frac{e^{-|x|}}{|x|^{(n-1)/2}} < u(x) < C_2 \frac{e^{-|x|}}{|x|^{(n-1)/2}} \quad \text{for } x \in \Omega$$

where C_1 and C_2 are positive constants.

Proof. For some positive R_1 and R_2 , $B_{R_1}^e(0) \subset \Omega \subset B_{R_2}^e(0)$. By Theorem 18 and Corollary 2 to Theorem 18

$$u(x) \leq w(|x|, R_1, 0) \leq C \frac{\exp(-|x|)}{|x|^{(n-1)/2}}$$

and $u(x) = o(1)$ as $|x| \rightarrow \infty$. Thus by Theorem 19: $w(|x|, R_2, \gamma_0) \leq u(x)$. By Theorem 16: $w(|x|, R_2, \gamma_0) \sim C(R_2, \gamma_0)K(|x|)$. Combining estimates gives the result.

REMARK. In this case we have determined the rate of decay.

We now improve upon the upper bounds given in § 10.

THEOREM 20. *Let u be a solution to problem (1) with $\gamma_1 \leq \gamma(\sigma) \leq \pi$, $0 < \gamma_1 < \pi/2$. Suppose $B_R(y) \subset \Omega$ and $B_{\bar{R}}(y) \cap \Omega$ is piecewise smooth and convex, where $B_{\bar{R}}(y)$ is the maximal domain of existence of $v(r, R, \gamma_1)$, $r = |x - y|$, then $u(x) \leq v(r, R, \gamma_1)$ in $B_R(y)$.*

Proof. Let $\partial(B_{\bar{R}} \cap \Omega) = \Sigma^0 + \Sigma^1 + \Sigma^2$ where Σ^1 and $\Sigma^2 \in C^1$, $\Sigma^1 \subset \Sigma = \partial\Omega$ and $\Sigma^2 \subset (\partial B_{\bar{R}}) \cap \Omega$.

$$\text{For } x \in \Sigma^2, \quad Tv \cdot \nu = 1 > Tu \cdot \nu.$$

$$\text{For } x \in \Sigma^1, \quad Tv \cdot \nu = (\cos \theta) \sin \psi(r)$$

where θ is the angle between ν and $x - y$. Let π_x be the tangent plane to Σ at x and let $r_1 = \text{dist}(y, \pi_x)$. We have $\cos \theta = r_1/r$ and $r \geq r_1 \geq R$. Thus

$$Tv \cdot \nu = r_1 \frac{\sin \psi(r)}{r} \geq \sin \psi(r_1) \geq \sin \psi(R) = \cos \gamma_1 \geq Tu \cdot \nu.$$

The first inequality comes from Lemma 6(ii).

Therefore CP applies, giving $u \leq v$ in $B_{\bar{R}} \cap \Omega$.

COROLLARY 1. *Let u be solution to problem (1) with $0 \leq \gamma(\sigma) \leq \pi/2$. Suppose $B_R(y) \subset \Omega$ and $B_{R_i}(y) \cap \Omega$ is piecewise smooth and convex, with $\lim_{i \rightarrow \infty} R_i = \infty$, then $u(x) \geq 0$ in $B_R(y)$.*

Proof. Define γ_i so that $v(r, R_i, \pi) = v(r, R, \gamma_i)$ for $0 \leq r \leq R$. Note: $\pi/2 < \gamma_i < \pi$. By Corollary 1 to Theorem 11 $\lim_{i \rightarrow \infty} v(r, R_i, \pi) = 0$

for $0 \leq r \leq R$. By § 3, $-v(r, R, \gamma_i) = v(r, R, \pi - \gamma_i)$ and $-u$ is a solution to problem (1) with boundary data $\pi - \gamma(\sigma)$. Since $0 < \pi - \gamma_i < \pi/2 \leq \pi - \gamma(\sigma)$, Theorem 20 gives $-u \leq -v$. Thus $v(r, R, \gamma_i) \leq u(x)$ in $B_R(y)$. Letting $i \rightarrow \infty$ gives the result.

COROLLARY 2. *Let $H = \{x: x_1 \geq 0\}$. There is a unique solution to $Nu = u$ in H and $Tu \cdot \nu = \cos \gamma$ on ∂H , γ constant: the one-dimensional solution, $u = z(x_1, \gamma)$.*

Proof. We need only consider $0 \leq \gamma \leq \pi/2$. By Corollary 1, $u(x) \geq 0$ on H . By Theorem 19, if $H \subset B_R^+(y)$ then $w(r, R, \gamma) \leq u(x)$. For $x \in H$, choose $y = (-R, x_2, \dots, x_n)$, then $r = R + x_1$. Thus $w(R + x_1, R, \gamma) \leq u(x)$. Take the limit $R \rightarrow \infty$: $z(x_1, \gamma) \leq u(x)$.

By Theorem 20, if $B_R(y) \subset H$ then $u(x) \leq v(r, R, \gamma)$. For $x \in H$ choose $y = (R, x_2, \dots, x_n)$, then $r = R - x_1$. Thus $u(x) \leq v(R - x_1, R, \gamma)$. Take the limit $R \rightarrow \infty$: $u(x) \leq z(x_1, \gamma)$. Therefore $u(x) = z(x_1, \gamma)$.

REMARK. Uniqueness also holds for $n = 1$, since a solution to problem (1) in H is a solution to problem (1) in H for $n = 2$.

For an interior domain, we define the “interior rolling number”:

$$\mathcal{R}_1 = \max \{R: \text{for each } x \in \Sigma \text{ there is a ball } B_R \subset \Omega, \text{ with } x \in \bar{B}_R\}.$$

THEOREM 21. *Let Ω be bounded and $\Sigma \in C^1$. Let $u \in C^1(\bar{\Omega}) \cap C^2(\Omega)$ be a solution to problem (1), with $\gamma_0 \leq \gamma(\sigma) < \pi$, $0 < \gamma_0 \leq \pi/2$. Suppose \mathcal{R}_1 exists, then*

- (i) $\max u(x) \leq v(\mathcal{R}_1, \mathcal{R}_1, \gamma_0)$.
- (ii) if $B_R(y) \subset \Omega$ and $R \leq \mathcal{R}_1$, then $u(x) \leq v(r, R, \gamma_0)$ in $B_R(y)$, with $r = |x - y|$.

Proof.

Part (i). If $u \leq 0$ there is nothing to prove. If $u(x_1) > 0$ for $x_1 \in \Omega$, then by the maximum principle, $\max u(x)$ occurs on Σ , say at x_0 . Suppose $u(x_0) > v(\mathcal{R}_1, \mathcal{R}_1, \gamma_0)$. By Theorems 6 and 9, $v(R, R, \gamma_0)$ is continuous and strictly decreasing in R . By Theorem 7, $\lim_{R \rightarrow 0} v(R, R, \gamma_0) = \infty$. We can thus choose $R < \mathcal{R}_1$ so that $v(R, R, \gamma_0) = u(x_0)$. There is a ball $B_R \subset \Omega$ with $x_0 \in \bar{B}_R$. By CP: $u(x) < v(r, R, \gamma_0)$ in B_R . Thus $(\partial u / \partial \nu)(x_0) \geq (\partial v / \partial \nu)(x_0)$. However, $(\partial v / \partial \nu)(x_0) = \cot \gamma_0$ and $(\partial u / \partial \nu)(x_0) = \cot \gamma(x_0)$, since at x_0 :

$$\cot \gamma(x_0) = \frac{\nabla u \cdot \nu}{\sqrt{1 + |\nabla u|^2}} = \frac{|\nabla u|}{\sqrt{1 + |\nabla u|^2}}$$

and $(\partial u/\partial \nu)(x_0) = |\nabla u|$. Therefore $(\partial u/\partial \nu)(x_0) = (\partial v/\partial \nu)(x_0)$, contradicting Lemma 1.

Part (ii). This follows immediately from Part (i) and CP.

REMARK. The proof shows that equality holds in Part (i) only if Ω is a ball and $\gamma(\sigma) \equiv \gamma_0$.

For an exterior domain, we define the “exterior rolling number”:

$$\mathcal{R}_2 = \min \{R: \text{for each } x \in \Sigma \text{ there is a ball } B_R \supset \Omega^c, \text{ with } x \in \bar{B}_R\}.$$

THEOREM 22. *Let Ω be an exterior domain, $\Sigma \in C^1$ and Ω^c convex. Let $u(x) \in C^1(\bar{\Omega}) \cap C^2(\Omega)$ be a solution to problem (1) with $\gamma_0 \leq \gamma(\sigma) < \pi$, $0 < \gamma_0 \leq \pi/2$. Then*

(i) $\max u(x) < z(0, \gamma_0)$.

(ii) if \mathcal{R}_2 exists, then $\max u(x) \leq w(\mathcal{R}_2, \mathcal{R}_2, \gamma_0)$.

(iii) if \mathcal{R}_2 exists and $B_R^c(y) \subset \Omega$, $R \geq \mathcal{R}_2$, then $u(x) \leq w(r, R, \gamma_0)$ in $B_R^c(y)$, with $r = |x - y|$.

Proof.

Part (i). If $u \leq 0$ there is nothing to prove. If $u(x_1) \geq 0$ for $x_1 \in \Omega$, then by the maximum principle and the fact that $u(x) = o(1)$ as $|x| \rightarrow \infty$: $\max u(x)$ occurs on Σ , say at x_0 . Suppose $u(x_0) \geq z(0, \gamma_0) = \sqrt{2}(1 - \sin \gamma_0)$. Choose $\gamma_1 \leq \gamma_0$ so that $u(x_0) = z(0, \gamma_1)$. For convenience suppose $x_0 = 0$, that the tangent plane to Σ at x_0 is given by $\{x: x_1 = 0\}$, and that $H = \{x: x_1 > 0\} \subset \Omega$. Since $u - z \leq o(1)$ as $|x| \rightarrow \infty$, by CP: $u(x) < z(x, \gamma_1)$ in H . Hence $(\partial u/\partial \nu)(0) \geq (\partial z/\partial \nu)(0)$. However, $(\partial z/\partial \nu)(0) = \cot \gamma_1$ and $(\partial u/\partial \nu)(0) = \cot \gamma(0)$; thus $(\partial u/\partial \nu)(0) = (\partial v/\partial \nu)(0)$, contradicting Lemma 1.

Part (ii). Suppose $u(x_0) = \max u(x) > w(\mathcal{R}_2, \mathcal{R}_2, \gamma_0)$, where $x_0 \in \Sigma$. By Theorems 13 and 15: $w(R, R, \gamma_0)$ is continuous and strictly increasing in R , with $\lim_{R \rightarrow \infty} w(R, R, \gamma_0) = z(0, \gamma_0)$. We can choose $R > \mathcal{R}_2$ so that $w(R, R, \gamma_0) = u(x_0)$, since by Part (i) $u(x_0) < z(0, \gamma_0)$. There is a ball B_R , with $\Omega^c \subset B_R$ and $x_0 \in \bar{B}_R$. By CP, $u(x) < w(r, R, \gamma_0)$ in B_R^c , thus $(\partial u/\partial \nu)(x_0) \geq (\partial w/\partial \nu)(x_0)$. However, $(\partial w/\partial \nu)(x_0) = \cot \gamma_0$ and $(\partial u/\partial \nu)(x_0) = \cot \gamma(x_0)$. Therefore $(\partial w/\partial \nu)(x_0) = (\partial u/\partial \nu)(x_0)$, contradicting Lemma 1.

Part (iii). This follows immediately from CP and Part (ii).

REMARKS. (1) The proof shows that equality can hold in Part (ii) only if Ω^c is a ball and $\gamma(\sigma) \equiv \gamma_0$.

(2) If $\Sigma \in C^2$ and Ω° is strictly convex, then \mathcal{R}_2 will exist, $\mathcal{R}_2 = (\bar{k})^{-1}$ where $\bar{k} = \min k(\sigma)$ over Σ and $k =$ the minimum principal curvature of Σ at σ (see [2]).

12. Two examples. We give two examples that show the appropriateness of the convexity condition of Theorems 20 and 22 and the “rolling number” condition of Theorem 21.

EXAMPLE 1. We construct a domain $\Omega = B_R(y) \cup \mathcal{H}$ where \mathcal{H} is part of a cone $\mathcal{H} = \{x: x_1 > |x| \cos \alpha, x_1 < \bar{x}_1\}$, $0 < \alpha < \pi/2$, and $y = (R + \varepsilon, 0, \dots, 0)$; here $\varepsilon > 0$ and

$$(71) \quad \bar{x}_1 = \varepsilon \cdot \frac{2R + \varepsilon}{R + \varepsilon} \left\{ 1 + \left[1 - \varepsilon \sec^2 \alpha \frac{2R + \varepsilon}{(R + \varepsilon)^2} \right]^{1/2} \right\}^{-1}$$

where \bar{x}_1 is the x_1 coordinate of the first intersection of \mathcal{H} with B_R (see Figure 1).




FIGURE 1

Let $v = v(|x - y|, R, \gamma)$ and let $u(x)$ be the solution to problem (1) in Ω , where $\gamma = \pi/2 - \alpha$. For ε sufficiently small we will show that $u \not\leq v$ in B_R .

Let w_n be the volume of the unit ball in n -dimensions.

We modify an argument of Finn [5]. By integration by parts:

$$\int_{\Omega} u dx = \int_{\partial\Omega} Tu \cdot \nu d\sigma = (\cos \gamma) |\partial\Omega|$$

$$\int_{B_R} v dx = (\cos \gamma) |\partial B_R| .$$

Thus

$$\int_{B_R} (u - v) dx = \cos \gamma (|\partial\Omega| - |\partial B_R|) - \int_{\Omega - B_R} u dx .$$

Now $|\partial\Omega| - |\partial B_R| = |\Sigma^1| - |\Sigma^2|$ where $\Sigma^1 \subset \partial\mathcal{H}$ and $\Sigma^2 \subset \partial B_R$. We calculate:

$$|\Sigma^1| = w_{n-1} (\tan^{n-2} \alpha) \sec \alpha \bar{x}_1^{n-1}$$

$$|\Sigma^2| = (n - 1) w_{n-1} \int_0^a r_1^{n-2} ds \leq w_{n-1} a^{n-1}$$

[where r_1 is the distance from the x_1 axis along a profile curve and s is arc length, thus $r_1(s) \leq s$]

$$a = R \sin^{-1} \left(\frac{\bar{x}_1 \tan \alpha}{R} \right) = \bar{x}_1 \tan \alpha + O(\bar{x}_1^3)$$

$$|\Omega - B_R| \leq |\mathcal{H}| = w_{n-1} \frac{\tan^{n-1} \alpha}{n} x_1^n.$$

Let $0 < R_1 < R$ and $y_1 = (R_1, 0, \dots, 0)$.

Claim. For ε sufficiently small $\Omega - B_R \subset B_{R_1}(y_1)$ and $u(x) < h(x)$ in $B_{R_1} \cap \Omega$, where

$$h(x) = \frac{n}{R_1} + R_1 + \sqrt{R_1^2 - |x - y_1|^2}.$$

Assuming the claim:

$$\int_{B_R} (u - v) dx \geq \cos \gamma w_{n-1} \tan^{n-2} \alpha (\sec \alpha - \tan \alpha) \bar{x}_1^{n-1}$$

$$- \left(\frac{n}{R_1} + R_1 \right) \frac{w_{n-1}}{n} \tan^{n-1} \alpha \bar{x}_1^n$$

$$+ O(\bar{x}_1^{n+2}).$$

For \bar{x}_1 sufficiently small

$$\int_{B_R} (u - v) dx > 0$$

by equation (71) this is true if ε is sufficiently small. Therefore $u \not\leq v$ in B_R .

Proof of claim. Since $Nh \leq h$ (cf. Corollary 1 to Theorem 1) we need show only that $Th \cdot \nu \geq \cos \gamma$ on $[\partial(\Omega \cap B_{R_1})]^* = \Sigma^1 + \Sigma^3 + \Sigma^4$, where $\Sigma^3 \subset \partial B_{R_1}$ and $\Sigma^4 \subset \partial B_R$. On Σ^1 , $Th \cdot \nu = \cos \gamma$ and on Σ^3 , $Th \cdot \nu = 1$. For $x \in \Sigma^4$ we must show

$$(72) \quad \cos \theta \geq \frac{R_1}{r} \cos \gamma$$

where $r = |x - y_1|$ and θ is the angle between $x - y_1$ and ν .

By changing the x_1 coordinate, $x \rightarrow R_1 - x_1$, $-(R_1, 0, \dots, 0)$, we calculate

$$(73) \quad \cos \theta = \frac{1}{R} \left[r + \frac{x_1}{r} (R - R_1 + \varepsilon) \right].$$

We check that $\cos \theta$ is smallest for $x_1 = \hat{x}_1$ determined by $\partial B_R \cap \partial B_{R_1}$:

$$\hat{x}_1 = \frac{(R - R_1)(R_1 - \varepsilon) - \varepsilon^2/2}{R - R_1 + \varepsilon}.$$

Putting this value into equation (73) gives

$$(74) \quad \cos \theta \geq 1 - \varepsilon \frac{(R - R_1)}{RR_1} - \frac{\varepsilon^2}{2RR_1}.$$

For ε sufficiently small, so that

$$1 - \varepsilon \frac{(R - R_1)}{RR_1} - \frac{\varepsilon^2}{2RR_1} > \frac{R_1 \cos \gamma}{R_1 - \varepsilon}$$

then condition (72) will be satisfied.

REMARKS. (1) The domain Ω is not convex.

(2) The example can be modified so that Σ is smooth $B_R \subset \Omega$, $R > R_1$, and $u(x) \prec v(r, R, \gamma)$ in B_R .

EXAMPLE 2. We construct a domain $\Omega = B_R^c(y) \cup \mathcal{K}$ (see Figure 2) by the method of Example 1 so that $u(x) \prec w(r, R, \gamma)$ in $B_R^c(y)$.




FIGURE 2

REMARKS. (1) Ω^c is not convex.

(2) The example can be modified so that Σ is smooth, Ω^c is not convex, $B_R^c \subset \Omega$, and $u(x) \prec w(r, R, \gamma)$ in B_R^c .

13. The infinite wedge. We consider solutions to problem (1) in

$$\mathcal{K}_\alpha = \{(x_1, x_2): x_1 > |x_2| \cos \alpha\}$$

with $0 < \alpha < \pi$, and γ constant, $0 < \gamma \leq \pi/2$.

THEOREM 23. For $0 < \gamma < \pi/2$ problem (1) has a unique positive solution in \mathcal{K}_α . Furthermore, for $\alpha < \pi/2$: on Σ^* , $u(x) \geq z(0, \gamma)$ and $\lim_{|x| \rightarrow \infty} u(x) = z(0, \gamma)$. For $\alpha > \pi/2$: on Σ^* , $u(x) \leq z(0, \gamma)$ and $\lim_{|x| \rightarrow \infty} u(x) = z(0, \gamma)$. Here $\Sigma^* = \partial \mathcal{K}_\alpha - \{(0, 0)\}$.

Proof. Existence comes from Theorem 4. Let $d = \text{dist}(x, \Sigma)$.

Case $\alpha < \pi/2$. Any solution $u(x)$ to problem (1) must be non-negative since \mathcal{K}_α is convex, by Corollary 1 to Theorem 20. By

Theorem 19, if $\Omega \subset B_R^c$ then $w(r, R, \gamma) < u(x)$ in B_R^c . By a limit argument (cf. the proof of Corollary 2 to Theorem 20) we obtain

$$(75) \quad z(d, \gamma) \leq u(x) \quad \text{for } x \in \mathcal{H}_\alpha.$$

If $B_R \subset \mathcal{H}_\alpha$, then $u(x) < v(r, R, \gamma)$ in B_R , by Theorem 20. Given $x \in \mathcal{H}_\alpha$ we choose $B_R \subset \mathcal{H}_\alpha$ with center on the x_1 axis so that the radius through x is perpendicular to Σ . This determines $R = x_1 \tan \alpha \sec \alpha - d \tan^2 \alpha$. Thus,

$$(76) \quad u(x) \leq v(R - d, R, \gamma).$$

Suppose there are two solutions $u_1(x)$ and $u_2(x)$ to problem (1). We show that $u_1 - u_2 = o(1)$ as $|x| \rightarrow \infty$.

Given $\varepsilon > 0$, choose d_1 so that $v(0, d_1, 0) < \varepsilon/2$. For $d > d_1$: $|u_1(x) - u_2(x)| < |u_1(x)| + |u_2(x)| < \varepsilon$. For $x_1 > \bar{x}_1$ and $d \leq d_1$, with \bar{x}_1 sufficiently large: $0 < v(R - d, R, \gamma) - z(d, \gamma) < \varepsilon$, by Theorem 9. Thus, $|u_1(x) - u_2(x)| < \varepsilon$ for $x_1 > \bar{x}_1$, by estimates (75) and (76). Therefore $u_1 - u_2 = o(1)$ as $|x| \rightarrow \infty$. By CP, $u_1 \equiv u_2$.

Taking $d = 0$ in estimates (75) and (76) gives the limit statement and estimate on Σ^* .

Case $\alpha > \pi/2$. We first show that for any solution $u(x)$ to problem (1): $u \geq o(1)$ as $|x| \rightarrow \infty$.

Given $\varepsilon > 0$, choose d_1 as above. For $d > d_1$: $u(x) > -\varepsilon/2$. Choose $\gamma_1 > \pi/2$ so that $v(r, d_1, \gamma_1) > -\varepsilon$ for $0 \leq r \leq d_1$. For $d \leq d_1$ and $x_1 < \bar{x}_1$, \bar{x}_1 sufficiently small so that $x \in B_{d_1}(y) \subset \mathcal{H}_\alpha$, with $\text{dist}(y, \Sigma) = d$, and $B_{\bar{x}_1}(y) \cap \mathcal{H}_\alpha$ is convex, we have: $u(x) \geq v(r, d_1, \gamma) > -\varepsilon$ (cf. the proof of Corollary 1 to Theorem 20). Therefore $u \geq o(1)$ as $|x| \rightarrow \infty$.

Now, if $\mathcal{H}_\alpha \subset B_R^c$ then $w(r, R, \gamma) \leq u(x)$ in B_R^c , by Theorem 19. This shows $u(x) > 0$ in \mathcal{H}_α . Furthermore, given $x \in \mathcal{H}_\alpha$ with $x_1 < 0$, we choose $B_{R_x}(y) \subset \mathcal{H}_\alpha^c$, with center on the x_1 axis so that segment joining x and y is perpendicular to Σ . This determines $R_x = -x_1 \sec \alpha \tan \alpha + d \tan^2 \alpha$. We have

$$(77) \quad w(R_x + d, R_x, \gamma) \leq u(x).$$

Given $\varepsilon > 0$, choose d_1 as above. Thus,

$$(78) \quad 0 < u(x) < \varepsilon/2 \quad \text{for } d > d_1.$$

Choose \bar{x}_1 so that $|w(R_x + d, R_x, \gamma) - z(d, \gamma)| < \varepsilon/2$ for $x_1 < \bar{x}_1$ and $0 \leq d \leq d_1$. Choose R so that $|v(R - d, R, \gamma) - z(d, \gamma)| < \varepsilon/2$ for $d \leq d_1$. For $x \in \mathcal{H}_\alpha$ with $d \leq d_1$, there is a ball $B_R(y) \subset \mathcal{H}_\alpha$, $\text{dist}(y, \Sigma) = R$, the radius through x is perpendicular to Σ , and $B_{\bar{R}}(y) \cap \mathcal{H}_\alpha$ convex, if $x_1 < \hat{x}_1$, \hat{x}_1 sufficiently small. Thus

$$(79) \quad u(x) \leq v(R - d, R, \gamma) \quad \text{for } d \leq d_1 \text{ and } x_1 < \hat{x}_1.$$

Suppose there are two solutions $u_1(x)$ and $u_2(x)$ to problem (1). They must both satisfy estimates (77), (78), and (79). Thus $|u_1(x) - u_2(x)| < \varepsilon$ for $x_1 < \min(\bar{x}_1, \hat{x}_1)$. This implies $u_1 - u_2 = o(1)$ as $|x| \rightarrow \infty$; by CP, $u_1 \equiv u_2$.

The limit statements come from estimates (77) and (79) with $d = 0$.

We now obtain the estimate on Σ^* . Let $\Omega = \mathcal{H}_\alpha \cap \{x: x_2 < 0\}$. Let $B_R(y) \subset \mathcal{H}_\alpha$ be tangent to Σ^* at a point $x_0 \in \partial\Omega$. Let $v = (r, R, \gamma)$ and $[\partial B_{\bar{R}}(y)]^* = \Sigma^1 + \Sigma^2 + \Sigma^3$, where $\Sigma^1 \subset \Sigma^*$, $\Sigma^2 \subset \{x: x_1 > 0, x_2 = 0\}$ and $\Sigma^3 \subset \partial B_{\bar{R}}(y)$. On Σ^1 , $Tv \cdot \nu \geq \cos \gamma$, by the proof of Theorem 20. On Σ^2 , $Tv \cdot \nu > 0$. On Σ^3 , $Tv \cdot \nu = 1$. Because of uniqueness, $u(x_1, x_2) = u(x_1, -x_2)$. Thus $\partial u / \partial x_2 = 0$ on Σ^2 , hence $Tu \cdot \nu = 0$ on Σ^2 . Therefore $Tv \cdot \nu \geq Tu \cdot \nu$ on $[\partial B_{\bar{R}}(y)]^*$; by CP, $u(x) < v(r, R, \gamma)$ in $B_R(y)$. Taking $R \rightarrow \infty$, with $B_R(y)$ tangent at x_0 , gives:

$$(80) \quad \begin{aligned} u(x) &< z(\bar{n}_1 \cdot x, \gamma) && \text{in } H_1 = \{x: \bar{n}_1 \cdot x > 0\} \\ u(x) &< z(\bar{n}_2 \cdot x, \gamma) && \text{in } H_2 = \{x: \bar{n}_2 \cdot x > 0\} \end{aligned}$$

where $\bar{n}_1 = (\sin \alpha, -\cos \alpha)$ and $\bar{n}_2 = (\sin \alpha, \cos \alpha)$. Thus on Σ^* , $u(x) \leq z(0, \gamma)$.

We now study the behavior at the vertex.

THEOREM 24. *For $\alpha + \gamma \geq \pi/2$ and $0 < \alpha < \pi/2$, the solution to problem (1) in \mathcal{H}_α satisfies*

$$(i) \quad \liminf_{x \rightarrow 0} u(x) \geq \sqrt{2}(1 - \sqrt{1 - k^2})^{1/2}$$

$$(ii) \quad \limsup_{x \rightarrow 0} u(x) \leq 2\sqrt{2}(1 - \sqrt{1 - k^2})^{1/2}$$

where $k = \cos \gamma / \sin \alpha$.

Proof.

Part (i). As a comparison surface choose $z(x_1, \hat{\gamma})$, with $\hat{\gamma} = \cos^{-1} k$. On $(\partial \mathcal{H}_\alpha)^*$, $Tz \cdot \nu = -\sin \psi(x_1) \sin \alpha \leq \cos \hat{\gamma} \sin \alpha = \cos \gamma = Tu \cdot \nu$. Since $u > 0$, then $z - u \leq o(1)$ as $|x| \rightarrow \infty$, in \mathcal{H}_α . Hence by CP, $z < u$ in \mathcal{H}_α . Therefore $\liminf_{x \rightarrow 0} u \geq z(0, \hat{\gamma})$. This gives the result.

Part (ii). As a comparison surface choose

$$h(x) = \frac{2}{R} + R - \sqrt{R^2 - |x - y|^2}$$

where $y = (kR, 0)$. We have $Nh \leq h$ in $B_R(y)$. We check: $Th \cdot \nu = \cos \gamma$ on $(\partial \mathcal{H}_\alpha)^*$ and $Th \cdot \nu = 1$ on ∂B_R . Thus, $Th \cdot \nu \geq Tu \cdot \nu$ on $[\partial(B_R \cap \mathcal{H}_\alpha)]^*$; by CP, $u(x) < h(x)$ in $B_R \cap \mathcal{H}_\alpha$. Therefore,

$$\limsup_{x \rightarrow 0} u(x) \leq h(0) = \frac{2}{R} + R(1 - \sqrt{1 - k^2}) .$$

Minimize this bound by choosing

$$R = \sqrt{2}(1 - \sqrt{1 - k^2})^{-1/2} .$$

This gives the result.

REMARKS. In n -dimensions, with $\mathcal{H}_\alpha = \{x: x_1 > |x| \cos \alpha\}$, $0 < \alpha < \pi/2$, Theorems 23 and 24 still hold with $2\sqrt{2}$ replaced by $2\sqrt{n}$ in Part (ii) of Theorem 24. We note

$$\sqrt{2}(1 - \sqrt{1 - k^2})^{1/2} > z(0, \gamma) .$$

This shows that there is a "rise" at the vertex. For the case $\alpha + \gamma < \pi/2$, Concus and Finn [3] show that the solution to problem (1) is unbounded at the vertex.

For

$$\mathcal{H}_\alpha = \{x: x_1 > \sqrt{x_1^2 + x_2^2} \cos \alpha\} , \quad 0 < \alpha < \pi .$$

Theorems 23 and 24 still hold, with the same modification.

Acknowledgment. This paper is based on the author's Ph.D. Dissertation written under the direction of Professor R. Finn at Stanford University.

REFERENCES

1. G. Bakker, *Handbook der Experimental Physik*, Vol. 6, Akademische Verlagsgesellschaft M.B.H., Leipzig, 1928.
2. W. Blaschke, *Kreis und Kugel*, Walter de Gruyter and Co., Berlin, 1956.
3. P. Concus and R. Finn, *On capillary free surfaces in a gravitational field*, Acta Math., **132** (1974), 207-223.
4. R. Finn, *Capillarity phenomena*, Uspehi Mat. Nauk., **29** (1974), 131-152.
5. ———, *Some comparison properties and bounds for capillary surfaces*, (to appear).
6. R. Finn and C. Gerhardt, *The internal sphere condition and the capillarity problem*, Ann. Mat. Pura Appl. IV., **112** (1977), 13-31.
7. C. Gerhardt, *On the exterior capillary problem*, Math. Zeit., **136** (1974), 345-349.
8. ———, *A further note on the exterior capillary problem*, Math. Zeit., **150** (1976), 149-157.
9. ———, *Global regularity of the solutions to the capillary problem*, Ann. Scuola Norm. Sup. Pisa IV, **3** (1976), 157-175.
10. ———, *A note on the exterior capillary problem*, Math. Zeit., **142** (1975), 203-204.
11. D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, New York, 1977.
12. W. E. Johnson and L. M. Perko, *Interior and exterior boundary value problems from the theory of the capillary tube*, Arch. Rat. Mech. Anal., **29** (1968), 125-143.
13. D. A. Ladyzhenskaya and N. N. Ural'tseva, *Linear and Quasilinear Elliptic Equations*, Academic Press, New York, 1966.

14. N. N. Lebedev, *Special Functions and Their Applications*, Dover, New York, 1972.
15. L. M. Perko, *Boundary layer analysis of the wide capillary tube*, Arch. Rat. Mech., **45** (1972), 120-133.
16. L. Simon and J. Spruck, *Existence and regularity of a capillary surface with prescribed contact angle*, Arch. Rat. Mech. Anal., **61** (1976), 19-34.

Received September 28, 1979.

UNIVERSITY OF SOUTHERN CALIFORNIA
LOS ANGELES, CA 90007

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$84.00 a year (6 Vols., 12 issues). Special rate: \$42.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1980 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Reinhold Böhme, Stefan Hildebrandt and Engelbert Tausch, <i>The two-dimensional analogue of the catenary</i>	247
Jean Ellen Taylor, <i>Nonexistence of F-minimizing embedded disks</i>	279
Claus Gerhardt, <i>A free boundary value problem for capillary surfaces</i>	285
Enrico Giusti, <i>Generalized solutions for the mean curvature equation</i>	297
Jin-Tzu Chen, <i>On the existence of capillary free surfaces in the absence of gravity</i>	323
Leon M. Simon, <i>Regularity of capillary surfaces over domains with corners</i>	363
Nicholas Jacob Korevaar, <i>On the behavior of a capillary surface at a re-entrant corner</i>	379
Henry Wente, <i>The symmetry of sessile and pendent drops</i>	387
E. Gonzalez, Umberto Massari and I. Tamanini, <i>Existence and regularity for the problem of a pendent liquid drop</i>	399
Henry Wente, <i>The stability of the axially symmetric pendent drop</i>	421
David Siegel, <i>Height estimates for capillary surfaces</i>	471
Bruce Edward Turkington, <i>Height estimates for exterior problems of capillarity type</i>	517
Robert Finn, <i>The sessile liquid drop. I. Symmetric case</i>	541