

# Pacific Journal of Mathematics

**IMBEDDING SMOOTH INVOLUTIONS IN TRIVIAL BUNDLES**

KENNETH JOSEPH PREVOT

## IMBEDDING SMOOTH INVOLUTIONS IN TRIVIAL BUNDLES

K. J. PREVOT

Let  $\tilde{M}^n$  denote a closed manifold with smooth involution, and  $\tilde{M}^n \rightarrow M^n$  the corresponding principal  $Z_2$ -bundle  $\xi$  classified by  $w_1(\xi)$  in  $H^1(M^n; Z_2)$ . The existence of a nested sequence  $\tilde{N}^{n-k+1} \subset \tilde{N}^{n-k+2} \subset \dots \subset \tilde{N}^{n-1} \subset \tilde{N}^n = \tilde{M}^n$  of characteristic submanifolds (corresponding to principal  $Z_2$ -bundles  $\xi_{n-k+1} \subset \xi_{n-k+2} \subset \dots \subset \xi_{n-1} \subset \xi_n = \xi$ ) satisfying  $(w_1(\xi_i))^{n-k+i} = 0$ , for  $1 \leq k \leq n$  and all  $i$  with  $n-k+1 \leq i \leq n$ , provides a necessary and sufficient condition for imbedding  $\xi$  in vector bundle  $M^n \times R^k \rightarrow M^n$  (preserving fibers).

For closed  $n$ -manifolds, the condition  $(w_1(\xi))^k = 0$  allows  $\xi$  to be imbedded in the vector bundle  $M^n \times R^k \rightarrow M^n$  "up to free  $Z_2$ -cobordism."

1. Introduction. In the following, all manifolds will be smooth and compact, and maps are understood to be smooth where appropriate. Let  $\pi: \tilde{M}^n \rightarrow M^n$  denote a principal  $Z_2$ -bundle  $\xi$ , with  $\tilde{M}^n$  and  $M^n$  being  $n$ -manifolds. An *imbedding* of  $\xi$  into the trivial fiber bundle  $M^n \times F$  over  $M^n$  is a smooth map  $h: \tilde{M}^n \rightarrow M^n \times F$  which maps  $\tilde{M}^n$  diffeomorphically onto its image  $h(\tilde{M}^n)$  in  $M^n \times F$ , such that the following diagram is commutative,

$$\begin{array}{ccc}
 \tilde{M}^n & \xrightarrow{h} & M^n \times F \\
 \searrow \pi & & \swarrow \text{proj} \\
 & & M^n .
 \end{array}$$

Recently, V. L. Hansen [3] has shown that for connected topological spaces  $X$  with nondegenerate base point and with the homotopy type of a CW complex of dimension  $m \geq 1$ , any finite covering map  $\pi: E \rightarrow X$  can be imbedded into the trivial real  $(m+1)$ -plane bundle over  $X$ . Our notion of imbedding is the same as his, except we will work in the smooth category.

One naturally asks the following question. To what extent does the vanishing of some iterated cup product of the characteristic class  $w_1(\xi)$  of the involution  $\tilde{M}^n$  determine an imbedding of  $\xi$  into the trivial vector bundle  $M^n \times R^k$  over  $M^n$ ? The purpose of this paper is to investigate this question.

The vanishing of  $w_1^2$  has already appeared in the literature in other contexts, for example ([2], [8], [9]). We will be dealing, however, with general two-fold covers rather than just orientation covers.

A good reference then is [1], page 60.

2. Imbeddings. We begin with a normalization lemma.

LEMMA 2.1. *If the principal  $Z_2$ -bundle  $\pi: \tilde{M} \rightarrow M$  imbeds in the trivial vector bundle  $M \times R^k \rightarrow M$ , then there is an imbedding map  $g$  sending  $\tilde{M}$  to  $M \times S^{k-1}$ , such that fibers of  $\tilde{M} \rightarrow M$  are mapped antipodally into  $S^{k-1}$ .*

*Proof.* Let  $h$  be an imbedding of  $\tilde{M}$  into  $M \times R^k$ . For each  $x$  in  $M$ , let  $y$  and  $y'$  be the two distinct elements in the fiber  $\pi^{-1}(x)$ . Define  $g: \tilde{M} \rightarrow M \times S^{k-1}$  by

$$g(y) = \left( x, \frac{h_2(y) - h_2(y')}{\|h_2(y) - h_2(y')\|} \right)$$

where  $h = (h_1, h_2)$  and  $\| \quad \|$  is the standard norm in  $R^k$ . The map  $g$  is as required.

In light of the previous lemma, one has the following result.

PROPOSITION 2.2. *The principal  $Z_2$ -bundle  $\xi: \tilde{M} \rightarrow M$  imbeds in  $(M \times R^k \rightarrow M)$  if and only if  $\xi$  is classified by the diagram below with  $\tilde{f}$   $Z_2$ -equivariant.*

$$\begin{array}{ccc} \tilde{M} & \xrightarrow{\tilde{f}} & S^{k-1} \subset S^\infty \\ \downarrow & & \downarrow \\ M & \xrightarrow{f} & R p^{k-1} \subset R p^\infty . \end{array}$$

COROLLARY 2.3. *If the principal  $Z_2$ -bundle  $\xi: \tilde{M} \rightarrow M$  imbeds in  $(M \times R^k \rightarrow M)$ , then the  $k$ -fold cup product  $(w_1(\xi))^k$  of the characteristic class  $w_1(\xi)$  of the involution  $\tilde{M}$  is zero.*

*Proof.* One has the following diagram classifying  $\xi$ .

$$\begin{array}{ccc} \tilde{M} & \xrightarrow{\tilde{f}} & S^{k-1} \\ \downarrow & & \downarrow \\ M & \xrightarrow{f} & R p^{k-1} . \end{array}$$

Let  $c$  be the generator of  $H^1(R p^{k-1}; Z_2)$ . Then

$$(w_1(\xi))^k = (f^*(c))^k = f^*(c^k) = f^*(0) = 0 .$$

For  $k = 1$ , there is the following result.

LEMMA 2.4. *Let  $\xi$  denote the principal  $\mathbf{Z}_2$ -bundle  $\tilde{M} \rightarrow M$ . Then  $w_1(\xi) = 0$  if and only if  $\xi$  imbeds in  $(M \times \mathbf{R}^1 \rightarrow M)$ .*

*Proof.* If  $\xi$  imbeds in  $M \times \mathbf{R}^1$ , then  $w_1(\xi) = 0$ , by Corollary 2.3. On the other hand if  $w_1(\xi) = 0$ , the line bundle  $\gamma$  associated to  $\xi$  is orientable, and so the structure group  $\mathbf{Z}_2 = O(1)$  of  $\gamma$  reduces to  $SO(1) = \{1\}$ , the trivial group. Thus  $\gamma$  is a trivial  $\mathbf{R}$ -bundle, since it is classified in  $BSO(1) = \text{point}$ . The natural inclusion of  $\xi$  in  $\gamma$  shows that  $\xi$  imbeds in  $M \times \mathbf{R}^1$ .

Now let  $\xi$  denote the principal  $\mathbf{Z}_2$ -bundle  $\tilde{M}^n \rightarrow M^n$ , and let  $\eta$  denote the principal  $\mathbf{Z}_2$ -bundle  $\tilde{N}^{n-1} \rightarrow N^{n-1}$ , where  $\tilde{N}^{n-1}$  is a characteristic submanifold for  $\tilde{M}^n$ . We will need the following lemma.

LEMMA 2.5. *An imbedding of  $\eta$  in the trivial vector bundle  $N^{n-1} \times \mathbf{R}^{k-1} \rightarrow N^{n-1}$  gives rise to an imbedding of  $\xi$  in the trivial vector bundle  $M^n \times \mathbf{R}^k \rightarrow M^n$ .*

*Proof.* Since  $\tilde{N}$  is a characteristic submanifold for  $\tilde{M}$ , write  $\tilde{M}$  as

$$\tilde{P}_- \cup (\tilde{N} \times [-1, 1]) \cup \tilde{P}_+$$

with the involution on  $\tilde{M}$  interchanging the two copies  $\tilde{P}_-$  and  $\tilde{P}_+$  of  $\tilde{P}$ . Also, write  $M$  as

$$P \cup Q$$

with  $Q$  being the disk-bundle of  $N$  in  $M$ .

Given the imbedding  $h: \tilde{N} \rightarrow N \times \mathbf{R}^{k-1}$  covering  $N$ , extend  $h$  in the obvious fashion to an imbedding

$$g: \tilde{N} \times [-1, 1] \longrightarrow Q \times \mathbf{R}^{k-1} \times [-1, 1] \subseteq Q \times \mathbf{R}^k \text{ covering } Q.$$

Lemma 2.1 says that  $g$  may be taken as an imbedding of  $\tilde{N} \times [-1, 1]$  into  $Q \times S^{k-1}$ ; moreover,  $g$  may be chosen to take  $\tilde{N} \times \{-1, 1\}$  into  $\partial Q \times S^0$  for some inclusion of  $S^0$  in  $S^{k-1}$ . Since the disjoint union  $\tilde{P}_- + \tilde{P}_+$  is a trivial two-fold cover of  $P$ , the map  $g$  extends to an imbedding of  $\tilde{M}$  into  $M \times S^{k-1}$  covering  $M$ . Hence  $\xi$  imbeds in the trivial vector bundle  $M \times \mathbf{R}^k$ .

We are now ready to state the first theorem.

THEOREM 2.6. *Let  $\xi$  denote the principal  $\mathbf{Z}_2$ -bundle  $\tilde{M}^n \rightarrow M^n$  of closed manifold classified by  $w_1(\xi)$  in  $H^1(M^n; \mathbf{Z}_2)$ . The existence of a nested sequence  $\tilde{N}^{n-k+1} \subset \tilde{N}^{n-k+2} \subset \dots \subset \tilde{N}^{n-1} \subset \tilde{N}^n = \tilde{M}^n$  of characteristic submanifolds (corresponding to principal  $\mathbf{Z}_2$ -bundles  $\xi_{n-k+1} \subset$*

$\xi_{n-k+2} \subset \dots \subset \xi_{n-1} \subset \xi_n = \xi$ ) satisfying  $(w_1(\xi_i))^{k-n+i} = 0$ , for  $1 \leq k \leq n$  and all  $i$  with  $n - k + 1 \leq i \leq n$ , provides a necessary and sufficient condition for imbedding  $\xi$  in  $M \times \mathbf{R}^k$ .

*Proof.* Assume that  $\xi$  imbeds in  $M \times \mathbf{R}^k$ . By Proposition 2.2,  $\xi$  is classified by the diagram

$$\begin{array}{ccc} \tilde{M} & \xrightarrow{\tilde{f}} & S^{k-1} \\ \downarrow & & \downarrow \\ M & \xrightarrow{f} & \mathbf{R}p^{k-1}. \end{array}$$

Thus  $(w_1(\xi))^k = 0$ . Deform  $f$  to be transverse regular on  $\mathbf{R}p^{k-2}$ . Then  $f^{-1}(\mathbf{R}p^{k-2})$  is a 1-codimensional submanifold  $N^{n-1}$  of  $M^n$  with the pull-back covering  $\xi_{n-1}: \tilde{N}^{n-1} \rightarrow N^{n-1}$  satisfying  $(w_1(\xi_{n-1}))^{k-1} = 0$  and  $\tilde{N}^{n-1}$  a characteristic submanifold for  $\tilde{M}^n$ . Continuing to deform  $f$ , one gets  $(w_1(\xi_i))^{k-n+i} = 0$  for the rest of the  $\xi_i$  and the nested sequence of characteristic submanifolds. Thus necessity is proved.

To prove sufficiency induct on  $k \geq 1$ . For  $k = 1$ , see Lemma 2.4. As an inductive step,  $\xi_{n-1}$  imbeds in  $N^{n-1} \times \mathbf{R}^{k-1}$ , and along with Lemma 2.5 sufficiency is proved.

**3. Imbedding up to  $\mathbf{Z}_2$ -cobordism.** Let  $\mathcal{N}_*(\mathbf{Z}_2)$  denote the bordism module of fixed point free involutions as defined in [1, p. 59]. It is well-known that  $\mathcal{N}_*(\mathbf{Z}_2)$  is a free module over the unoriented bordism ring  $\mathcal{N}_*$  with base the class of the antipodal map on the  $j$ -sphere  $[A, S^j]$ , for  $j = 0, 1, 2, \dots$ . In  $\mathcal{N}_n(\mathbf{Z}_2)$ , the bordism class of the involution  $\tilde{M}^n$  is determined uniquely by the integers mod 2  $\langle w_{i(1)} \cdots w_{i(k)} (w_1(\xi))^m, [M^n] \rangle$  where  $\xi: \tilde{M}^n \rightarrow M^n$  is the 2-fold cover, the  $w_{i(j)}$  are the Stiefel-Whitney classes of  $M^n$ ,  $[M^n]$  is the fundamental class of  $M^n$ , and  $i(1) + \dots + i(k) + m = n$ . See [1, p. 60].

**THEOREM 3.1.** *Let  $\xi: \tilde{M}^n \rightarrow M^n$  be a principal  $\mathbf{Z}_2$ -bundle of closed manifolds and let  $k \geq 1$ . If  $(w_1(\xi))^k = 0$ , then there is a representative  $\tilde{H}^n$  of the class of  $\tilde{M}^n$  in  $\mathcal{N}_n(\mathbf{Z}_2)$  such that  $\eta: \tilde{H}^n \rightarrow H^n$  imbeds in  $H^n \times \mathbf{R}^k$ .*

*Proof.* Notice first of all that if  $k \geq n + 1$  or  $k = 1$ , then previous results allow one to take  $\tilde{H}^n = \tilde{M}^n$ . Suppose, therefore, that  $1 < k \leq n$ . Write  $\tilde{M}^n$  up to fixed point free  $\mathbf{Z}_2$ -cobordism as the disjoint union

$$\sum_{j=0}^n (A, S^j) \times V^{n-j}$$

where  $(A, S^j)$  denotes the antipodal map on the  $j$ -sphere and  $V^{n-j}$

denotes a closed  $(n - j)$ -manifold with trivial action. Let  $u \in H^1(\ ; \mathbf{Z}_2)$  generally denote the characteristic class of an involution. Note that for  $j < r$ ,  $(A, S^j) \times V^{n-j}$  has  $u^r = 0$ .

Assume  $(w_1(\xi))^n = 0$ , then

$$0 = \langle (w_1(\xi))^n, [M] \rangle = \left\langle u^n, \left[ \sum_{j=0}^n R p^j \times V^{n-j} \right] \right\rangle = \sum_{j=0}^n \langle u^n, [R p^j \times V^{n-j}] \rangle \\ = \langle u^n, [R p^n \times V^0] \rangle .$$

Thus  $V^0$  must represent the trivial element in  $\mathcal{N}_0$ . Now write  $\tilde{M}^n$  up to fixed point free  $\mathbf{Z}_2$ -cobordism as  $\sum_{j=0}^{n-1} (A, S^j) \times V^{n-j}$ . Moreover, since  $\mathcal{N}_1 = 0$ , one may as well write  $\sum_{j=0}^{n-2} (A, S^j) \times V^{n-j}$ .

Suppose inductively that the vanishing of  $(w_1(\xi))^n, (w_1(\xi))^{n-1}, \dots, (w_1(\xi))^{r+1}$  allows one to remove  $(A, S^n) \times V^0, (A, S^{n-1}) \times V^1, \dots, (A, S^{r+1}) \times V^{n-(r+1)}$  from the bordism representation. Let  $(w_1(\xi))^r = 0$ , and let  $p^{n-r}$  generally denote a homogeneous polynomial of degree  $(n - r)$  in the Stiefel-Whitney classes of a manifold. Then  $(w_1(\xi))^r = 0$  implies that

$$0 = \langle w_1(\xi)^r p^{n-r}, [M] \rangle = \left\langle u^r p^{n-r}, \sum_{j=0}^r [R p^j \times V^{n-j}] \right\rangle \\ = \sum_{j=0}^r \langle u^r p^{n-r}, [R p^j \times V^{n-j}] \rangle = \langle u^r p^{n-r}, [R p^r \times V^{n-r}] \rangle .$$

Since  $u$  is "dual" to  $R p^{r-1} \times V^{n-r}$ , (see [7, p. 155]),

$$0 = \langle u^r p^{n-r}, [R p^r \times V^{n-r}] \rangle = \langle u^{r-1} p^{n-r}, [R p^{r-1} \times V^{n-r}] \rangle .$$

Successive use of this duality shows  $\langle p^{n-r}, [V^{n-r}] \rangle = 0$  for all  $p^{n-r}$  on the Stiefel-Whitney classes of  $V^{n-r}$ . Thus  $V^{n-r}$  represents a trivial element in  $\mathcal{N}_{n-r}$ .

Hence  $\tilde{M}^n$  represents the same element in  $\mathcal{N}_*(\mathbf{Z}_2)$  as  $\tilde{H}^n$  which equals  $\sum_{j=0}^{k-1} (A, S^j) \times V^{n-j}$ . Clearly,  $\eta: \tilde{H}^n \rightarrow H^n$  imbeds  $H^n \times R^k$ .

REFERENCES

1. P. E. Conner and E. E. Floyd, *Differentiable Periodic Maps*, Springer-Verlag, Berlin-Heidelberg-New York, 1964.
2. J. L. Dupont and G. Lusztig, *On manifolds satisfying  $w_1^2 = 0$* , *Topology*, **10** (1971), 81-92.
3. V. L. Hansen, *Embedding finite covering spaces into trivial bundles*, *Math. Ann.*, **236** (1978), 239-243.
4. S.-T. Hu, *Cohomology Theory*, Markham, Chicago, 1968.
5. J. W. Milnor and J. D. Stasheff, *Characteristic Classes*, Princeton University Press, Princeton, 1974.
6. R. E. Mosher and M. C. Tangora, *Cohomology Operations and Applications in Homotopy Theory*, Harper and Row, New York-Evanston-London, 1968.
7. R. E. Stong, *Notes on Cobordism Theory*, Princeton University Press, Princeton, 1968.
8. E. Thomas, *Vector fields on manifolds*, *Bull. Amer. Math. Soc.*, **75** (1969), 643-683.

9. C. T. C. Wall, *Determination of the cobordism ring*, Ann. of Math., **72** (1960), 292-311.

Received May 4, 1979.

UNIVERSITY OF TENNESSEE  
KNOXVILLE, TN 37916

*Current address:* Bell Laboratories  
11900 No. Pecos St.  
Denver, CO 80234

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California  
Los Angeles, California 90024

HUGO ROSSI

University of Utah  
Salt Lake City, UT 84112

C. C. MOORE AND ANDREW OGG

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. FINN AND J. MILGRAM

Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON



David Bressoud, <i>A note on gap-frequency partitions</i> .....	1
John David Brillhart, <i>A double inversion formula</i> .....	7
Frank Richard Deutsch, Günther Nürnberger and Ivan Singer, <i>Weak Chebyshev subspaces and alternation</i> .....	9
Edward Richard Fadell, <i>The relationship between Ljusternik-Schnirelman category and the concept of genus</i> .....	33
Harriet Jane Fell, <i>On the zeros of convex combinations of polynomials</i> .....	43
John Albert Fridy, <i>An addendum to: "Tauberian theorems via block dominated matrices"</i> .....	51
Andrzej Granas, Ronald Bernard Guenther and John Walter Lee, <i>Applications of topological transversality to differential equations. I. Some nonlinear diffusion problems</i> .....	53
David E. Handelman and G. Renault, <i>Actions of finite groups on self-injective rings</i> .....	69
Michael Frank Hutchinson, <i>Local <math>\Lambda</math> sets for profinite groups</i> .....	81
Arnold Samuel Kas, <i>On the handlebody decomposition associated to a Lefschetz fibration</i> .....	89
Hans Keller, <i>On the lattice of all closed subspaces of a Hermitian space</i> .....	105
P. S. Kenderov, <i>Dense strong continuity of pointwise continuous mappings</i> .....	111
Robert Edward Kennedy, <i>Krull rings</i> .....	131
Jean Ann Larson, Richard Joseph Laver and George Frank McNulty, <i>Square-free and cube-free colorings of the ordinals</i> .....	137
Viktor Losert and Harald Rindler, <i>Cyclic vectors for <math>L^p(G)</math></i> .....	143
John Rowlay Martin and Edward D. Tymchatyn, <i>Fixed point sets of 1-dimensional Peano continua</i> .....	147
Augusto Nobile, <i>On equisingular families of isolated singularities</i> .....	151
Kenneth Joseph Prevot, <i>Imbedding smooth involutions in trivial bundles</i> .....	163
Thomas Munro Price, <i>Spanning surfaces for projective planes in four space</i> .....	169
Dave Riffelmacher, <i>Sweedler's two-cocycles and Hochschild cohomology</i> .....	181
Niels Schwartz, <i>Archimedean lattice-ordered fields that are algebraic over their <math>o</math>-subfields</i> .....	189
Chao-Liang Shen, <i>A note on the automorphism groups of simple dimension groups</i> .....	199
Kenneth Barry Stolarsky, <i>Mapping properties, growth, and uniqueness of Vieta (infinite cosine) products</i> .....	209
Warren James Wong, <i>Maps on simple algebras preserving zero products. I. The associative case</i> .....	229