

Pacific Journal of Mathematics

A CHARACTERIZATION OF THE WEIGHTS IN A DIVIDED DIFFERENCE

DONALD J. NEWMAN AND THEODORE JOSEPH RIVLIN

A CHARACTERIZATION OF THE WEIGHTS IN A DIVIDED DIFFERENCE

D. J. NEWMAN AND T. J. RIVLIN

Given points x_0, x_1, \dots, x_n of the real line which satisfy $x_0 < x_1 < \dots < x_n$ put

$$w(u) = (u - x_0) \cdots (u - x_n).$$

If function values $f(x_0), \dots, f(x_n)$ are given, the divided difference of f with respect to x_0, \dots, x_n may be defined by

$$(1) \quad f(x_0, \dots, x_n) = \sum_{i=0}^n \frac{f(x_i)}{w'(x_i)}.$$

If we put

$$(2) \quad a_i = \frac{1}{w'(x_i)}, \quad i = 0, \dots, n$$

we call a_0, \dots, a_n the weights of the divided difference. Our problem is to characterize the vectors $a: (a_0, \dots, a_n)$ which satisfy (2) for some $x: (x_0, \dots, x_n)$ and to show that such a correspond to an essentially unique x .

Note that

$$(3) \quad w'(x_i) = (x_i - x_0) \cdots (x_i - x_{i-1})(x_i - x_{i+1}) \cdots (x_i - x_n), \\ i = 0, \dots, n,$$

so that translating the point x leaves a unchanged. Therefore we add the further (normalizing) assumption that $x_0 = 0$. We can now state our result.

THEOREM. *If $n > 0$ (2) holds for unique x_0, \dots, x_n satisfying $0 = x_0 < x_1 < \dots < x_n$ if, and only if,*

(i) $(-1)^{n-i} a_i > 0, \quad i = 0, \dots, n$

(ii) $\sum_{i=0}^n a_i = 0$

and

(iii) $(-1)^{n-j} \sum_{i=j}^n a_i > 0, \quad j = 1, \dots, n.$

Note that this theorem may be viewed as characterizing the slopes at its zeros of a polynomial $w(u) = u(u - x_1) \cdots (u - x_n)$.

Proof. (a) Suppose x_0, \dots, x_n satisfying $0 = x_0 < x_1 < \dots < x_n$ are given. (i) follows immediately from (3). Since $f(x_0, \dots, x_n) = 0$ if f is a polynomial of degree $< n$, taking $f \equiv 1$ in (1) yields (ii). As

for (iii) consider the fundamental polynomials of Lagrangian interpolation in x_0, \dots, x_n

$$l_i(x) = \frac{w(x)}{(x - x_i)w'(x_i)}, \quad i = 0, \dots, n.$$

Put

$$p_j(x) = \sum_{i=0}^j l_i(x), \quad j = 0, \dots, n - 1.$$

Then $p_j(x_i) = 1$, $i = 0, \dots, j$, $p_j(x_i) = 0$, $i = j + 1, \dots, n$ and so p_j is of positive degree. If c_j is the coefficient of x^n in p_j then

$$c_j = \sum_{i=0}^j a_i.$$

Now p'_j is of degree at most $n - 1$ and by Rolle's theorem has j zeros in (x_0, x_j) and $n - 1 - j$ zeros in (x_{j+1}, x_n) which accounts for all its zeros. Thus $c_j \neq 0$ and, since the behavior of $p_j(x)$ at ∞ determines the sign of c_j , this sign is that of $p'_j(x_n)$. We proceed to determine that sign. To this end note that $p'_j(x_j) \neq 0$. If $p'_j(x_j) > 0$ then since $p_j(x_j) = 1$ and $p_j(x_{j+1}) = 0$, $p'_j(x)$ has a zero in (x_j, x_{j+1}) , implying that $p_j = 0$, and contradicting the fact that p_j is of positive degree. Thus $p'_j(x_j) < 0$ which implies that $p'_j(x_{j+1}) < 0$ and hence that $p'_j(x_n)$ has the sign $(-1)^{n-j}$. We have shown that

$$(-1)^{n-j} \sum_{i=0}^j a_i > 0, \quad j = 0, \dots, n - 1,$$

and hence for $j = 0, \dots, n - 1$

$$(-1)^{n-j} \sum_{i=j+1}^n a_i = (-1)^{n-j} \left(\sum_{i=0}^n a_i - \sum_{i=0}^j a_i \right) < 0$$

in view of (ii), from which (iii) follows.

(b) We prove next that for each a satisfying (i), (ii), (iii) there exist unique x_0, \dots, x_n satisfying $0 = x_0 < \dots < x_n$ such that (2) holds.

Let K denote the convex set in R^n defined by the inequalities

$$(4) \quad (-1)^{n-t} t_i > 0, \quad i = 1, \dots, n,$$

and

$$(5) \quad (-1)^{n-j} \sum_{i=j}^n t_i > 0, \quad j = 1, \dots, n.$$

In view of (i) and (iii) K is not empty. Let S be the convex set of all $(x_1, \dots, x_n) \in R^n$ such that $0 < x_1 < x_2 < \dots < x_n$. Let ϕ be the continuous mapping of S into K defined by

$$\phi: (x_1, \dots, x_n) \longrightarrow (a_1, \dots, a_n)$$

where the a_i are as defined in (2) with $x_0 = 0$. We intend to show that ϕ is a global homeomorphism of S onto K . If this were so then given a_0, \dots, a_n satisfying (i), (ii) and (iii), (a_1, \dots, a_n) is in K and its unique preimage (x_1, \dots, x_n) satisfies $0 = x_0 < x_1 < \dots < x_n$ and $a_i = (w'(x_i))^{-1}$, $i = 1, \dots, n$. But then in view of (ii)

$$a_0 = -\sum_{i=1}^n a_i = -\sum_{i=1}^n \frac{1}{w'(x_i)} = \frac{1}{w'(x_0)},$$

and the proof of the theorem would be complete. It remains, therefore, to show that ϕ is a homeomorphism of S onto K .

We begin by showing that the Jacobian matrix of ϕ , $J(\phi)$ is nonsingular in S . Given $0 = x_0 < x_1 < \dots < x_n$ the divided difference $f(x_0, \dots, x_n) = 0$ if f is a polynomial of degree at most $n - 1$ while if $f = x^n$, $f(x_0, \dots, x_n) = 1$. If we take f to be x, x^2, \dots, x^n successively we thus obtain

$$\sum_{i=1}^n a_i x_i^j = 0, \quad j = 1, \dots, n - 1,$$

$$\sum_{i=1}^n a_i x_i^n = 1.$$

Partial differentiation with respect to x_i gives

$$\begin{pmatrix} x_1 & x_2 & \dots & x_n \\ x_1^2 & x_2^2 & \dots & x_n^2 \\ \vdots & \vdots & \ddots & \vdots \\ x_1^n & x_2^n & \dots & x_n^n \end{pmatrix} \begin{pmatrix} \frac{\partial a_1}{\partial x_i} \\ \frac{\partial a_i}{\partial x_i} \\ \vdots \\ \frac{\partial a_n}{\partial x_i} \end{pmatrix} + \begin{pmatrix} 1 \\ 2x_i \\ 3x_i^2 \\ \vdots \\ nx_i^{n-1} \end{pmatrix} a_i = 0$$

so that,

$$AJ(\phi) + B: = \begin{pmatrix} x_1 & x_2 & \dots & x_n \\ \vdots & \vdots & \ddots & \vdots \\ x_1^n & x_2^n & \dots & x_n^n \end{pmatrix} J(\phi) + \begin{pmatrix} a_1 & \dots & a_n \\ 2a_1 x_1 & & 2a_n x_n \\ \vdots & & \vdots \\ na_1 x_1^{n-1} & & na_n x_n^{n-1} \end{pmatrix} = 0.$$

Since A and B are nonzero multiples of the Vandermonde matrix they are nonsingular and so, therefore, is $J(\phi)$.

Next we show that ϕ maps the boundary of S (which includes the point at infinity) into the boundary of K (which includes the point at infinity) in the sense that if a sequence of points of S has a limit point in ∂S then the image sequence under ϕ has its limit points in ∂K . The boundary points of S arise either from coalescence

of coordinates or coordinates tending to zero or infinity, or combinations of these. (3) reveals that if 2 or more coordinates coalesce with x_i and stay bounded away from 0 and ∞ then $w'(x_i) \rightarrow 0$ and hence $a_i \rightarrow \infty$. If $x_1 \rightarrow 0$ and $w'(x_1) \rightarrow 0$, or $x_n \rightarrow \infty$ and $w'(x_n) \rightarrow \infty$, boundary goes to boundary. The only remaining possibility is that $x_1 \rightarrow 0$ and $x_n \rightarrow \infty$ in such a way that $\{|w'(x_i)|\}$, $i = 1, \dots, n$ stays bounded away from 0 and ∞ . This can only happen if for some j satisfying $2 \leq j \leq n - 1$ we have $x_{j-1} \rightarrow 0$ and $x_j \rightarrow \infty$. But in this case put $p(x) = x(x - x_1) \cdots (x - x_{j-1})$ and $q(x) = (x - x_j) \cdots (x - x_n)$. Then on the one hand eventually

$$\frac{1}{2\pi i} \int_{|z|=1} \frac{dz}{p(z)q(z)} = \sum_{i=0}^{j-1} \frac{1}{w'(x_i)}$$

by the calculus of residues, while on the other the integral tends to zero since $|q| \rightarrow \infty$. Thus

$$\lim_{i=0}^{j-1} a_i = \lim_{i=j}^n a_i = 0,$$

so that the image sequence again tends to the boundary of K .

We are now in a position to show that ϕ maps S onto K . Since $J(\phi)$ is nonsingular, $\phi(S)$ is open in K . Let $w \in \phi(S)^\circ \cap K$, we claim that some neighborhood of w lies in $\phi(S)^\circ \cap K$. Otherwise there is a sequence $w_i = \phi(z_i)$ with $w_i \rightarrow w$. If the z_i went to the boundary, ∂S , then the $\phi(z_i) (= w_i)$ would converge to the boundary ∂K and we would have $w \in \partial K$. This contradicts our assumption and we conclude, indeed, that $\phi(S)^\circ \cap K$ is open. Thus $K = (\phi(S)^\circ \cap K) \cup (\phi(S) \cap K)$ is the union of two disjoint open sets. K , being convex, is certainly connected, however, and so we are forced to the conclusion that $\phi(S) \cap K = K$ or $\phi(S) = K$.

Thus, to sum up, we have shown that ϕ is a continuous map of convex S onto convex K which is locally one-to-one and takes the boundary of S into the boundary of K .

Next we show, following Favard (Cf. J. Favard, Cours d'Analyse de l'École Polytechnique, Vol. I, Paris, 1960, pp. 295-296.) that there is a fixed integer $k \geq 1$ such that each point of K has exactly k preimages in S .

First we observe that if $t \in K$ then $t = \phi(x)$ cannot hold for infinitely many $x \in S$. For if $t = \phi(x)$ for infinitely many $x \in S$ then the set of such x cannot have a boundary point of S as limit point since $t \in K$ and ϕ takes ∂S into ∂K . Thus the set of solutions $\{x\}$ has a limit point in S contradicting the fact that ϕ is a local homeomorphism.

Suppose that $\bar{t} \in K$ and $x^{(1)}(\bar{t}), \dots, x^{(s)}(\bar{t})$ are its preimages. The inverse functions $x = \Psi_1(t), \dots, x = \Psi_s(t)$ are homeomorphisms of

some neighborhood of \bar{t} , and these neighborhoods have a nonempty intersection. Let $r(\bar{t})$ be the radius of the largest ball centered at \bar{t} in which Ψ_1, \dots, Ψ_s are homeomorphisms. It is easy to see that $|r(t^{(1)}) - r(t^{(2)})| \leq \|t^{(1)} - t^{(2)}\|$, hence $r(t)$ is continuous in K and has a positive lower bound on every compact subset of K . Now every point t' contained in the largest ball of homeomorphy centered at t has at least as many preimages as does t , i.e., $s(t') \geq s(t)$. On the other hand in the closed ball $\|t - t'\| \leq r(t)/2$, $r(t')$ has a positive lower bound, b , and if we further restrict t' so that $\|t - t'\| < b$ then the largest ball of homeomorphy centered at t' contains t so that $s(t) \geq s(t')$, hence $s(t) = s(t')$ and $s(t)$ is a continuous integer-valued function of t in the convex set K , hence a constant, $k \geq 1$.

We next conclude the proof of our theorem by exhibiting an explicit t which has a *unique* pre-image. To this end consider the points

$$\eta_j = -\cos \frac{j\pi}{n}, \quad j = 0, \dots, n,$$

the extrema of the Chebyshev polynomial of the first kind of degree n . An elementary computation yields

$$\begin{aligned} f(\eta_0, \dots, \eta_n) &= (-1)^n \frac{2^{n-1}}{n} \left(\frac{f(\eta_0)}{2} + \sum_{j=1}^{n-1} (-1)^j f(\eta_j) + \frac{(-1)^n}{2} f(\eta_n) \right) \\ &= : (-1)^n \frac{2^{n-1}}{n} \sum_{j=0}^n (-1)^j f(\eta_j) \end{aligned}$$

and taking $f(u)$ to be u, u^2, \dots, u^{n-1} successively yields

$$(6) \quad \sum_{j=0}^n (-1)^j \eta_j^i = 0, \quad i = 1, \dots, n-1.$$

Suppose $y_0 < y_1 < \dots < y_n$ has the property that,

$$(7) \quad \frac{(-1)^n}{W'(y_i)} = \frac{(-1)^n}{V'(\eta_i)} = \begin{cases} \frac{2^{n-1}}{2n}; & i = 0, \\ \frac{2^{n-1}}{n} (-1)^i; & i = 1, \dots, n-1, \\ \frac{2^{n-1}}{2n} (-1)^n; & i = n \end{cases}$$

where $V(u) = (u - \eta_0) \dots (u - \eta_n)$, $W(u) = (u - y_0) \dots (u - y_n)$ and we may assume, with no loss of generality, that $y_0 = -y_n$. Then we have

$$\sum_{j=0}^n (-1)^j y_j^i = 0, \quad i = 1, \dots, n-1,$$

and if we put

$$x_j = \frac{y_j}{y_n} =: Cy_j, \quad j = 0, \dots, n$$

then

$$(8) \quad \sum_{j=0}^n (-1)^j x_j^i = 0, \quad i = 1, \dots, n-1,$$

$$x_0 = \eta_0 = -1; \quad x_n = \eta_n = 1$$

and $x_0 < x_1 < \dots < x_n$.

Equating the left-hand sides in (6) to the corresponding ones in (8) yields the systems

$$(9a) \quad x_1^i + \eta_2^i + \dots + x_{n-1}^i = \eta_1^i + x_2^i + \dots + \eta_{n-1}^i, \quad i = 1, \dots, n-1;$$

n even or

$$(9b) \quad x_1^i + \eta_2^i + \dots + \eta_{n-1}^i = \eta_1^i + x_2^i + \dots + x_{n-1}^i, \quad i = 1, \dots, n-1;$$

n odd. We claim that (9a) or (9b) implies that $\eta_i = x_i$, $i = 1, \dots, n-1$. To establish this we use the following lemma.

LEMMA. *If*

$$\sum_{j=1}^k b_j^i = \sum_{j=1}^k c_j^i, \quad i = 1, \dots, k$$

and $b_1 < \dots < b_k$; $c_1 < \dots < c_k$ then $b_j = c_j$, $j = 1, \dots, k$.

Proof. Let

$$S_i(b) = \sum_{j=1}^k b_j^i, \quad i = 1, \dots, k.$$

We assume $S_i(b) = S_i(c)$, $i = 1, \dots, k$. The elementary symmetric functions $\sigma_1(b), \dots, \sigma_k(b)$ are uniquely determined by $S_1(b), \dots, S_k(b)$ hence $\sigma_i(b) = \sigma_i(c)$, $i = 1, \dots, k$. Thus (c_1, \dots, c_k) and (b_1, \dots, b_k) are the complete set of zeros of the same polynomial. Since they are ordered by hypothesis, the lemma follows.

The lemma applied to (9a), say, implies that $x_1, \eta_2, \dots, x_{n-1}$ is a rearrangement of $\eta_1, x_2, \dots, \eta_{n-1}$. Thus in the sequence $\eta_1, \dots, \eta_{n-1}, x_1, \dots, x_{n-1}$ each of $\eta_1, \dots, \eta_{n-1}$ appears twice. But then the monotonicity of the vectors η and x implies that $x_j = \eta_j$, $j = 1, \dots, n-1$. The same argument prevails if (9b) holds. Thus we have shown that

$$y_j = \frac{\eta_j}{C}, \quad j = 0, \dots, n.$$

But, in view of the analogue of (3), (7) implies that $C = 1$.

We have established that the weight vector

$$(-1)^n \frac{2^{n-1}}{n} \left(\frac{1}{2}, -1, +1, \dots, (-1)^{n-1}, \frac{(-1)^n}{2} \right)$$

corresponds uniquely to the (normalized) points

$$1 + \eta_j, \quad j = 0, \dots, n$$

and so we may conclude that $((1 + \eta_1), \dots, (1 + \eta_n))$ is the unique element of S whose image under ϕ is

$$(-1)^n \frac{2^{n-1}}{n} \left(-1, 1, \dots, (-1)^{n-1}, \frac{(-1)^n}{2} \right).$$

This completes the proof of the theorem.

Postscript. I. J. Schoenberg reports (Cardinal Spline Interpolation, Vol. 12, Regional Conference Series in Applied Mathematics, Philadelphia, 1973, p. 9) that S. Karlin told him that he had solved the problem of characterizing the weights in a divided difference. Professor Karlin has confirmed this in a phone conversation with one of us, but not wishing to publish his solution suggested that we present ours.

Received June 13, 1979.

TEMPLE UNIVERSITY
PHILADELPHIA, PA 19122
AND
IBM
Box 218
YORKTOWN HEIGHTS, NY 10598

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$102.00 a year (6 Vols., 12 issues). Special rate: \$51.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1981 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Ilya Eugene Blum and Srinivasa Swaminathan, Continuous selections and realcompactness	251
Lawrence James Brenton, Differentiably k -normal analytic spaces and extensions of holomorphic differential forms	261
Jo-Ann Deborah Cohen, Topologies on the ring of integers of a global field	269
Robert Jay Daverman, Detecting the disjoint disks property	277
Edmund H. Feller, Rings where the annihilators of α -critical modules are prime ideals	299
Richard Elam Heisey and Henryk Toruńczyk, On the topology of direct limits of ANRs	307
Gerald William Johnson and David Lee Skoug, Notes on the Feynman integral. I	313
Michael S. Keane and Stuart Jay Sidney, Distinguishing a plane curve from other curves similar to it	325
Leonid A. Luxemburg, On compact metric spaces with noncoinciding transfinite dimensions	339
Chun Ming Ma, A uniqueness theorem for Navier-Stokes equations	387
Donald J. Newman and Theodore Joseph Rivlin, A characterization of the weights in a divided difference	407
Marc Aristide Rieffel, C^* -algebras associated with irrational rotations	415
Kichi-Suke Saito, Invariant subspaces for finite maximal subdiagonal algebras	431
Frederic W. Shultz, Dual maps of Jordan homomorphisms and $*$ -homomorphisms between C^* -algebras	435
Vsevolod Alekseevich Solonnikov, On the solvability of boundary and initial-boundary value problems for the Navier-Stokes system in domains with noncompact boundaries	443
Tavan Thomas Trent, New conditions for subnormality	459
L. E. Ward, Extending Whitney maps	465
Leslie Wilson, Jets with regular zeros	471
Sergio Eduardo Zarantonello, The sheaf of H^p -functions in product domains	479