

Pacific Journal of Mathematics

NOTE ON BOUNDED L^p -SOLUTIONS OF A GENERALIZED LIÉNARD EQUATION

ALLAN J. KROOPNICK

NOTE ON BOUNDED L^p -SOLUTIONS OF A GENERALIZED LIÉNARD EQUATION

ALLAN J. KROOPNICK

Two theorems are presented giving sufficient conditions for all solutions to $y'' + c(t)f(y)y' + a(t)b(y) = 0$ to be bounded. Furthermore, two other theorems are given guaranteeing when these solutions are L^p -solutions. Asymptotic stability is then discussed as well as several applications of these results.

In this paper, sufficient conditions will be given so that all solutions to a generalized Liénard equation,

$$(1) \quad y'' + c(t)f(y)y' + a(t)b(y) = 0$$

will be L^p -solutions ($p \geq 1$) on $[0, \infty)$. By an L^p -solution we shall mean a solution to (1) such that $\left(\int_0^\infty |y|^p dt\right) < \infty$. This note shall generalize some previous results (see [1] - [3]). We first need the following theorem.

THEOREM I. *Suppose $a(t)$ and $c(t)$ are continuous functions on $[0, \infty)$ and let $b(y)$ and $f(y)$ be continuous on $(-\infty, +\infty)$. Furthermore, suppose for some positive constant a_0 , $a(t) \geq a_0$, $a'(t) \leq 0$, $c(t) \geq 0$ for $0 \leq t < \infty$, and $f(y) > 0$. Finally, if $B(y) = \int_0^y b(u)du \rightarrow +\infty$ as $|y| \rightarrow \infty$, then every solution of (1) exists on $[0, \infty)$ and $|y(t)|, |y'(t)|$ are bounded as $t \rightarrow \infty$.*

Proof. By standard existence theory (1) has at least one solutions satisfying $y(0) = y_0$, $y'(0) = \dot{y}_0$, and existing on some interval $[0, T)$, $T > 0$. Consider any such solutions of (1) on $[0, T)$. Multiply (1) by y' and integrate from 0 to $t < T$ to obtain,

$$(2) \quad \frac{1}{2}y'(t)^2 + \int_0^t c(s)f(y)y'^2 ds + \int_0^t a(s)b(y)y' ds = \frac{1}{2}\dot{y}_0^2.$$

Integrating (2) by parts we have

$$(3) \quad \frac{1}{2}y'(t)^2 + \int_0^t c(s)f(y)y'^2 ds + a(t)B(y(t)) - \int_0^t a'(s)B(y(s)) ds = \frac{1}{2}\dot{y}_0^2 + a(0)B(y_0) \quad (0 \leq t < T).$$

For $|y|, |y'|$ large all terms on the LHS of (3) are positive except,

perhaps, for $-\int_0^t a'(s)B(y)ds$. We show that this term is bounded from below. By our hypotheses, $B(y) > -d^2$ for some real constant d and $-\int_0^t a'(s)B(y(s))ds > d^2\int_0^t a'(s)ds > -a(0)d^2$. Since

$$\int_0^t c(s)f(y(s))(y'(s))^2ds \geq 0$$

and since $a(t)B(y(t)) > -a(0)d^2$, (3) becomes

$$\begin{aligned} (3') \quad -2a(0)d^2 &< \frac{1}{2}y'^2(t) + a(t)B(y(t)) - \frac{1}{2}\int_0^t a'(s)B(y(s))ds \\ &\leq \frac{1}{2}y_0^2 + a(0)B(y_0) \quad (0 \leq t < T). \end{aligned}$$

By assumption $B(y) \rightarrow +\infty$ as $|y| \rightarrow \infty$, and the left and right sides of the inequality (3') are a priori lower and upper bounds, independent of T . Therefore, a standard argument completes the proof of global existence of solutions y such that $|y(t)|$ and $|y'(t)|$ are bounded on $[0, \infty)$.

REMARKS (1). If one also assumes $yb(y) \geq 0$, then all terms on the LHS of (3) are positive making the proof considerably simpler. (2) If the assumptions concerning a, c in Theorem I hold on an interval $[t_0, \infty)$ for some $t_0 \neq 0$, the conclusions are valid on the interval $[t_0, \infty)$. The same remark applies to Theorems II, III, and IV below.

We now proceed to our main theorems.

THEOREM II. *Let the hypotheses of Theorem I hold. In addition, let there exist constants $c_0 > 0, f_0 > 0, M > 0, p \geq 1$ such that $c(t) \geq c_0, c'(t) \leq 0$ ($0 \leq t < \infty$), and $f(y) \geq f_0, yb(y) \geq M|y|^p$ ($y \in \mathbf{R}$). Then $\int_0^\infty |y(t)|^p dt < \infty$.*

Proof. From (3) (which now holds on $0 < t < \infty$) we see that $c_0f_0 \int_0^\infty y'(t)^2 dt \leq (1/2)y_0^2 + a(0)B(y_0)$, so y' is square integrable. Multiply (1) by y and integrate by parts from 0 to t obtaining,

$$(4) \quad y(t)y'(t) - \int_0^t y'(s)^2 ds + \int_0^t c(s)f(y)yy'ds + \int_0^t a(s)b(y)yds = y_0y_0.$$

Letting $F(y) = \int_0^y uf(u)du$ we have upon another integration by parts,

$$(5) \quad \begin{aligned} y(t)y'(t) - \int_0^t y'(s)^2 ds + c(t)F(y(t)) - \int_0^t c'(s)F(y) ds \\ + \int_0^t a(s)b(y)y ds = y_0\dot{y}_0 + c(0)F(y_0). \end{aligned}$$

As $t \rightarrow \infty$ all terms on the LHS of (5) stay finite. Specifically, (5) shows that

$$(6) \quad 0 \leq \alpha_0 M \int_0^t |y(s)|^p ds \leq K_1 \quad (0 \leq t < \infty)$$

where $K_1 = |y_0||\dot{y}_0| + K^2 + 3c(0) \sup_{-K \leq y \leq K} F(y) + \int_0^\infty y'(s)^2 ds$ and K is the bound of $|y|$ and $|y'|$ on $[0, \infty)$. Since K_1 is independent of t , the result follows.

REMARK. Theorem II is still true under the assumptions $c_0 \leq c(t) \leq c_1$ ($c_0, c_1 > 0$) and $c' \in L^1[0, \infty)$ because

$$(7) \quad \left| \int_0^t c'(s)F(y) ds \right| \leq \sup_{-K \leq y \leq K} F(y) \int_0^\infty |c'(s)| ds.$$

This implies all terms on the LHS of (5) are bounded so that inequality (6) still holds although K_1 will be different.

The last two theorems use the well-known Gronwall-Bellman inequality.

THEOREM III. *The hypotheses are the same as Theorem I except that $a'(t) \geq 0$ ($0 \leq t < \infty$). Furthermore, if $yb(y) \geq 0$ ($y \in \mathbf{R}$), then all solutions to (1) are bounded as $t \rightarrow \infty$.*

Proof. From (3) we obtain the following inequality,

$$(8) \quad 0 \leq a(t)B(y(t)) \leq K + \int_0^t a'(s)B(y(s)) ds$$

where $K = \alpha_0 B(y_0) + (1/2)\dot{y}_0^2$. (8) may be rewritten as

$$(9) \quad 0 \leq a(t)B(y(t)) \leq K + \int_0^t \frac{a'(s)}{a(s)} a(s)B(y(s)) ds.$$

Applying the Gronwall-Bellman inequality to (9) yields,

$$(10) \quad a(t)B(y(t)) \leq K \left(\exp \int_0^t \frac{a'(s)}{a(s)} ds \right) = K \frac{a(t)}{a(0)}$$

or

$$(11) \quad B(y(t)) \leq \frac{K}{a_0}.$$

Since $B(y) \rightarrow \infty$ as $|y| \rightarrow \infty$, $y(t)$ must stay finite on $0 \leq t < \infty$.

REMARK. If one does not assume $a'(t) \leq 0$ (or $a'(t) \geq 0$) for $0 \leq t < \infty$, but only that $0 < a_0 \leq a(t) < \infty$, and $a' \in L^1[0, \infty)$, one can still obtain the conclusion that $y(t)$ remains bounded, for from (8) one has

$$(12) \quad a(t)B(y(t)) \leq K + \frac{1}{a_0} \int_0^t |a'(s)|a(s)B(y(s))ds .$$

Thus by the Gronwall-Bellman inequality

$$(13) \quad a(t)B(y(t)) \leq K \left(\exp \left(\frac{1}{a_0} \int_0^\infty |a'(s)| ds \right) \right) < \infty .$$

By strengthening the hypotheses of Theorem III slightly, we are able to show that all solutions are in $L^p[0, \infty)$. This is the substance of our final theorem.

THEOREM IV. *The hypotheses are the same as Theorem III. In addition, let there exist positive constants c_0 and f_0 such that $c(t) \geq c_0$ and $f(y) \geq f_0$, $c'(t) \leq 0$, $yb(y) \geq M|y|^p$ (for some positive constants M and $p \geq 1$), and if $a(t) \leq A_0$ where A_0 is a fixed positive constant, then $\int_0^\infty |y|^p dt < \infty$.*

Proof. From (3) we see that if $a(t)$ is bounded from above, then $\int_0^t a'(s)B(y)ds$ is bounded and, therefore, y' is bounded since all other terms in (3) are bounded. The remainder of the proof is identical to Theorem II and the remark after theorem is still true.

REMARKS. Had we only required $yb(y) \geq 0$, then Theorems II and IV would yield $\int_0^\infty y(t)b(y(t))dt < \infty$.

Actually, more may be proven. Under the hypotheses of Theorem II, the solutions to (1) are asymptotically stable (cf. Lemma 2.2 and Remarks 2.1 and 2.2 of [2]). In order to see this, write (1) as the following two dimensional system,

$$(14) \quad \begin{aligned} y_1' &= y_2 \\ y_2' &= -c(t)f(y_1)y_2 - a(t)b(y_1) . \end{aligned}$$

Now consider the following Liapunov function $V(t, y_1, y_2) = (1/2)y_2^2 + a(t)B(y_1)$. Differentiating we have $dV/dt = y_2y_2' + a'(t)B(y_1) + a(t)b(y_1)y_1' = a'(t)B(y_1) - c(t)f(y_1)y_2^2 < 0$ for $(y_1, y_2) \neq (0, 0)$ implying the solutions are asymptotically stable.

EXAMPLE 1. Consider the differential equation

$$(15) \quad y'' + \frac{1}{t^n}y = 0 \quad (0 < t_0 \leq t < \infty, n \text{ a positive integer}).$$

The substitution $y = x(t) \exp(t)$ transforms (15) into

$$(16) \quad x'' + 2x' + \left(1 + \frac{1}{t^n}\right)x = 0 \quad (0 < t_0 \leq t < \infty).$$

Therefore, applying Theorem II and Remark 2 following Theorem I, any solution to (15) is the product of the exponential function and a function in $L^2[a, \infty)$ ($a > 0$) since $xb(x) = x^2$.

EXAMPLE 2. It is well-known that the solutions to the homogeneous Duffing equation

$$(17) \quad y'' + cy' + ay(1 + dy^2) = 0 \quad (a, c, d > 0)$$

are asymptotically stable. By Theorem II the solutions are in both $L^2[0, \infty)$ and $L^4[0, \infty)$ since $yb(y) = y^2(1 + dy^2) \geq y^2$ and $yb(y) \geq dy^4$.

EXAMPLE 3. Emden's equation

$$(18) \quad y'' + \frac{2}{x}y' + y^n = 0 \quad (n \text{ a positive integer})$$

has important applications in astrophysics. When n is an odd positive integer, all solutions to (18) are bounded on $0 < x_0 \leq x < \infty$ using Theorem I and Remark 2 following Theorem I.

The author would like to thank the referee for several helpful suggestions concerning this paper.

REFERENCES

1. A. Kroopnick, L^2 -solutions to $y'' + c(t)y' + a(t)b(y) = 0$, Proc. Amer. Math. Soc., **39** (1973), 217-218.
2. J. J. Levin and J. A. Nohel, *Global asymptotic stability for nonlinear systems of differential equations and applications to reactor dynamics*, Archive Rat. Mech. Anal., **5** (1960), 194-211.
3. A. Strauss, *Liapunov functions and L^p -solutions of differential equations*, Trans. Amer. Math. Soc., **119** (1956), 37-50.

Received April 10, 1978 and in revised form August 10, 1980.

SOCIAL SECURITY ADMINISTRATION
OFFICE OF RESEARCH AND STATISTICS
MEADOWS EAST BUILDING
6401 SECURITY BLVD
BALTIMORE, MD 21235

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE AND ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Willy Brandal, Conditions for being an FGC domain	1
Allan Calder and Frank Williams, Incompressibility of maps and the homotopy invariance of Čech cohomology	13
Jacques Chaumat, Quelques propriétés du couple d'espaces vectoriels $(L^1m/H^{\infty\perp}, H^{\infty})$	21
Manfred Droste and Rüdiger Göbel, Products of conjugate permutations ...	47
Jean Esterle, Rates of decrease of sequences of powers in commutative radical Banach algebras	61
Allan Fryant, Ultraspherical expansions and pseudo analytic functions	83
John Hannah, Homogenization of regular rings of bounded index. II	107
Shigeru Haruki, On the theorem of S. Kakutani-M. Nagumo and J. L. Walsh for the mean value property of harmonic and complex polynomials	113
Hugh M. Hilden, Representations of homology 3-spheres	125
Craig Huneke, A characterization of locally Macaulay completions	131
Takesi Isiwata, Closed ultrafilters and realcompactness	139
Joseph Weston Kitchen, Jr. and David A. Robbins, Tensor products of Banach bundles	151
Allan J. Kroopnick, Note on bounded L^p -solutions of a generalized Liénard equation	171
Ajay Kumar and Ajit Kaur Chilana, Spectral synthesis in products and quotients of hypergroups	177
Charles Livingston, Homology cobordisms of 3-manifolds, knot concordances, and prime knots	193
Hans Opolka, Projective representations of finite groups in cyclotomic fields	207
V. D. Pathak, Isometries of $C^{(n)}[0, 1]$	211
Mark Allan Pinsky, On the spectrum of Cartan-Hadamard manifolds	223
Judith Roitman, The number of automorphisms of an atomic Boolean algebra	231
Kai Wang, Locally smooth torus group actions on integral cohomology complex projective spaces	243