

# Pacific Journal of Mathematics

## **SMALL DOWKER SPACES**

WILLIAM ALBERT RODERICK WEISS

## SMALL DOWKER SPACES

W. WEISS

**We construct a normal, locally compact, first countable, separable, real compact topological space which is not countably paracompact. This construction is performed under the (relatively consistent) set-theoretic hypotheses: Martin's Axiom plus  $\diamond_\omega(E)$ .**

Years ago, C. Dowker proved the now well-known theorem that for any Hausdorff topological space  $X$ , topological product of  $X$  with the closed unit interval is normal iff  $X$  is both normal and countably paracompact [3]. Since then researchers have called Hausdorff spaces which are normal but *not* countably paracompact *Dowker spaces*.

Dowker spaces seem to be extremely rare and difficult to construct. In fact, in the literature there is only one which is constructed using just the usual ZFC axioms of set theory [8]. It was discovered by M. E. Rudin who then asked [10] if there were any "small" Dowker spaces, i.e., ones which are, for example, first countable, separable, realcompact, or of small cardinality.

Examples have been constructed, using extra set-theoretic axioms. M. E. Rudin [9] used the existence of a Suslin line to obtain a Dowker space which is hereditarily separable and first countable. In [5] I. Juhász, K. Kunen and M. E. Rudin construct, using CH, a first countable, hereditarily separable, realcompact Dowker space and claim that with the stronger assumption  $\diamond$  they could construct one which is locally compact as well.

It was unknown if Martin's axiom plus not CH allowed the construction of small Dowker spaces since Martin's axiom plus not CH often implies topological results contrary to CH or  $\diamond$ . In this paper we construct a Dowker space using axioms of set theory consistent with Martin's axiom. This Dowker space is locally compact, first countable, separable and realcompact. By the results of [11] we cannot hope to prove that this Dowker space is hereditarily separable.

Recently, M. Bell has constructed a first countable realcompact Dowker space, assuming Martin's axiom. In fact, only a weakened form of Martin's axiom, called MA ( $\sigma$ -centered) is used. We use it repeatedly in our construction.

MA ( $\sigma$ -centered). If  $P$  is a partial order such that  $P = \bigcup_{n \in \omega} P_n$  where each  $P_n$  is centered and  $\kappa < 2^{\aleph_0}$  and  $\{D_\alpha: \alpha < \kappa\}$  is a collection of dense subsets of  $P$ , then  $P$  contains a  $\{D_\alpha: \alpha < \kappa\}$ -generic subset.

Recall that a subset  $S$  of a partial order  $P$  is centered iff each

finite subset of  $S$  has a lower bound. A subset  $D$  of  $P$  is called dense iff for each  $p \in P$  there is some element of  $D$  less than or equal to  $p$ . If  $\{D_\alpha: \alpha < \kappa\}$  is a collection of dense subset of  $P$ , a subset  $G$  of  $P$  is called  $\{D_\alpha: \alpha < \kappa\}$ -generic iff  $D_\alpha \cap G \neq \emptyset$  for each  $\alpha < \kappa$ .

Another useful axiom is variation of  $\diamond$ , called  $\diamond_{\omega_2}(E)$ . We let  $E = \{\alpha \in \omega_2: \text{cf}(\alpha) = \omega\}$ .

$\diamond_{\omega_2}(E)$ . There is a sequence  $\{X_\alpha: \alpha \in E\}$  such that each  $X_\alpha \subseteq \alpha$  and for any  $X \subseteq \omega_2$   $\{\alpha \in E: X = X \cap \alpha\}$  is stationary in  $\omega_2$ .

**PROPOSITION 1.** *If ZFC is consistent, so is ZFC plus MA ( $\sigma$ -centered) plus  $2^{\aleph_0} = \aleph_2$  plus  $\diamond_{\omega_2}(E)$ .*

*Proof.* This seems to be folklore. The model needed can be the “usual” model of Martin’s axiom plus  $2^{\aleph_0} = \aleph_2$  where the iterated forcing is done over a ground model of  $V = L$ . A modification of the proof of Theorem 7 of Chapter 18 in [2] shows that this model is a model of  $\diamond_{\omega_2}(E)$ . □

The role of  $\aleph_2$  in the collection of axioms in Proposition 1 might be played, in this article, by any regular uncountable cardinal, but for concreteness we choose to assume  $2^{\aleph_0} = \aleph_2$ . These assumptions are used by Hajnal and Juhász for a topological theorem of a completely different nature in [4].

**PROPOSITION 2.**  $\diamond_{\omega_2}(E)$  implies there exists a sequence of ordered pairs  $\{\langle S_\alpha, T_\alpha \rangle: \alpha \in E\}$  such that for all  $\alpha \in E$   $S_\alpha \subseteq \alpha$  and  $T_\alpha \subseteq \alpha$  and for any two subsets  $S$  and  $T$  of  $\omega_2$ ,  $\{\alpha \in E: \text{both } S_\alpha = S \cap \alpha \text{ and } T_\alpha = T \cap \alpha\}$  is stationary in  $\omega_2$ .

*Proof.* This too is folklore. Briefly, let  $f: \omega_2 \rightarrow \omega_2 \times \omega_2$  be any bijection. Let  $S_\alpha = \{\text{first coordinates of elements of } f''X_\alpha\}$  and  $T_\alpha = \{\text{second coordinates of elements of } f''X_\alpha\}$ . Then for any two subsets  $S$  and  $T$  of  $\omega_2$   $\{\alpha \in E: X_\alpha = f^{-1}(S \times T) \cap \alpha\} \cap \{\alpha \in \omega_2: f''\alpha = \alpha \times \alpha\}$  is contained in  $\{\alpha \in E: S_\alpha \times T_\alpha = (S \cap \alpha) \times (T \cap \alpha)\}$  which must therefore be stationary. □

We denote the real numbers by  $\mathbf{R}$  and the rationals by  $\mathbf{Q}$ . In [5], Lusin sets are used to construct a Dowker space. Here, we use:

**PROPOSITION 3.** *MA ( $\sigma$ -centered) plus  $2^{\aleph_0} = \aleph_2$  implies there is an  $L \subseteq \mathbf{R}$  such that*

- (i)  $L \cap \mathbf{Q} = \emptyset$ ,

- (ii) for any open  $U \subseteq R$ ,  $|L \cap U| = \aleph_2$ ,
- (iii) for any nowhere dense  $N \subseteq R$ ,  $|L \cap N| < \aleph_2$ .

*Proof.* Folklore again. Note that the hypotheses imply that  $R$  is not the union of  $\leq \aleph_1$  nowhere dense sets and then make a simple modification to the usual inductive construction of a Lusin set.  $\square$

We now do the promised construction. We will eventually define a topology  $\tau$  on a subset  $X$  of  $L \cup Q$ , where  $L$  is as given in Proposition 3. Fix  $\leq$  as a well ordering of  $L$ . Let  $\{A_\alpha: \alpha \in \omega_2 \setminus E\}$  enumerate  $\{A \subseteq L: |A| < \aleph_2\}$  with each  $A$  being repeated  $\aleph_2$  times. In addition, let  $\{B_\alpha: \alpha \in \omega_2 \setminus E\}$  enumerate all countable sequences of subsets of  $Q$ . Finally, let  $\langle \{S_\alpha, T_\alpha\}: \alpha \in E \rangle$  be as in Proposition 2.

From now on we shall assume ZFC plus MA ( $\sigma$ -centered) plus  $2^{\aleph_0} = \aleph_2$  plus  $\diamond_{\omega_2}(E)$ .

We shall define the topology  $\tau$  on  $X \subseteq L \cup Q$  by recursively defining a sequence of points  $\{x_\alpha: \alpha \in \omega_2\} \subseteq L$  and topologies  $\tau_\beta$  on  $X_\beta = \{x_\alpha: \alpha \in \beta\} \cup Q$ , for all  $\beta \in \omega_2$ . Simultaneously we define a collection  $\{C_\alpha: \alpha \in \omega_2 \setminus E\}$  of subsets of  $L \cup Q$  and a function  $\rho: X \rightarrow \omega$  recursively.

The inductive hypothesis is that for all  $\alpha < \beta$

- (i)  $\tau_\alpha = \tau_\beta \cap \mathcal{P}(X_\alpha)$ ,
- (ii)  $\tau_\beta$  refines the Euclidean topology on  $X_\beta$ ,
- (iii)  $Q$  is dense in  $\langle X_\beta, \tau_\beta \rangle$ ,
- (iv)  $\tau_\beta$  is a locally compact, locally countable topology,
- (v) each  $C_\alpha$  is a clopen subset of  $\langle X_\beta, \tau_\beta \rangle$ ,
- (vi)  $\rho(x_\alpha) = n$  implies there exists  $U \in \tau_{\alpha+1}$  such that for all  $y \in U$ ,  $\rho(y) \leq n$ .
- (vii) for all finite  $F \subseteq \beta$ ,  $Q \setminus \bigcup_{\alpha \in F} C_\alpha$  is dense in the Euclidean topology on  $R$ .

We shall make the construction so that each subset of  $L$  of cardinality  $< \aleph_2$  is contained in some  $C_\alpha$ .

We begin the construction by letting  $X_0 = Q$  and  $\tau_0$  the discrete topology on  $Q$ .

At limit ordinals  $\lambda$ , including  $\lambda = \omega_2$ , we let  $\tau_\lambda$  be the topology on  $X_\lambda = \bigcup \{X_\beta: \beta \in \lambda\}$  generated by  $\bigcup \{\tau_\beta: \beta \in \lambda\}$ . The inductive hypothesis is easily seen to be satisfied. All the action takes place at successor stages and there are two cases to consider: first, when  $\beta = \alpha + 1$  for some  $\alpha \in E$  and second, when  $\beta = \alpha + 1$  for some  $\alpha \notin E$ .

At stage  $\alpha + 1$  when  $\alpha \in E$  we consider  $\langle S_\alpha, T_\alpha \rangle$  and an increasing sequence  $\{\alpha_n: n \in \omega\}$  converging to  $\alpha$ .

*Case (a).* There is  $x \in L \setminus X_\alpha$ , there is a sequence  $\{s_n: n \in \omega\} \subseteq S_\alpha$

and a sequence  $\{t_n: n \in \omega\} \subseteq T_\alpha$  such that:

(i) each  $s_n \in X_\alpha \setminus X_{\alpha_n}$  and each  $t_n \in X_\alpha \setminus X_{\alpha_n}$ ,

(ii) there exists  $l \in \omega$  such that  $\{\rho(s_n): n \in \omega\} \subseteq l$  and  $\{\rho(t_n): n \in \omega\} \subseteq l$ ,

(iii) in the Euclidean topology on  $\mathbf{R}$   $\{S_n\} \rightarrow x$  and  $\{t_n\} \rightarrow x$ .

In this case, let  $x_\alpha = x$ . We need a lemma.

**LEMMA 1.** *There exist open  $U_n \in \tau_\alpha$  containing  $s_n$  such that for all  $\xi < \alpha \{n \in \omega: C_\xi \cap U_n \neq \emptyset\}$  is finite.*

*Proof.* Since  $\langle X_\alpha, \tau_\alpha \rangle$  is first countable, let  $V(n, m)$  be the  $m$ th basic open neighborhood of  $s_n$ .

Let  $P$  be the set of all pairs  $\langle f, g \rangle$  such that

(i)  $f$  is a finite partial function from  $\alpha$  into  $\omega$ ,

(ii)  $g$  is a finite partial function from  $\omega$  into  $\omega$ , and

(iii) for all  $\xi \in \text{dom } f$ , for all  $n \in \text{dom } g$ ,  $n > f(\xi)$  implies

$$s_n \in C_\xi \quad \text{or} \quad V(n, g(n)) \cap C_\xi = \emptyset,$$

with the partial ordering of function extension in both coordinates.

Let  $D_n^1 = \{\langle f, g \rangle: n \in \text{dom } g\}$ . Since each  $C_\xi$  is closed in  $\langle X_\alpha, \tau_\alpha \rangle$ , each  $D_n^1$  is dense in  $P$ .

Let  $D_\xi^2 = \{\langle f, g \rangle: \xi \in \text{dom } f\}$ . Clearly each  $D_\xi^2$  is dense in  $P$ .

Now, if  $\langle f_1, g \rangle$  and  $\langle f_2, g \rangle$  are elements of  $P$  and  $f_1$  and  $f_2$  are compatible as functions, then  $\langle f_1 \cup f_2, g \rangle$  is in  $P$ . Hence for each  $\tilde{g}$ ,  $\{\langle f, g \rangle: g = \tilde{g}\}$  is isomorphic to the partial order of finite partial functions from  $\alpha$  into  $\omega$  ordered by set containment, and so  $P$  is  $\sigma$ -centered.

By MA ( $\sigma$ -centered), there is a  $G \subseteq P$  generic for all  $D_n^1$  and  $D_\xi^2$ . Let  $F = \bigcup \{f: \text{for some } g, \langle f, g \rangle \in G\}$  and  $G = \bigcup \{g: \text{for some } f, \langle f, g \rangle \in G\}$ .

Let  $U_n = V(n, G(n))$ . For each  $\xi \in \alpha$  there is some  $n \in \omega$  such that  $C_\xi \subseteq X_{\alpha_{n\xi}}$ . If  $n > \max \{n_\xi, F(\xi)\}$ , then  $U_n \cap C_\xi = \emptyset$ . □

Returning to the construction, let  $\{O_s(n): n \in \omega\}$  refine  $\{U_n: n \in \omega\}$  of Lemma 1 such that

(i)  $s_n \in O_s(n) \subseteq U_n$ ,  $\rho''O_s(n) \subseteq l$ ,

(ii)  $O_s(n)$  is countable, compact and open in  $\langle X_\alpha, \tau_\alpha \rangle$ ,

(iii) the Euclidean diameter  $\text{diam } O_s(n) < 1/n$  for each  $n$ .

Similarly obtain  $\{O_t(n): n \in \omega\}$  with each  $t_n \in O_t(n)$ . Let  $V_m = \{x_\alpha\} \cup \bigcup \{O_s(n): n \geq m\} \cup \bigcup \{O_t(n): n \geq m\}$ . Let  $\tau_\alpha \cup \{V_m: m \in \omega\}$  generate  $\tau_{\alpha+1}$ . Let  $\rho(x_\alpha) = l$ . A routine check shows that the inductive hypothesis is still satisfied.

*Case (b).* If the conditions for Case (a) are not satisfied, we

waste time with the help of this lemma.

**LEMMA 2.** *There is a  $Q^* \subseteq Q$  which is dense in the Euclidean topology and for all  $\xi < \alpha$ ,  $(Q^* \cap C_\xi)$  is finite.*

*Proof.* Enumerate  $Q$  as  $\{q_n: n \in \omega\}$ . Let  $P$  be the set of all ordered pairs  $\langle M, f \rangle$  such that

- (i)  $M$  is a finite subset of  $Q$ ,
- (ii)  $f$  is a finite partial function from  $\alpha$  into  $\omega$ ,
- (iii) if  $\xi \in \text{dom } f$  then  $C_\xi \cap M \subseteq \{q_n: n < f(\xi)\}$  with the partial order given by extension of each coordinate.

Let  $D_\xi = \{\langle M, f \rangle: \xi \in \text{dom } f\}$ . Clearly  $D_\xi$  is dense in  $P$  for each  $\xi < \alpha$ .

Let, for each pair of rationals  $p, r$  with  $p < r$ ,  $D_{p,r} = \{\langle M, f \rangle: M \cap (p, r) \neq \emptyset\}$ , where  $(p, r)$  is the open interval. Condition (vii) of the inductive hypothesis ensures that each  $D_{p,r}$  is dense.

For each  $\tilde{M}$ ,  $\{\langle M, f \rangle: M = \tilde{M}\}$  is isomorphic to the partial order of finite partial functions from  $\alpha$  into  $\omega$  ordered by set containment, and hence  $P$  is  $\sigma$ -centered.

Hence by MA ( $\sigma$ -centered) there is a  $G \subseteq P$  which is generic with respect to the above-mentioned dense sets. Let  $Q^* = \bigcup \{M: \text{there is some } f \text{ such that } \langle M, f \rangle \in G\}$ , and let  $F = \bigcup \{f: \text{there is some } M \text{ such that } \langle M, f \rangle \in G\}$ . Then  $Q^*$  is dense and for each  $\xi < \alpha$ ,  $(Q^* \cap C_\xi) \subseteq \{q_n: n < F(\xi)\}$ . □

We can now pick any  $x \in L \setminus X_\alpha$  and find  $\{p_n: n \in \omega\} \subseteq Q^*$  from Lemma 2 such that  $\{p_n\} \rightarrow x$  in the Euclidean topology. Let  $x_\alpha = x$  and let  $V_m = \{x_\alpha\} \cup \{p_n: n \geq m\}$ . Let  $\tau_{\alpha+1}$  be generated by  $\tau_\alpha \cup \{V_m: m \in \omega\}$  and let  $\rho(x_\alpha) = 0$  to satisfy the inductive hypothesis in this case.

We can now proceed to stage  $\alpha + 1$  where  $\alpha \notin E$ . We must do two things: define  $x_\alpha$  and  $\tau_{\alpha+1}$  and define  $C_\alpha$ .

Embarking on the first task, let's consider  $B_\alpha = \langle B_{\alpha}^0, B_{\alpha}^1, B_{\alpha}^2, \dots \rangle$ . Using Lemma 2 obtain  $Q^* \subseteq Q$ , dense, such that  $C_\xi \cap Q^*$  is finite for each  $\xi < \alpha$ . There are now two cases to consider.

*Case (a).* Each  $x \in L \setminus X_\alpha$  is in the Euclidean closure of only finitely many of the sets  $B_\alpha^j \cap Q^*$ ,  $j \in \omega$ . In this case let  $x_\alpha$  be any element of  $L \setminus X_\alpha$ . Pick  $\{p_n: n \in \omega\} \subseteq Q^*$  such that  $\{p_n\} \rightarrow x_\alpha$  in the Euclidean topology. Let  $\tau_\alpha \cup \{\{x_\alpha\} \cup \{p_n: n \geq m\}: m \in \omega\}$  generate  $\tau_{\alpha+1}$ . Let  $\rho(x_\alpha) = 1 + \max \{j: x_\alpha \text{ is in the Euclidean closure of } B_\alpha^j \cap Q^*\}$ . Note that the inductive hypothesis is satisfied and that for all  $j \geq \rho(x_\alpha)$ ,  $x_\alpha \notin \tau_{\alpha+1}$ -closure of  $B_\alpha^j$ .

*Case (b).* There is some  $x \in L \setminus X_p$  such that  $x$  is in the Euclidean

closure of infinitely many of the sets  $B_\alpha^j \cap Q^*$ . In this case pick  $x_\alpha$  to be some such  $x$ . Pick an infinite  $J \subseteq \omega$  such that for each  $j \in J$ ,  $x_\alpha$  is in the Euclidean closure of  $B_\alpha^j \cap Q^*$ . For each  $j \in J$  pick  $\{p_n^j: n \in \omega\} \subseteq B_\alpha^j \cap Q^*$  such that  $\{p_n^j\} \rightarrow x_\alpha$  in the Euclidean topology. Let  $\{r_k: k \in \omega\}$  be a subsequence of  $\{p_n^j: n \in \omega, j \in J\}$  such that for each  $j \in J$  each tail of  $\{r_k\}$  contains an infinite subset of  $\{p_n^j: n \in \omega\}$  and such that  $\{r_k\} \rightarrow x_\alpha$  in the Euclidean topology. Let  $\tau_\alpha \cup \{x_\alpha\} \cup \{r_k: k \geq m\}: m \in \omega$  generate  $\tau_{\alpha+1}$ . Let  $\rho(x_\alpha) = 0$ . Note that the inductive hypothesis is satisfied and that  $x_\alpha$  is in the  $\tau_{\alpha+1}$ -closure of infinitely many  $E_\alpha^j$ .

We now proceed to define  $C_\alpha$ . Consider  $A_\alpha$ . If  $A_\alpha \not\subseteq X_{\alpha+1}$ , let  $C_\alpha = \emptyset$ . If  $A_\alpha \subseteq X_{\alpha+1}$ , then  $C_\alpha$  is given by the following lemma.

**LEMMA 3.** *Suppose  $A_\alpha \subseteq (X_{\alpha+1} \cap L)$  and  $\{C_\xi: \xi \in \alpha \setminus E\}$  is a collection of clopen subsets of  $X_{\alpha+1}$  such that for each finite  $F \subseteq (\alpha \setminus E)$ ,  $(Q \cup \{C_\xi: \xi \in F\})$  is dense in the Euclidean topology. Then there exists a clopen  $C_\alpha \subseteq X_{\alpha+1}$  such that  $A_\alpha \subseteq C_\alpha$  and for each finite  $F \subseteq ((\alpha + 1) \setminus E)$ ,  $(Q \cup \{C_\xi: \xi \in F\})$  is dense in the Euclidean topology.*

*Proof.* From Lemma 1 obtain  $Q^*$  such that for each  $\xi \in (\alpha \setminus E)$ ,  $(Q^* \cap C_\xi)$  finite, and  $Q^*$  is dense in the Euclidean topology. It now suffices to construct a clopen  $C_\alpha$  such that  $A_\alpha \subseteq C_\alpha \subseteq X_{\alpha+1}$  and  $(Q^* \setminus C_\alpha)$  is dense.

To this end let  $P$  be the set of all triples  $\langle a, b, c \rangle$  such that:

- (i)  $a$  and  $b$  are clopen, compact subsets of  $X_{\alpha+1}$ ,
- (ii)  $c$  is a finite subset of  $Q^*$ ,
- (iii)  $(a \cup b) \cap c = \emptyset$ ,
- (iv)  $a \cap c = \emptyset$

with the partial order of set containment in each coordinate.

$\{\langle a, b, c \rangle: x \in a \cup b\}$  is dense for each  $x \in X_{\alpha+1}$ . For each interval  $(p, q)$  with rational endpoints,  $\{\langle a, b, c \rangle: (p, q) \cap c \neq \emptyset\}$  is dense.

For each finite union  $I$  of interval with rational endpoints and each finite  $\tilde{c} \subseteq Q^*$ ,  $\{\langle a, b, c \rangle: a \subseteq I, (b \cap I) = \emptyset \text{ and } c = \tilde{c}\}$  is centered; hence  $P$  is  $\sigma$ -centered.

Let  $G \subseteq P$  be generic with respect to the above dense sets. Let  $C_\alpha = \bigcup \{a: \text{for some } b \text{ and } c, \langle a, b, c \rangle \in G\}$ . It is straightforward to show that  $C_\alpha$  is clopen,  $A_\alpha \subseteq C_\alpha$  and  $(Q^* \setminus C_\alpha)$  is dense.  $\square$

This completes the construction. We let  $X = X_{\omega_2}$  and  $\tau = \tau_{\omega_2}$ . Clearly  $\tau$  is a locally compact,  $T_2$ , first countable, separable and submetrizable topology on  $X$ . Since  $\langle X, \tau \rangle$  is first countable and submetrizable, it is real-compact [5] also. It remains to show that  $\langle X, \tau \rangle$  is a Dowker space.

We show that  $\langle X, \tau \rangle$  is not countably compact by showing that the open cover  $\{O_j: j \in \omega\}$  has no precise locally finite refinement,

where  $O_j = \{x \in X: \rho(x) \leq j\}$ . Due to (vi) of the inductive hypothesis  $\{O_j: j \in \omega\}$  is an open cover. In order to achieve a contradiction assume  $\{W_j: j \in \omega\}$  is a locally finite precise refinement. For some  $\alpha \in (\omega_2 \setminus E)$ ,  $\langle (W_0 \cap Q), (W_1 \cap Q), (W_2 \cap Q), \dots \rangle = B_\alpha$  and we look at what happened at stage  $\alpha + 1$ . Under Case (b)  $x_\alpha$  witnesses that  $\{W_j: j \in \omega\}$  is not locally finite. Under Case (a)  $x_\alpha$  witnesses that  $\{W_j: j \in \omega\}$  is not a cover, since if  $j < \rho(x_\alpha)$  then  $x_\alpha \notin O_j$  and if  $j \geq \rho(x_\alpha)$  then  $x_\alpha \notin \overline{W_j \cap Q} \supseteq W_j$ .

In order to show that  $\langle X, \tau \rangle$  is normal, we need a short lemma.

LEMMA 4. *If  $H$  and  $K$  are closed disjoint subsets of  $X$ , one of which has cardinality  $< \aleph_2$ , then there are disjoint open sets separating  $H$  and  $K$ .*

*Proof.* Assume  $|H| < \aleph_2$  so that there is some clopen  $C \subseteq X$  such that  $|C| < \aleph_2$  and  $H \subseteq C$ . Let  $K^* = K \cap C$ . By MA ( $\sigma$ -centered), since  $X$  is locally compact and submetrizable there exist disjoint open  $U$  and  $V$  such that  $H \subseteq U$  and  $K^* \subseteq V$  [1, 6]. Then  $H \subseteq U \cap C$  and  $K \subseteq U \cup (X \setminus C)$  separating  $H$  and  $K$ .  $\square$

Now let  $H$  and  $K$  be arbitrary closed disjoint subsets of  $X$ . Without loss of generality we assume  $H \cup K \subseteq L$  and we find disjoint open sets separating  $H$  and  $K$ . Let  $H_i = \{x \in H: \rho(x) \leq i\}$  and  $K_i = \{x \in K: \rho(x) \leq i\}$ . Define  $U_i^H$  to be the compliment in  $R$  of the Euclidean closure of  $\bigcup \{V: V \text{ is Euclidean open and } |V \cap H_i| < \aleph_2\}$ . Note that for each nonempty Euclidean open  $V \subseteq U_i^H |V \cap H_i| = \aleph_2$ . Similarly define  $U_i^K$  for each  $i \in \omega$ .

LEMMA 5. *With  $U_i^H$  and  $U_i^K$  defined as above for each  $i \in \omega$  we have  $\bigcup_{i \in \omega} U_i^K \cap \bigcup_{i \in \omega} U_i^H = \emptyset$ .*

*Proof.* Suppose  $U_i^H \cap U_j^K \neq \emptyset$ . Then there is some interval  $I \subseteq U_i^H \cap U_j^K$ . Let  $\hat{H} = H_i \cap I$  and  $\hat{K} = K_j \cap I$ . Let  $Y_H = \{\alpha \in \omega_2: \text{for all } \beta < \alpha \hat{H} \cap [\beta, \alpha) \text{ is dense in } I\}$  and  $Y_K = \{\alpha \in \omega_2: \text{for all } \beta < \alpha \hat{K} \cap [\beta, \alpha) \text{ is dense in } I\}$ .  $Y_H$  and  $Y_K$  are both cub subsets of  $\omega_2$ . So there exists, by  $\diamond_{\omega_2}(E)$ , some  $\alpha \in E \cap Y_H \cap Y_K$  such that  $\hat{H} \cap \alpha = S_\alpha$  and  $\hat{K} \cap \alpha = T_\alpha$ .

Let's see what happened at stage  $\alpha + 1$ . We had  $\{\alpha_n: n \in \omega\}$  increasing up to  $\alpha$ . Since  $\alpha \in Y_H \cap Y_K$  we can find  $x \in L \setminus X_\alpha$  and sequences  $\{s_n: n \in \omega\}$  and  $\{t_n: n \in \omega\}$  such that the conditions for Case (a) hold. Hence  $x_\alpha \in \bar{S}_\alpha \cap \bar{T}_\alpha \subseteq \bar{H} \cap \bar{K}$  which is a contradiction.  $\square$

Now, since  $L$  has the properties from Proposition 3, both  $H \setminus \bigcup_{i \in \omega} U_i^H$  and  $K \setminus \bigcup_{i \in \omega} U_i^K$  have cardinality less than  $\aleph_2$ . Hence by Lemma 4 there are open  $V_H$  and  $V_K$  such that:



$$(H \setminus \bigcup_{i \in \omega} U_i^H) \subseteq V_H \subseteq \bar{V}_H \subseteq (X \setminus K)$$

and

$$(K \setminus \bigcup_{i \in \omega} U_i^K) \subseteq V_K \subseteq \bar{V}_K \subseteq (X \setminus H).$$

Hence

$$H \subseteq (V_H \setminus \bar{V}_K) \cup \bigcup_{i \in \omega} U_i^H$$

and

$$K \subseteq (V_K \setminus \bar{V}_H) \cup \bigcup_{i \in \omega} U_i^K$$

which separates  $H$  and  $K$ . The proof is complete.

We could do the above constructions and proofs with  $\aleph_2$  replaced everywhere by  $\aleph_1$  thus reducing the set theoretic assumptions to ZFC plus  $\diamond$ . A Dowker space with properties similar to  $X$  was claimed in [5] to follow from  $\diamond$ . The additional property of hereditary separability mentioned in [5] can be obtained by weaving in the idea of constructing hereditarily separable spaces from  $\diamond$  in [7] to the above construction.

The space  $\langle X, \tau \rangle$  above is normal and each bounded or closed unbounded (referring to the indices) subspace is also normal. However, there is no reason to believe that  $\langle X, \tau \rangle$  is hereditarily normal.

#### REFERENCES

1. K. Alster and T. Przymusiński, *Normality and Martin's axiom*, *Fund. Math.*, **41** (1976), 123-131.
2. K. J. Devlin, *Aspects of Constructibility*, *Lecture Notes in Math.*, Vol. **354**, Springer-Verlag, Berlin and New York, 1973.
3. C. H. Dowker, *On countably paracompact spaces*, *Canad. J. Math.*, **3** (1951), 219-224.
4. A. Hajnal and I. Juhász, *Weakly separated subspaces and networks*, preprint.
5. I. Juhász, K. Kunen and M. E. Rudin, *Two more hereditarily separable non-Lindelöf spaces*, *Canad. J. Math.*, **28** (1976), 998-1005.
6. I. Juhász and W. Weiss, *Martin's axiom and normality*, *General Topology and Appl.*, **9** (1978), 263-274.
7. A. Ostaszewski, *On countably compact, perfectly normal spaces*, *J. London Math. Soc.*, **14** (1976), 505-516.
8. M. E. Rudin, *A normal space for which  $X \times I$  is not normal*, *Fund. Math.*, **78** (1971-1972), 179-168.
9. ———, *A separable Dowker space*, *Symposia Mathematica*, Instituto Nazionale di Alta Matematica, 1973.
10. ———, *Lectures in Set-theoretic Topology*, *Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics No.*, **23** (1974); Amer. Math. Soc., Providence, R.I.
11. Z. Szentmiklóssy, *S and L spaces under MA*, to appear.

Received October 9, 1979 and in revised form December 14, 1979. Research partially supported by NSTRC grant no. A3185.

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California  
Los Angeles, CA 90024

HUGO ROSSI

University of Utah  
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University  
Stanford, CA 94305

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$102.00 a year (6 Vols., 12 issues). Special rate: \$51.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1981 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

Thomas E. Armstrong and William David Sudderth, Nearly strategic measures .....	251
John J. Buoni, Artatrana Dash and Bhushan L. Wadhwa, Joint Browder spectrum .....	259
Jack Paul Diamond, Hypergeometric series with a $p$ -adic variable .....	265
Raymond Frank Dickman, Jack Ray Porter and Leonard Rubin, Completely regular absolutes and projective objects .....	277
James Kenneth Finch, On the local spectrum and the adjoint .....	297
Benno Fuchssteiner, An abstract disintegration theorem .....	303
Leon Gerber, The volume cut off a simplex by a half-space .....	311
Irving Leonard Glicksberg, An application of Wermer's subharmonicity theorem .....	315
William Goldman, Two examples of affine manifolds .....	327
Yukio Hirashita, On the Weierstrass points on open Riemann surfaces .....	331
Darrell Conley Kent, A note on regular Cauchy spaces .....	333
Abel Klein and Lawrence J. Landau, Periodic Gaussian Osterwalder-Schrader positive processes and the two-sided Markov property on the circle .....	341
Brenda MacGibbon, $\mathcal{H}$ -Borelian embeddings and images of Hausdorff spaces .....	369
John R. Myers, Homology 3-spheres which admit no PL involutions .....	379
Boon-Hua Ong, Invariant subspace lattices for a class of operators .....	385
Chull Park, Representations of Gaussian processes by Wiener processes .....	407
Lesley Millman Sibner and Robert Jules Sibner, A sub-elliptic estimate for a class of invariantly defined elliptic systems .....	417
Justin R. Smith, Complements of codimension-two submanifolds. III. Cobordism theory .....	423
William Albert Roderick Weiss, Small Dowker spaces .....	485
David J. Winter, Cartan subalgebras of a Lie algebra and its ideals. II .....	493