

Pacific Journal of Mathematics

CONVEXITY AND THE TABLE THEOREM

MARK D. MEYERSON

CONVEXITY AND THE TABLE THEOREM

MARK D. MEYERSON

In "The Table Theorem" ([1]), Roger Fenn proves that one can "balance" a square "table" on any "hill". One of the hypotheses he suggests relaxing is that the hill has convex support. We show here the necessity of such a hypothesis.

More precisely, Fenn showed that if $D \subset \mathbf{R}^2$ is a compact, convex disk and $f: \mathbf{R}^2 \rightarrow \mathbf{R}$ is a map which is zero outside D and nonnegative inside D , then, given $s > 0$, there is a square of side s in \mathbf{R}^2 with center in D such that f takes the same values at all the vertices of the square. Below, we shall give an example where D is not convex and the conclusion fails to hold.

The two following lemmas will be useful:

LEMMA 1. *Suppose $K \subset \mathbf{R}^2$ is compact and $\varepsilon > 0$ is given. Then there is a $\delta > 0$ such that if all the points of K' are within δ of K , then any square with vertices in K' has each of its vertices within ε of a point of K such that these 4 points of K form the vertices of a square. Four copies of a single point are viewed as the vertices of a (degenerate) square.*

Proof. Each square in \mathbf{R}^2 corresponds to a point in \mathbf{R}^8 . The square with vertices (x_1, y_1) , (x_2, y_2) , (x_3, y_3) , and (x_4, y_4) , traveling counterclockwise around the square, is given by equations: $x_3 = x_2 + y_1 - y_2$, $y_3 = y_2 + x_2 - x_1$, $x_4 = x_1 + y_1 - y_2$, and $y_4 = y_1 + x_2 - x_1$. Thus all squares are represented by the 4-dimensional flat, F , in \mathbf{R}^8 which is the intersection of these 4 hyperplanes. Also represented are degenerate squares.

Let $N \subset \mathbf{R}^8$ be the set of points within $\varepsilon/2$ of $K^4 \cap F$ ($K^4 = K \times K \times K \times K$). Let $\delta > 0$ be smaller than $\varepsilon/4$ and smaller than half the distance from F to $K^4 \setminus N$. Let V' be a square with vertices within δ of K . Then in \mathbf{R}^8 , V' is represented by a point $v \in F$ within $2\delta < \varepsilon/2$ of a point k of K^4 . Now v is more than 2δ from $K^4 \setminus N$ so $k \in K^4 \cap N$. Thus k is within $\varepsilon/2$ of $K^4 \cap F$. Hence v is within ε of $K^4 \cap F$. \square

The next lemma follows from the fact that an angle greater than 90° with vertex at the center of a square must contain a vertex of the square.

LEMMA 2. *Suppose at every point of region D there is the vertex of an angle greater than 90° with an ε -radius sector in the interior of D (except perhaps for the vertex of the angle). Then for $\delta < \sqrt{2}\varepsilon$, if a square of side δ has center in D , at least one vertex of the square is in the interior of D .* \square

We now construct the promised example.

EXAMPLE. A compact disk $D \subset \mathbf{R}^2$, a map $f: \mathbf{R}^2 \rightarrow \mathbf{R}$ zero outside D and nonnegative inside D , and $s > 0$, such that if f takes the same values on the vertices of a square of side s then the square's center is outside D .

Construction. Start with the unit circle in the plane. For $\varepsilon > 0$ small add the two segments from $(-\varepsilon, 0)$ to the circle which make angles of $\pm\varepsilon$ with the positive x -axis. Throw out the small arc of the circle subtended by the angle the two segments determine. "Round off" the two corners on the unit circle with arcs of small circles so that the resulting curve, C :

- (i) is strongly starlike from any point $(-\gamma, 0)$ with $\varepsilon < \gamma < 1$ and
- (ii) the points from but not including $(-\varepsilon, 0)$ to the origin lie outside C .

Note that the above can be achieved so that the only squares with all 4 vertices on C have center at the origin and vertices on the unit circle. One way to check this is to start with the unit circle and a single radius. On this compact set the claim is true. Then, using Lemma 1, we only need check for new squares which are very close to the old ones or very small.

Next, using Lemma 2, choose a small $s > 0$ so that no square of side s can have its center on or inside C and all its vertices on or outside C .

The boundary of $D(\partial D)$ is C translated to the right by γ , $\varepsilon < \gamma < 1$ (to be determined later). For $0 \leq r \leq 1$, the level curve on which f takes value r is given by multiplying the points of ∂D (treating them as vectors) by $1 - r$. So the level curves are all similar to C . By condition (i) above, this will indeed give us a continuous function. Now the only nonzero level on which our "table can balance" has $1 - r$ (the radius of the circle about the level curve) equal to $s/\sqrt{2}$, i.e., $r = 1 - s/\sqrt{2}$. For this r we choose γ so that $\varepsilon < \gamma < \varepsilon/r$ and hence $(1 - r)\gamma > \gamma - \varepsilon$. So the only center of a square "balancing" at a nonzero level lies over a point outside D (at $((1 - r)\gamma, 0)$).

Thus the only “tables which balance” on the graph of f are at the zero level or $r = 1 - s/\sqrt{2}$ level, and in both cases the centers lie outside D . \square

By putting the level curves at heights $g(r)$ for appropriate g , rather than r , we could have permitted a smoothing out of C at $(-\varepsilon, 0)$. In fact we could have made ∂D and f as smooth as desired.

REFERENCE

1. Roger Fenn, *The Table Theorem*, Bull. London Math. Soc., **2** (1970), 73-76.

Received September 3, 1980.

DEPARTMENT OF MATHEMATICS
U. S. NAVAL ACADEMY
ANNAPOLIS, MD 21402

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Charles A. Asmuth and Joe Repka, Tensor products for $SL_2(\mathbb{F})$. II. Supercuspidal representations	1
Joseph Barback, On finite sums of regressive isols	19
Matthew G. Brin and Daniel Russell McMillan, Jr., Generalized three-manifolds with zero-dimensional nonmanifold set	29
Kun Soo Chang, Converse measurability theorems for Yeh-Wiener space	59
Christopher Brian Croke, A "maximal torus" type theorem for complete Riemannian manifolds	65
Gustave Adam Efroymson, Sums of squares in planar Nash rings	75
John Robert Fisher, Axiomatic radical and semisimple classes of rings	81
Betty Kvarda, Consecutive integers for which $n^2 + 1$ is composite	93
Roosevelt Gentry, New diagram proofs of the Hausdorff-Young theorem and Young's inequality	97
Patrick M. Gilmer, Topological proof of the G -signature theorem for G finite	105
Chung Wei Ha, A noncompact minimax theorem	115
James J. Hebda, Manifolds admitting taut hyperspheres	119
Takayuki Kawada, Sample functions of Pólya processes	125
Peter K. F. Kuhfittig, Common fixed points of nonexpansive mappings by iteration	137
James Thomas Loats and Judith Roitman, Almost rigid Hopfian and dual Hopfian atomic Boolean algebras	141
Roger McCann, On embedding semiflows into a radial flow on I_2	151
John McDonald, Closed orbits of convex sets of operators on the disk algebra	159
Mark D. Meyerson, Convexity and the table theorem	167
Arnold William Miller, Generic Souslin sets	171
Takemi Mizokami, On the closed images of paracomplexes	183
Jagannadham Venkata Pakala and Thomas Stephen Shores, On compactly packed rings	197
Andrew Pletch, Strong completeness in profinite groups	203
Wilbur Carrington Whitten, Inverting double knots	209
James Juei-Chin Yeh, Existence of strong solutions for stochastic differential equations in the plane	217