

Pacific Journal of Mathematics

SEMIFREE FINITE GROUP ACTIONS ON HOMOTOPY SPHERES

RONALD DOTZEL

SEMIFREE FINITE GROUP ACTIONS ON HOMOTOPY SPHERES

RONALD M. DOTZEL

Suppose G is a finite group acting semifreely (i.e., free off the fixed set) on a finite CW complex X in the homotopy type of S^n . When X^G is also homotopy equivalent to S^n (as e.g., in $S^n \times D^k$) necessary and sufficient conditions are given to determine the degree of the inclusion $X^G \rightarrow X$. It follows that for instance, if G is the group of quaternions (nonabelian of order 8), the only attainable degrees are those $\pm 1 \pmod{8}$.

O. Introduction. Suppose that G is a finite group acting cellularly and semifreely (i.e., free off the fixed set) on a finite CW-complex X , in the homotopy type of S^n . Assume that X^G is also homotopy equivalent to S^n . The degree q of the inclusion $X^G \hookrightarrow X$ is defined and by Smith Theory is relatively prime to $|G|$ (see [1; Chap. III]). We give necessary and sufficient conditions for the converse to be true. First recall that when q is relatively prime to $|G|$, the trivial $\mathbb{Z}G$ -module \mathbb{Z}_q has projective dimension one and well-defines an element, $[\mathbb{Z}_q]$, in $\tilde{K}_0(\mathbb{Z}G)$, the reduced projective class group.

THEOREM. *Let G be a finite group acting cellularly and semifreely on a finite CW complex X . Assume that both X and X^G are in the homotopy type of S^n . Let q be the degree of $X^G \hookrightarrow X$. Then $[\mathbb{Z}_q] = 0$ in $\tilde{K}_0(\mathbb{Z}G)$. Conversely, suppose q is relatively prime to $|G|$ and $[\mathbb{Z}_q] = 0$ in $\tilde{K}_0(\mathbb{Z}G)$. Then there exists an action of G , as above, in which the degree of $X^G \hookrightarrow X$ is q .*

REMARKS. (i) From [2] it follows, e.g., that the quaternions, a nonabelian group of order 8, cannot act semifreely as above with degree of the inclusion $\pm 3 \pmod{8}$.

(ii) When G is cyclic and q is prime to $|G|$, $[\mathbb{Z}_q]$ is always zero [4; §6] and therefore it is easy to see how to construct an effective, unrestricted action of the quaternions realizing a degree 3 inclusion. In [1; page 391], Bredon gave examples of semifree (smooth) cyclic group actions realizing any (appropriate) degree. In general, of course, one cannot hope to find smooth semifree actions of this type.

(iii) The invariant introduced here is in fact the same as the invariant, $\chi(f)$, introduced in [3] (when applied to the inclusion map).

1. **Proof of the theorem.** Suppose G is a finite group acting as described and let q denote the degree of $X^G \rightarrow X$. X/X^G has semifree G action with one fixed point, is $n - 1$ connected and has $H_n(X/X^G) = \mathbf{Z}_q$. By an elementary construction (a nonequivariant version can be found in [5; Prop. 6.13]) one can replace X/X^G with a finite CW -complex Y , with semifree G -action, contractible fixed set, no m cells for $0 < m \leq n - 1$ and such that X/X^G is an equivariant strong deformation retract of Y . Let k denote the cellular dimension of Y . Then

$$0 \longrightarrow C_k(Y, Y^G) \longrightarrow \cdots \longrightarrow C_{n+1}(Y, Y^G) \longrightarrow C_n(Y, Y^G) \longrightarrow \mathbf{Z}_q$$

is a finite free resolution of \mathbf{Z}_q over $\mathbf{Z}G$. This implies that $[\mathbf{Z}_q] = 0$ in $\tilde{K}_0(\mathbf{Z}G)$.

Now suppose q is relatively prime to $|G|$ and that $[\mathbf{Z}_q] = 0$ in $\tilde{K}_0(\mathbf{Z}G)$. Let M_f be the mapping cylinder of a map $f: S^n \rightarrow S^n$ of degree q ; $M_f = S^n \times I \cup S^n/(x, 0) \sim f(x)$. Denote by Y the semifree G -complex obtained by taking the disjoint union of $|G|$ copies of M_f , identified along the sphere $S^n \times \{1\}$. G acts on Y via the action induced from the free action on the disjoint union; the fixed set is S^n .

It follows from Mayer-Vietoris that if I_ε denotes the augmentation ideal, i.e., the kernel of $\mathbf{Z}G \xrightarrow{\varepsilon} \mathbf{Z}$ (where $\varepsilon(\Sigma n_i g_i) = \Sigma n_i$), then $H_n(Y) = \mathbf{Z}G/qI_\varepsilon$.

There is the exact sequence of $\mathbf{Z}G$ -modules,

$$0 \longrightarrow I_\varepsilon/qI_\varepsilon \longrightarrow H_n(Y) \longrightarrow \mathbf{Z}.$$

Since $I_\varepsilon/qI_\varepsilon$ has projective dimension one, it well-defines an element $[I_\varepsilon/qI_\varepsilon]$ in $\tilde{K}_0(\mathbf{Z}G)$.

Now consider the exact sequence of $\mathbf{Z}G$ -modules (\mathbf{Z}_q has trivial G -action),

$$0 \longrightarrow \ker \varepsilon_q \longrightarrow \mathbf{Z}G \xrightarrow{\varepsilon_q} \mathbf{Z}_q$$

where ε_q is ε followed by reduction mod q). Since \mathbf{Z}_q has projective dimension one, it well-defines an element in $\tilde{K}_0(\mathbf{Z}G)$ which is $-[\ker \varepsilon_q]$. $\ker \varepsilon_q$ and $I_\varepsilon/qI_\varepsilon$ are related by the exact sequence

$$0 \longrightarrow \mathbf{Z}G \xrightarrow{q} \ker \varepsilon_q \longrightarrow I_\varepsilon/qI_\varepsilon.$$

The assumption that $[\mathbf{Z}_q] = 0$ forces $[I_\varepsilon/qI_\varepsilon] = 0$ and so $I_\varepsilon/qI_\varepsilon$ has a short exact projective resolution

$$0 \longrightarrow \tilde{F} \longrightarrow F_0 \longrightarrow I_\varepsilon/qI_\varepsilon$$

where F_0 is free and \tilde{F} is stably free. Adding free orbits of $n + 1$

cells to Y to kill $I_\varepsilon/qI_\varepsilon$ (realizing $F_0 \twoheadrightarrow I_\varepsilon/qI_\varepsilon$) produces \tilde{Y} with $H_{n+1}(\tilde{Y}) = \tilde{F}$, $H_n(\tilde{Y}) = \mathbf{Z}$. Using the surjective Hurewicz homomorphism (\tilde{Y} is $n - 1$ connected), $n + 2$ cells may be added ($H_{n+1}(\tilde{Y})$ is made free by wedging, at a fixed point, with a suitable bouquet of spheres) to produce the complex X which is homotopy equivalent to S^n . By considering the homology sequence of the triple (X, \tilde{Y}, Y^G) one can see that the degree $X^G \hookrightarrow X$ is q . \square

As in [4; §6], let (q, N) denote the ideal of $\mathbf{Z}G$ generated by q (relatively prime to $|G|$) and $N = \sum_{g \in G} g$. It is easy to see that $[\mathbf{Z}_q] = [(q, N)]$ in $\tilde{K}_0(\mathbf{Z}G)$. This explains Remark (i).

REFERENCES

1. G. Bredon, *Introduction to Compact Transformation Groups*, Academic Press, 1972.
2. J. Martinet, *Modules Sur L'Algèbre Du Groupe Quaternionen*, Ann. Scient. Ec. Norm. Sup., 4^o serie, t. 4, 1971, p. 309 à 408.
3. T. Petrie, *G maps and the projective class group*, Comment. Math. Helveticci, **39** (51), 611-626.
4. R. Swan, *Periodic resolutions for finite groups*, Annals of Math., **72** (1960), 267-291.
5. R. Switzer, *Algebraic Topology—Homotopy and Homology*, Springer-Verlag, 1975.

Received September 26, 1980.

THE UNIVERSITY OF TEXAS
AUSTIN, TX 78712

Current Address:

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MISSOURI-ST. LOUIS
ST. LOUIS, MISSOURI 63121

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR AGUS

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

R. ARNES

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Abdul Aziz , <i>On the zeros of composite polynomials</i>	1
Salomon Benzaquen and Enrique M. Cabaña , <i>The expected measure of the level sets of a regular stationary Gaussian process</i>	9
Claudio D'Antoni, Roberto Longo and László Zsidó , <i>A spectral mapping theorem for locally compact groups of operators</i>	17
Ronald Dotzel , <i>Semifree finite group actions on homotopy spheres</i>	25
Daniel H. Gottlieb , <i>The Lefschetz number and Borsuk-Ulam theorems</i>	29
Shui-Hung Hou , <i>On property (Q) and other semicontinuity properties of multifunctions</i>	39
Kevin Mor McCrimmon , <i>Compatible Peirce decompositions of Jordan triple systems</i>	57
Mitsuru Nakai , <i>Corona problem for Riemann surfaces of Parreau-Widom type</i>	103
Jack Ray Porter and R. Grant Woods , <i>Extensions of Hausdorff spaces</i>	111
Milton Rosenberg , <i>Quasi-isometric dilations of operator-valued measures and Grothendieck's inequality</i>	135
Joseph L. Taylor , <i>A bigger Brauer group</i>	163
Thomas Vogel , <i>Symmetric unbounded liquid bridges</i>	205
Steve Wright , <i>The splitting of operator algebras. II</i>	243