

Pacific Journal of Mathematics

NOTE ON EXPONENTIAL POLYNOMIALS

LÁSZLÓ SZÉKELYHIDI

NOTE ON EXPONENTIAL POLYNOMIALS

LÁSZLÓ SZÉKELYHIDI

It is known that every finite dimensional translation invariant subspace of measurable functions on a σ -compact locally compact Abelian group consists of exponential polynomials. This paper extends this result for continuous functions on arbitrary commutative topological groups. An analogous characterization is proved for trigonometric polynomials using Fourier transformation.

In this paper joining with the investigations of Engert [3] and Laird [5] we prove that every finite dimensional translation invariant subspace of continuous functions on arbitrary commutative topological groups consists of exponential polynomials. Our method is similar to that of [3] but we prove the important lemma of Engert in a simpler way using generalized polynomials. In contrast with [3] and [5] here the main emphasis is on functional equations. In the last part we prove an analogous result for bounded continuous functions using the Fourier transform of almost periodic functions. We note that translation invariant finite dimensional subspaces of the space of finite signed measures on a commutative topological group can be characterized in a similar way by the same technique.

If G is an Abelian group then an additive function on G is a complex valued function a such that $a(x + y) = a(x) + a(y)$ for all x and y in G . A multiplicative function on G is a complex valued function m such that $m(x + y) = m(x)m(y)$ for all x and y in G . If n is a positive integer then we mean by an n -additive function on G a complex valued function on G^n which is additive in each variable. We define generalized polynomials on G as functions satisfying the so called Fréchet equation: $\Delta_y^{n+1}f(x) = 0$. (Here Δ_y denotes the difference operator: $\Delta_y f(x) = f(x + y) - f(x)$ and $\Delta_y^{n+1}f = \Delta_y(\Delta_y^n f)$.) Functions with this property are called generalized polynomials of degree at most n . It is well-known (see e.g., [2], [6], [8]) that every complex valued generalized polynomial of degree at most n can be uniquely expressed in the form $\sum_{k=0}^n A^{(k)}$ where $A^{(k)}$ is the diagonalization of a k -additive, symmetric function A_k , that is $A^{(k)}(x) = A_k(x, x, \dots, x)$ ($A^{(0)}$ is a constant). For more about generalized polynomials on groups see [6], [8].

If G is a topological Abelian group then by a polynomial on G we mean a function of the form $p(x) = P(a_1(x), \dots, a_n(x))$ where P is a complex polynomial in n variables and a_i ($i = 1, \dots, n$) is a

continuous additive function. An exponential polynomial on G is a function of the form $\sum_{i=1}^n p_i \cdot m_i$ where p_i is a polynomial and m_i is a continuous multiplicative function. By a trigonometric polynomial on G we mean a linear combination of characters, that is continuous multiplicative functions into the complex unit circle.

A multi-index $p = (p_1, \dots, p_n)$ is an n -tuple of nonnegative integers and if (a_1, \dots, a_n) is a complex n -tuple, then a^p is defined to be $a_1^{p_1} \dots a_n^{p_n}$. (For more details on the notation see [3], [4], [5].)

THEOREM 1. *Let f be a continuous function on the topological group G such that the complex linear space spanned by $\{\Delta_y f: y \in G\}$ is a finite dimensional space of polynomials. Then f is a polynomial.*

Proof. Let a_1, \dots, a_n be a finite set of continuous additive functions such that all polynomials in the subspace V spanned by $\{\Delta_y f: y \in G\}$ are built up from these functions and $\{a^p\}$ are linearly independent for $|p| \leq N$. Then

$$\Delta_y f = \sum_{|p| \leq N} (P/y) a^p$$

holds for all y in G . We see that f satisfies the Fréchet equation $\Delta_y^{N+2} f = 0$ and hence we have the representation

$$f = \sum_{k=0}^{N+1} A^{(k)} .$$

This yields

$$\Delta_y f(x) = \sum_{k=0}^{N+1} [A^{(k)}(x+y) - A^{(k)}(x)] .$$

On the right hand side we have only one member which is of degree N in x . This is $A_{N+1}(x, x, \dots, x, y)$. It follows that

$$\sum_{|p|=N} (p/y) a^p = A_{N+1}(x, x, \dots, x, y)$$

holds for all x and y in G . Since the right hand side is additive in y we have

$$\sum_{|p|=N} [c_p(y+z) - c_p(y) - c_p(z)] a^p = 0 .$$

Here the functions a^p ($|p| = N$) are linearly independent and we conclude that c_p is additive for all $|p| = N$. Hence $A^{(N+1)}$ is a polynomial. Repeating this argument for the function $f - A^{(N+1)}$ we get the statement by induction.

THEOREM 2. *Let V be a translation invariant finite dimen-*

sional vectorspace of continuous functions on a topological Abelian group. Then every function in V is an exponential polynomial.

Proof. Let g_1, \dots, g_n be a basis for V , then for every function f in V the functional equation

$$f(x + y) = \sum_{i=1}^n g_i(x)h_i(y)$$

holds. Let x_1, \dots, x_n be elements of the group G for which the matrix $(g_i(x_j))$ is regular. We may suppose without loss of generality that this matrix is the identity matrix. We introduce the notations $\hat{f}(x) = (f(x_1 + x), \dots, f(x_n + x))$, $M(x) = (g_i(x_j + x))$, $\tilde{h}(x) = (h_1(x), \dots, h_n(x))$ for all x in G . Then we have the functional equation

$$\tilde{f}(x + y) = M(x)\tilde{h}(y).$$

which shows that $\tilde{f} = \tilde{h}$ and the subspace in C^n generated by the range of \tilde{f} is invariant under $M(x)$ for all x . (C denotes the set of complex numbers.) We may suppose that this subspace is C^n , then we have for all x and y in G

$$M(x + y) = M(x)M(y).$$

The matrices $M(x)$ commute for all x hence they can be transformed into triangular form simultaneously. Since after a similarity transformation our equation remains valid we may suppose that the matrices are all triangular from above. It is easy to see that the diagonal elements are all multiplicative functions. Let $D(x)$ be the diagonal matrix for which $M(x) - D(x)$ is strictly triangular from above for all x . Then $D(x + y) = D(x)D(y)$ and with the notation $A(x) = D^{-1}(x)M(x)$ we have

$$A(x + y) = A(x)A(y)$$

and all diagonal elements of $A(x)$ are 1. This equation means for the components A_{ij} of A

$$A_{ij}(x + y) = \sum_{k=1}^n A_{ik}(x)A_{kj}(y) = \sum_{k=i+1}^{j-1} A_{ik}(x)A_{kj}(y) + A_{ij}(x) + A_{ij}(y).$$

We prove by induction on $j - i$ that A_{ij} is a polynomial. For $i = j$ this is trivial. Supposing that it is valid for $j - i \leq l$ we see that

$$\Delta_y A_{i, i+l+1}(x) = A_{i, i+l+1}(y) + \sum_{k=i+1}^{i+l} A_{ik}(x)A_{k, i+l+1}(y)$$

hence the subspace spanned by $\{\Delta_y A_{i, i+l+1}, y \in G\}$ is contained in the

subspace spanned by $1, A_{ik}$ ($k = i + 1, \dots, i + l$) (l), that is, consists of polynomials. Thus by Theorem 1 all components of M are exponential polynomials. Finally, by

$$\tilde{f}(x) = M(x)\tilde{f}(0)$$

we conclude that f is an exponential polynomial.

THEOREM 3. *Let V be a translation invariant finite dimensional vectorspace of continuous bounded functions on a topological Abelian group which has sufficiently many characters. Then every function in V is a trigonometric polynomial.*

Proof. Using the notations of the previous theorem we have that for every function f in V the functional equation

$$f(x + y) = \sum_{i=1}^n g_i(x)h_i(y)$$

holds. Since the functions g_i, f are bounded and g_1, \dots, g_n are linearly independent, it follows from [7] that f, g_i, h_i are almost periodic functions on G . If \hat{G} denotes the dual of G we have by Fourier transformation, that

$$\hat{f}(\gamma)\gamma(y) = \sum_{i=1}^n \hat{g}_i(\gamma)h_i(y)$$

holds for every y in G and γ in \hat{G} . Repeating this argument with respect to y we have that

$$\tilde{f}(\gamma)\hat{\gamma}(\tau) = \sum_{i=1}^n \hat{g}_i(\gamma)\hat{h}_i(\tau)$$

holds for all γ and τ in \hat{G} . If $\gamma_i, \dots, \gamma_n$ are elements of \hat{G} such that the matrix $(\hat{g}_i(\gamma_j))$ is regular, then substituting γ_j for γ we have a linear system of equations for the unknowns $\hat{h}_i(\tau)$ ($i = 1, \dots, n$) which is homogeneous if $\tau \neq \gamma_j$ ($j = 1, \dots, n$). Hence we conclude that $\hat{h}_i(\tau) = 0$ for $\tau \neq \gamma_j$ ($i, j = 1, \dots, n$) and by the inversion theorem h_i is a trigonometric polynomial for $i = 1, \dots, n$. Since f is a linear combination of the functions h_i , we have that f is a trigonometric polynomial.

REFERENCES

1. P. M. Anselone and J. Korevaar, *Translation invariant subspaces of finite dimension*, Proc. Amer. Math. Soc., **15** (1964), 747-752.
2. D. Z. Djokovic, *A representation theorem for $(X_1-1)(X_2-1)\cdots(X_n-1)$ and its applications*, Ann. Polon. Math., **22** (1969), 189-198.
3. M. Engert, *Finite dimensional translation invariant subspaces*, Pacific J. Math.,

32 (1970), 333-343.

4. L. Hörmander, *Linear Partial Differential Operators*, Springer Verlag, Berlin, 1963.

5. P. G. Laird, *On characterizations of exponential polynomials*, Pacific J. Math., **80** (1979), 503-507.

6. L. Székelyhidi, *Polynomials on groups*, Publ. Math., Debrecen, (to appear).

7. ———, *Almost periodic functions and functional equations*, Acta Math. Szeged, **42** (1980), 165-169.

8. G. Van der Lijn, *La définition fonctionnelle des polynômes dans les groupes abéliens*, Fund. Math., **33** (1939), 42-50.

Received January 28, 1980.

KOSSUTH LAJOS UNIVERSITY
4010 DEBRECEN, HUNGARY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR AGUS

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF AAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies,

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966, Regular subscription rate: \$114.00 a year (6 Vol., 12 issues). Special rate: \$57.00 a year to individual members of supporting institution.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Pacific Journal of Mathematics

Vol. 103, No. 2

April, 1982

Alberto Alesina and Leonede De Michele , A dichotomy for a class of positive definite functions	251
Kahtan Alzubaidy , Rank ₂ p -groups, $p > 3$, and Chern classes	259
James Arney and Edward A. Bender , Random mappings with constraints on coalescence and number of origins	269
Bruce C. Berndt , An arithmetic Poisson formula	295
Julius Rubin Blum and J. I. Reich , Pointwise ergodic theorems in l.c.a. groups	301
Jonathan Borwein , A note on ε -subgradients and maximal monotonicity	307
Andrew Michael Brunner, Edward James Mayland, Jr. and Jonathan Simon , Knot groups in S^4 with nontrivial homology	315
Luis A. Caffarelli, Avner Friedman and Alessandro Torelli , The two-obstacle problem for the biharmonic operator	325
Aleksander Całka , On local isometries of finitely compact metric spaces	337
William S. Cohn , Carleson measures for functions orthogonal to invariant subspaces	347
Roger Fenn and Denis Karmen Sjerve , Duality and cohomology for one-relator groups	365
Gen Hua Shi , On the least number of fixed points for infinite complexes	377
George Golightly , Shadow and inverse-shadow inner products for a class of linear transformations	389
Joachim Georg Hartung , An extension of Sion's minimax theorem with an application to a method for constrained games	401
Vikram Jha and Michael Joseph Kallaher , On the Lorimer-Rahilly and Johnson-Walker translation planes	409
Kenneth Richard Johnson , Unitary analogs of generalized Ramanujan sums	429
Peter Dexter Johnson, Jr. and R. N. Mohapatra , Best possible results in a class of inequalities	433
Dieter Jungnickel and Sharad S. Sane , On extensions of nets	437
Johan Henricus Bernardus Kemperman and Morris Skibinsky , On the characterization of an interesting property of the arcsin distribution	457
Karl Andrew Kosler , On hereditary rings and Noetherian V -rings	467
William A. Lampe , Congruence lattices of algebras of fixed similarity type. II	475
M. N. Mishra, N. N. Nayak and Swadeenananda Pattanayak , Strong result for real zeros of random polynomials	509
Sidney Allen Morris and Peter Robert Nickolas , Locally invariant topologies on free groups	523
Richard Cole Penney , A Fourier transform theorem on nilmanifolds and nil-theta functions	539
Andrei Shkalikov , Estimates of meromorphic functions and summability theorems	569
László Székelyhidi , Note on exponential polynomials	583
William Thomas Watkins , Homeomorphic classification of certain inverse limit spaces with open bonding maps	589
David G. Wright , Countable decompositions of E^n	603
Takayuki Kawada , Correction to: "Sample functions of Pólya processes"	611
Z. A. Chanturia , Errata: "On the absolute convergence of Fourier series of the classes $H^\omega \cap V[v]$ "	611