

Pacific Journal of Mathematics

**ON ATTACHING 3-HANDLES TO A 1-CONNECTED
4-MANIFOLD**

BRUCE STEWART TRACE

ON ATTACHING 3-HANDLES TO A 1-CONNECTED 4-MANIFOLD[†]

BRUCE TRACE

This paper studies various properties of 3-handle presentations of 1-connected, smooth, 4-manifolds—with-connected-boundary. Although the question of whether or not such 4-manifolds have handle presentations consisting solely of 0-, 1- and 2-handles remains open, the main results of this paper do imply that the essential structure of such 4-manifolds is contained in a neighborhood of their 2-skeleton.

Throughout this article all maps and manifolds will be of class C^∞ . The key idea exploited here is that a knot in the boundary of a 4-manifold which is slice in this 4-manifold (i.e., which bounds a smooth, properly embedded 2-disc) may be viewed as the cocore of a 2-handle—the 2-handle being a tubular neighborhood of the slice disc. Thus, if M^4 is obtained from W^4 by attaching a 3-handle, h^3 , along an embedded $S^2 \times [-1, 1]$ in ∂W^4 , which is denoted $W^4 \bigcup_{\Sigma^2} h^3$ where Σ^2 is the image of $S^2 \times \{0\}$ under this embedding, then in order to construct a 2-handle which is complementary to h^3 we need only construct a knot $K \subset \partial W^4$ which meets Σ^2 transversely in a single point (i.e., K and Σ^2 are complementary in ∂W^4) and which is slice in W^4 . If the boundary of W^4 is connected, the existence of a knot $K \subset \partial W^4$ which is complementary to Σ^2 is clearly equivalent to Σ^2 not separating ∂W^4 . If W^4 is 1-connected, the following proposition tells us that a knot complementary to Σ^2 must be slice.

PROPOSITION 1. *Suppose W^4 is a 1-connected 4-manifold—with-connected-boundary. If K is a knot in ∂W^4 which is complementary to an embedded 2-sphere Σ^2 in ∂W^4 , then K is slice in W^4 .*

The proof of Proposition 1 is based on the Norman trick [6], which is discussed later in the paper.

The discussion above implies.

PROPOSITION 2. *Suppose $M = W^4 \bigcup_{\Sigma^2} h^3$ and $\tilde{M}^4 = W^4 \bigcup_{\tilde{\Sigma}^2} \tilde{h}^3$ are 1-connected 4-manifolds—with-connected-boundary. If there exists a knot K in ∂W^4 such that K is simultaneously complementary to both Σ^2 and $\tilde{\Sigma}^2$, then M^4 is diffeomorphic to \tilde{M}^4 .*

Standard 3-manifold techniques yield.

PROPOSITION 3. *Suppose W^3 is a connected, orientable, 3-manifold. If Σ^2 and $\tilde{\Sigma}^2$ are disjoint, nonseparating, embedded 2-spheres in the interior of W^3 , then there exists a knot K in W^3 which is simultaneously complementary to both Σ^2 and $\tilde{\Sigma}^2$.*

If $\Sigma^2 \cap \tilde{\Sigma}^2 \neq \emptyset$, by isotoping Σ^2 to Σ_0^2 where Σ_0^2 meets $\tilde{\Sigma}^2$ transversely, then by performing cutting and pasting techniques to innermost circles of intersection in Σ_0^2 together with an induction argument we arrive at

PROPOSITION 4. *Suppose W^3 is a connected, orientable, 3-manifold. If Σ^2 and $\tilde{\Sigma}^2$ are nonseparating, embedded 2-spheres in the interior of W^3 , then there exists a finite sequence $\Sigma_0^2 \rightarrow \Sigma_1^2 \rightarrow \cdots \rightarrow \Sigma_m^2 = \tilde{\Sigma}^2$ of nonseparating, embedded 2-spheres in the interior of W^3 such that $\Sigma_i^2 \cap \Sigma_{i+1}^2 = \emptyset$ for $i = 0, 1, \dots, m - 1$, and Σ_0^2 is isotopic to Σ^2 .*

Propositions 2, 3, and 4 combine to yield

PROPOSITION 5. *Suppose $M^4 = W^4 \mathbf{U}_{\Sigma^2} h^3$ and $\tilde{M}^4 = W^4 \mathbf{U}_{\tilde{\Sigma}^2} \tilde{h}^3$ are 1-connected 4-manifolds—with—connected-boundary, then M^4 is diffeomorphic to \tilde{M}^4 .*

Proposition 5 together with an inductive argument are used to prove the nontrivial implication in

THEOREM 1. *Suppose $M^4 = W^4 \mathbf{U}_{\Sigma_1^2} h_1^3 \cup \cdots \cup \mathbf{U}_{\Sigma_k^2} h_k^3$ and $\tilde{M}^4 = \tilde{W}^4 \mathbf{U}_{\tilde{\Sigma}_1^2} \tilde{h}_1^3 \cup \cdots \cup \mathbf{U}_{\tilde{\Sigma}_k^2} \tilde{h}_k^3$ are 1-connected, 4-manifolds—with—connected-boundary. Then M^4 is diffeomorphic to \tilde{M}^4 if and only if W^4 is diffeomorphic to \tilde{W}^4 .*

REMARK. In [4] F. Laudenbach and V. Poenaru showed (essentially) that Theorem 1 remains true if we assume $\partial M^4 = S^3$ rather than $\pi_1(M^4) = 1$. (See [5] for details.)

By introducing k complementary 2-, 3-handle pairs to W^4 then employing Theorem 1 to reattach these complementary 3-handles so as to form $M^4 \cup k$ (2-handles) from $W^4 \cup k$ (complementary 2-, 3-handle pairs), we observe that M^4 embeds in W^4 . Hence we observe

THEOREM 2. *Suppose M^4 and W^4 are as in the statement of Theorem 1. Then, if N^4 is an arbitrary 4-manifold, M^4 embeds in N^4 if and only if W^4 embeds in N^4 .*

Theorem 1 together with the cutting and pasting argument of Proposition 4 imply

THEOREM 3. *Suppose M^4 and W^4 are as in the statement of Theorem 1, and $W^4 = W_1^4 \#_0 W_2^4$. Then $M^4 = M_1^4 \#_0 M_2^4$ where $M_i^4 = W_i^4 \cup (3\text{-handles})$ for $i = 1, 2$.*

I would like to thank R. D. Edwards for his suggestions. These suggestions served to simplify many points in the proof of Theorem 1.

DEFINITIONS AND NOTATION. We shall assume the reader is familiar with the basic definitions and results of differential topology—in particular handle theory.

We shall denote the unit ball in R^n by B^n . Any diffeomorph of B^n is called an n -disc. The $(n - 1)$ -sphere is denoted by S^{n-1} . If M^n is an n -manifold—with—boundary we denote the boundary of M^n by ∂M^n . The interior of M^n , denoted $\text{int } M^n$, is $\text{int } M^n = M^n - \partial M^n$. If X is a subset of M^n , then we shall use $\text{cl } X$ to denote the closure of X in M^n .

If M_1^n and M_2^n are two manifolds, then $M_1^n \# M_2^n$ denotes the (interior) connected sum of M_1^n with M_2^n and the boundary connected sum of M_1^n with M_2^n is denoted by $M_1^n \#_0 M_2^n$.

Throughout this paper lower case “ h ” is used only for “handle”. The symbols h^r , h_i^r and \tilde{h}_i^r all represent r handles. The subscript denotes the order of attaching.

An arc is an embedded 1 disc and a knot in a 3-manifold is an embedded S^1 .

Proof of Proposition 1. Since W^4 is 1-connected there exists an immersed 2-disc $\mathcal{A}_1^2 \subset W^4$ bounded by K . We may assume that \mathcal{A}_1^2 is properly immersed in W^4 (i.e. $\mathcal{A}_1^2 \cap \partial W^4 = \partial \mathcal{A}_1^2$) and that the singular set of \mathcal{A}_1^2 consists of a finite number of transverse double points. Since Σ^2 meets K transversely in a single point, there exists an embedded 2-sphere $\Sigma_1^2 \subset \text{int } W^4$ such that Σ_1^2 meets \mathcal{A}_1^2 transversely in a single point, say g .

If p is a double point of \mathcal{A}_1^2 , let α be an arc in \mathcal{A}_1^2 joining p to g such that α meets no other double point of \mathcal{A}_1^2 . We now apply the Norman trick. This means we pipe \mathcal{A}_1^2 to Σ_1^2 via a small annulus contained in the normal bundle of \mathcal{A}_1^2 restricted to α . The net result of this piping operation is to cancel the double point p of \mathcal{A}_1^2 against the point $g = \mathcal{A}_1^2 \cap \Sigma_1^2$. Thus we have replaced the 2-disc \mathcal{A}_1^2 with a new 2-disc, \mathcal{A}_2^2 , which is bounded by K and has one fewer double point than \mathcal{A}_1^2 . Inductively we obtain K bounds a properly embedded 2-disc, i.e., K is slice in W^4 .

Proof of Proposition 2. By Proposition 1, such a knot K is slice

in W^4 . Let \mathcal{A}^2 be a slice disc for K in W^4 . If $N(\mathcal{A}^2)$ is a small tubular neighborhood of \mathcal{A}^2 then $N(\mathcal{A}^2)$ is a trivial 2-discbundle over \mathcal{A}^2 . Hence, W^4 is obtained from $\text{cl}(W^4 - N(\mathcal{A}^2))$ by attaching a 2-handle, namely $N(\mathcal{A}^2)$, where \mathcal{A}^2 is the cocore disc of $N(\mathcal{A}^2)$. Since K is simultaneously complementary to both Σ^2 and $\tilde{\Sigma}^2$, we obtain the 2-handle $N(\mathcal{A}^2)$ is complementary to each of the 3-handle h^3 and \tilde{h}^3 . Therefore both M^4 and \tilde{M}^4 are diffeomorphic to $\text{cl}(W^4 - N(\mathcal{A}^2))$.

Proof of Proposition 3. Since neither Σ^2 nor $\tilde{\Sigma}^2$ separate W^3 we must have that $W^3 - (\Sigma^2 \cup \tilde{\Sigma}^2)$ consists of at most two components. If $W^3 - (\Sigma^2 \cup \tilde{\Sigma}^2)$ is connected, the existence of the desired knot K is obvious. If $W^3 - (\Sigma^2 \cup \tilde{\Sigma}^2)$ consists of two components, say C_1 and C_2 , choose points $p \in \Sigma^2$ and $\tilde{p} \in \tilde{\Sigma}^2$. Because both $W^3 - \Sigma^2$ and $W^3 - \tilde{\Sigma}^2$ are connected and C_i , $i = 1, 2$, is arc connected, there exist arcs α_i properly embedded in $\text{cl}(C_i)$, $i = 1, 2$, joining p to \tilde{p} . Thus $K = \alpha_1 \cup \alpha_2$ is the desired knot.

Proof of Proposition 4. We may isotope Σ^2 to Σ_0^2 , where $\Sigma_0^2 \cap \tilde{\Sigma}^2$ consists of a finite collection of transverse, disjoint circles. Let σ^1 be an innermost circle of intersection in $\tilde{\Sigma}^2$ bounding the innermost disc $\tilde{\mathcal{A}}^2 \subset \tilde{\Sigma}^2$, i.e., $\partial\tilde{\mathcal{A}}^2 = \sigma^1$ and $\Sigma_0^2 \cap (\text{int}\tilde{\mathcal{A}}^2) = \emptyset$.

Let $\mathcal{A}_1^2 \cup \mathcal{A}_2^2 = \Sigma_0^2$, where $\mathcal{A}_1^2 \cap \mathcal{A}_2^2 = \sigma^1$. We may isotope the 2-spheres $\tilde{\mathcal{A}}^2 \cup \mathcal{A}_i^2$ to disjoint 2-spheres Σ_{1i}^2 , $i = 1, 2$, where both Σ_{11}^2 and Σ_{12}^2 meet $\tilde{\Sigma}^2$ in at least one fewer component than Σ_0^2 and $\Sigma_{1i}^2 \cap \Sigma_0^2 = \emptyset$, $i = 1, 2$. Since the connected sum of Σ_{11}^2 with Σ_{12}^2 gives back Σ_0^2 (up to isotopy), we must have either Σ_{11}^2 or Σ_{12}^2 not separating W^3 . Let Σ_1^2 be this noseparating 2 sphere.

Since Σ_1^2 meets $\tilde{\Sigma}^2$ in fewer components than does Σ_0^2 , we may inductively continue the process until we come to a 2-sphere which is disjoint from $\tilde{\Sigma}^2$. Having done so, we will have constructed the desired sequence of 2-spheres.

Proof of Proposition 5. Since both M^4 and \tilde{M}^4 have connected boundary, it follows that Σ^2 and $\tilde{\Sigma}^2$ are nonseparating 2-spheres in the connected, orientable 3-manifold ∂W^4 . Thus, by Proposition 4, there exists a 2-sphere $\Sigma_0^2 \subset \partial W^4$ which is isotopic to Σ^2 and a finite sequence of nonseparating 2-spheres $\Sigma_0^2 \rightarrow \Sigma_1^2 \rightarrow \dots \rightarrow \Sigma_m^2 = \tilde{\Sigma}^2$ where $\Sigma_i^2 \cap \Sigma_{i+1}^2 = \emptyset$, $i = 0, 1, \dots, m-1$. By Proposition 3 there exists a knot $K_i \subset \partial W^4$ such that K_i is simultaneously complementary to both Σ_i^2 and Σ_{i+1}^2 for $i = 0, 1, \dots, m-1$.

Set $M_i^4 = W^4 \bigcup_{\Sigma_i^2} h_i^3$ for $i = 0, 1, \dots, m-1$. Since Σ^2 is isotopic to Σ_0^2 , we have M^4 is diffeomorphic to M_0^4 . We apply Proposition 2 to obtain M_i^4 is diffeomorphic to M_{i+1}^4 , $i = 0, \dots, m-1$. Thus M^4 is diffeomorphic to \tilde{M}^4 .

Proof of Theorem 1. Suppose $f: M^4 \rightarrow \tilde{M}^4$ is a diffeomorphism. Since the cocore of 3-handle is 1-dimensional and M^4 is simply connected, there exists an ambient isotopy of \tilde{M}^4, F_i , such that F_0 is the identity and $F_i[f(\text{cocore of } h_i^3)] = (\text{cocore of } \tilde{h}_i^3)$, $i = 1, \dots, k$. Then both $F_i(f(h_i^3))$ and \tilde{h}_i^3 are tubular neighborhoods of the cocore of \tilde{h}_i^3 , $i = 1, \dots, k$. Therefore there exists an isotopy G_i of \tilde{M}^4 such that G_0 is the identity and $G_i(F_i(f(h_i^3))) = \tilde{h}_i^3$, $i = 1, \dots, k$. Set $g = G_1 \circ F_1 \circ f: M^4 \rightarrow \tilde{M}^4$. Then $g|W^4$ is a diffeomorphism of W^4 onto \tilde{W}^4 .

As previously stated, the nontrivial implication in Theorem 1 proceeds via induction on k . Let $g: W^4 \rightarrow \tilde{W}^4$ be a diffeomorphism assume $k = 1$. Set $g(M^4) = \tilde{W} \bigcup_{g(\Sigma_1^2)} h^3$. Then g induces a diffeomorphism of M^4 onto $g(M^4)$. Therefore $g(M^4)$ and \tilde{M}^4 satisfy the hypothesis of Proposition 5. This yields $g(M^4)$ is diffeomorphic to \tilde{M}^4 and hence the theorem is true for $k = 1$.

Set

$$W_*^4 = W^4 \bigcup_{\Sigma_1^2} h_1^3 \cup \dots \bigcup_{\Sigma_{k-1}^2} h_{k-1}^3 \quad \text{and} \quad \tilde{W}_*^4 = \tilde{W}^4 \bigcup_{\tilde{\Sigma}_1^2} \tilde{h}_1^3 \cup \dots \bigcup_{\tilde{\Sigma}_{k-1}^2} \tilde{h}_{k-1}^3.$$

By induction, if W^4 is diffeomorphic to \tilde{W}^4 then W_*^4 is diffeomorphic to \tilde{W}_*^4 . By repeating the argument used in the $k = 1$ case to $M^4 = W_*^4 \bigcup_{\Sigma_1^2} h_k^3$ and $\tilde{M}^4 = \tilde{W}_*^4 \bigcup_{\tilde{\Sigma}_k^2} \tilde{h}_k^3$ we obtain M^4 is diffeomorphic to \tilde{M}^4 which completes the proof of Theorem 1.

Proof of Theorem 2. Let $\mathcal{L}^3 \subset \partial W^4$ be a 3-cell disjoint from $\bigcup_{i=1}^k \Sigma_i^2$. Attach k complementary 2-,3-handle pairs to W^4 with attaching tubes meeting ∂W^4 in the interior of \mathcal{L}^3 . Observing that $W^4 \cup k$ (2-handles) is simply connected if W^4 is simply connected, we apply Theorem 1 to alter the fashion in which 3-handles are attached to $W^4 \cup k$ (2-handles). We reattach the 3-handle which initially is complementary to the i th 2-handle to Σ_i^2 .

Since W^4 union k complementary 2-,3-handle pairs is diffeomorphic to W^4 upon altering the fashion in which 3-handles are attached to $W^4 \cup k$ (2-handles), we have that W^4 is diffeomorphic to $M^4 \cup k$ (2-handles). Hence any embedding of W^4 into N^4 yields an embedding of M^4 into N^4 .

The converse direction is trivial.

Proof of Theorem 3. Let $\Sigma^2 \subset \partial W^4$ be the 2-sphere along which we connect sum ∂W_1^4 to ∂W_2^4 to obtain ∂W^4 . We may isotope the Σ_i^2 , $i = 1, \dots, k$, so that they meet Σ^2 transversely in a finite collection of circles. Proceeding as in Proposition 4, the collection Σ_i^2 , $i = 1, \dots, k$, may be replaced with another collection of 2-spheres $\tilde{\Sigma}_i^2$, $i = 1, \dots, k$ such that $\tilde{\Sigma}_i^2 \cap \Sigma^2 = \emptyset$ for $i = 1, \dots, k$, and $\partial W^4 - (\bigcup_{i=1}^k \tilde{\Sigma}_i^2)$ is connected.

Set $\tilde{M}^4 = W^4 \mathbf{U}_{\tilde{\Sigma}_k^2} \tilde{h}_1^3 \cup \dots \cup \mathbf{U}_{\tilde{\Sigma}_k^2} \tilde{h}_k^3$ and observe that M^4 and \tilde{M}^4 satisfy the hypothesis of Theorem 1. Hence M^4 is diffeomorphic to \tilde{M}^4 . Since $\tilde{\Sigma}_i^2 \cap \Sigma^2 = \emptyset$ for $i = 1, \dots, k$, we have $\tilde{M}^4 = \tilde{M}_1^4 \#_0 \tilde{M}_2^4$ where $\tilde{M}_i^4 = \tilde{W}_i^4 \cup (3\text{-handles})$, $i = 1, 2$. This implies the desired result.

Some Corollaries.

COROLLARY 1. *Suppose $M^4 = B^4 \cup k(2\text{-handles}) \cup k(3\text{-handles}) \cong pt$. Then M smoothly embeds in S^4 if and only if the link along which the 2-handles are attached to B^4 is strongly slice in B^4 .*

Sketch of Proof. Since M^4 is contractible, each 2-handle is attached to B^4 via 0-framing. We leave as an exercise the fact that $B^4 \cup k(2\text{-handles})$ smoothly embed in S^4 if and only if the link along which 2-handles are attached to B^4 is strongly slice, i.e., each component is slice and the slice discs can be chosen to be mutually disjoint, and each 2-handle is attached via 0-framing. Theorem 2 implies M^4 smoothly embeds in S^4 if $B^4 \cup k(2\text{-handles})$ does.

COROLLARY 2. *Suppose $M^4 = B^4 \mathbf{U}_K(2\text{-handle}) \cup (3\text{-handle}) \cong pt$. Then the punctured double of M^4 smoothly embeds in $B^4 \mathbf{U}_K(2\text{-handle})$. Hence, $\text{Punct}(DM^4)$ smoothly embeds in S^4 if and only if K is slice.*

Proof. Write $M^4 = B^4 \mathbf{U}_K h^2 \cup h^3$. Then $\text{Punct}(DM^4) = B^4 \mathbf{U}_K h^2 \cup h^3 \cup (h^3)^* \cup (h^2)^*$, where $(h^3)^*$ is a 1-handle dual to h^3 and $(h^2)^*$ is a 2-handle dual to h^2 .




FIGURE 1

We attach a 2-handle, $(h^2)_m^*$, to $\text{Punct}(DM^4)$. This 2-handle is attached via 0-framing along a meridian of the attaching sphere of $(h^2)^*$, see Figure 1.

Then by sliding h^2 over $(h^2)_m^*$ we can untangle the a -spheres of h^2 and $(h^2)^*$. Since M^4 is contractible, we must have $(h^2)^*$ algebraically cancelling $(h^3)^*$. By sliding $(h^2)^*$ over $(h^2)_m^*$ we can arrange that $(h^2)^*$ geometrically cancels $(h^3)^*$. Thus, $\text{Punct}(DM^4) \subset B^4 \bigcup_K h^2 \cup h^3 \cup (h^2)_m^*$. Since the a -sphere of h^2 and the a -sphere of $(h^2)_m^*$ lie in disjoint 3-cells of ∂B^4 and since the a -sphere for $(h^2)_m^*$ is the unknot—we may apply Theorem 1 so as to reattach h^3 so that it geometrically cancels $(h^2)_m^*$. Thus $\text{Punct}(DM^4) \subset B^4 \bigcup_K h^2$.

Final Remarks. (1) In [3], R. Kirby and P. Melvin observed that K as in Corollary 2 is slice if $\partial M = S^3$.

(2) Recently, S. Akbulut and R. Kirby have obtained a $M^4 = B^4 \cup 2$ (2-handles) $\cup 2$ (3-handles) $\cong pt.$ where the link along which 2-handles are attached is not known to be slice, (see [1] and [2]).

REFERENCES

1. S. Akbulut and R. Kirby, *An Exotic Involution of S^4* , *Topology*, **18** (1979), 75-81.
2. S. Akbulut and R. Kirby, to appear.
3. R. Kirby and P. Melvin, *Slice Knots and Property R*, *Inventiones Math.*, **45** (1978), 57-59.
4. F. Laudenbach and V. Poenaru, *A Note on 4-Dimensional Handlebodies*, *Bull. Soc. Math., France*, **100** (1972), 337-347.
5. J. Montesinos, *Heegard Diagrams For 4-Manifolds*, Preprint.
6. R. Norman, *Dehn's Lemma for Certain 4-Manifolds*, *Inventiones Math.*, **7** (1969), 143-147.

Received October 13, 1980. Research partially supported by NSF grant MCS 7606903.

UNIVERSITY OF CALIFORNIA
LOS ANGELES, CA 920024

Present Address: University of Utah
Salt Lake City, UT 84112

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR AGUS

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

Mariano Giaquinta, Jindrich Necas, O. John and J. Stará, On the regularity up to the boundary for second order nonlinear elliptic systems . . .	1
Siegfried Graf, Realizing automorphisms of quotients of product σ -fields . . .	19
Alfred Washington Hales and Ernst Gabor Straus, Projective colorings . . .	31
Sandra Hayes, The weak Nullstellensatz for finite-dimensional complex spaces	45
Gerald Norman Hile and Murray Harold Protter, The Cauchy problem and asymptotic decay for solutions of differential inequalities in Hilbert space	57
Robert D. Little, Projective space as a branched covering with orientable branch set	89
Jaroslav Mach, On the proximality of Stone-Weierstrass subspaces	97
John C. Morgan, II, On product bases	105
K. Balakrishna Reddy and P. V. Subrahmanyam, Altman's contractors and fixed points of multivalued mappings	127
James Ted Rogers Jr., Decompositions of homogeneous continua	137
Ahmed Ramzy Sourour, Characterization and order properties of pseudo-integral operators	145
Robert Moffatt Stephenson Jr., Pseudocompact and Stone-Weierstrass product spaces	159
Bruce Stewart Trace, On attaching 3-handles to a 1-connected 4-manifold	175
Akihito Uchiyama, The construction of certain BMO functions and the corona problem	183
Thomas Alva Whitehurst, An application of orthogonal polynomials to random walks	205
David J. Winter, Root locologies and idempotents of Lie and nonassociative algebras	215
William Robin Zame, The classification of uniform algebras on plane domains	231