

# Pacific Journal of Mathematics

## **BMO FROM DYADIC BMO**

JOHN BRADY GARNETT AND PETER WILCOX JONES

## BMO FROM DYADIC BMO

JOHN B. GARNETT AND PETER W. JONES

**We give new proofs of four decomposition theorems for functions of bounded mean oscillation by first obtaining each theorem in the easier dyadic case and then averaging the results of the dyadic decomposition over translations in  $R_m$ .**

**1. Introduction.** Let  $\varphi$  be a locally integrable real function on  $R^m$ , let  $Q$  be a bounded cube in  $R^m$ , with sides parallel to the axes, and let  $|Q|$  be the Lebesgue measure of  $Q$ . Then

$$\varphi_Q = \frac{1}{|Q|} \int_Q \varphi dx$$

is the average of  $\varphi$  over  $Q$ . We say  $\varphi$  has *bounded mean oscillation*,  $\varphi \in \text{BMO}$ , if

$$\|\varphi\| = \sup_Q \frac{1}{|Q|} \int_Q |\varphi - \varphi_Q| dx < \infty .$$

A *dyadic cube* is a cube of the special form

$$Q = \{k_j 2^{-n} < x_j < (k_j + 1)2^{-n}; 1 \leq j \leq m\}$$

where  $n$  and  $k_j$ ,  $1 \leq j \leq m$ , are integers, and  $\varphi$  has *bounded dyadic mean oscillation*,  $\varphi \in \text{BMO}_d$ , if

$$\|\varphi\|_d = \sup_{Q \text{ dyadic}} \frac{1}{|Q|} \int_Q |\varphi - \varphi_Q| dx < \infty .$$

Then clearly  $\text{BMO} \subset \text{BMO}_d$  with  $\|\varphi\|_d \leq \|\varphi\|$ , but  $\text{BMO}$  and  $\text{BMO}_d$  are not the same space; the function  $\log|x_j| \chi_{\{x_j > 0\}}$  is in  $\text{BMO}_d$  but not in  $\text{BMO}$ . In analysis  $\text{BMO}$  is more important than  $\text{BMO}_d$  because  $\text{BMO}$  is translation invariant, but  $\text{BMO}_d$  is not. On the other hand,  $\text{BMO}_d$  is very much the easier space to work with because dyadic cubes are nested (if two open dyadic cubes intersect then one of them is contained in the other). For example, for  $\text{BMO}$  the original proofs [1], [6], [8], [11] of the four theorems stated below were rather technical, while for  $\text{BMO}_d$  the analogous results are comparatively trivial. In this paper we derive the four theorems from their dyadic counterparts.

Here is the idea. Let  $T_\alpha \varphi(x) = \varphi(x - \alpha)$ . Then

$$\varphi(x) = \lim_{N \rightarrow \infty} \frac{1}{(2N)^m} \int_{|\alpha_j| \leq N} T_\alpha \varphi(x + \alpha) d\alpha .$$

Each of the theorems amounts to showing  $\varphi \in \text{BMO}$  has the form  $\varphi = F_1 + F_2$  where  $F_1$  and  $F_2$  are BMO functions satisfying certain additional growth conditions. By the  $\text{BMO}_d$  result we have

$$T_\alpha \varphi = F_1^{(\alpha)} + F_2^{(\alpha)}$$

where  $F_1^{(\alpha)}, F_2^{(\alpha)} \in \text{BMO}_d$  satisfy the extra growth conditions on dyadic cubes. To prove each theorem we show the averages

$$F_j(x) = \lim_{N \rightarrow \infty} \frac{1}{(2N)^m} \int_{|\alpha_j| \leq N} F_j^{(\alpha)}(x + \alpha) d\alpha$$

are in BMO and have the correct growth. The method yields this general result.

**THEOREM.** *Suppose that  $\alpha \rightarrow \varphi^{(\alpha)}$  is a measurable mapping from  $\mathbf{R}^m$  to  $\text{BMO}_d$  such that all  $\varphi^{(\alpha)}(x)$  have support a fixed dyadic cube, such that  $\|\varphi^{(\alpha)}\|_d \leq 1$  and such that*

$$\int \varphi^{(\alpha)}(x) dx = 0 .$$

Then

$$\varphi_N(x) = \frac{1}{(2N)^m} \int_{|\alpha_j| \leq N} \varphi^{(\alpha)}(x + \alpha) d\alpha$$

is in BMO and  $\|\varphi_N\| \leq C$ .

By duality, this theorem implies Davis's result connecting  $H^1$  and  $H^1_{\text{dyadic}}$  on the unit circle. The proof of theorem is implicit in the arguments below. In §4 we show

$$\varphi_N = g + \sum_{n=1}^{\infty} f_n$$

where  $g \in L^\infty$  and where  $f_n(x)$  satisfies the Lipschitz condition (3.3) and the thinness condition (4.2). From these  $\|\varphi_N\| \leq C$  follows easily. This general result is not explicitly used in the proofs of Theorem 1 to 4.

Let  $\ell(Q)$  denote the sidelength of the cube  $Q$ . A Carleson measure is a signed measure on the upper half space  $\mathbf{R}^{m+1}_+ = \mathbf{R}^m \times (0, \infty)$  such that for some constant  $N(\sigma)$ ,

$$|\sigma|(Q \times (0, \ell(Q))) \leq N(\sigma) |Q|$$

for all cubes  $Q \subset \mathbf{R}^m$ . Here  $|\sigma|$  is the total variation of  $\sigma$ . Let  $K(x)$  be a positive function for which

$$(1.1) \quad K(x) = O((1 + |x|)^{-m-1})$$

and

$$\int K(x)dx = 1 .$$

Write  $\check{K}_y(x) = y^{-m}K(x/y)$ ,  $y > 0$ .

**THEOREM 1** (Carleson [1]). *If  $\varphi \in \text{BMO}$  has compact support, then there is  $g \in L^\infty$  and there is a Carleson measure  $\sigma$  such that*

$$(1.2) \quad \varphi(x) = g(x) + \int_{\mathbf{R}_+^{m+1}} K_y(x - t)d\sigma(t, y) ,$$

where

$$\|g\|_\infty \leq C\|\varphi\|$$

and

$$N(\sigma) \leq C\|\varphi\| ,$$

where the constant  $C$  depends only on  $K(x)$ .

Theorem 1 implies Fefferman's Theorem [5] that  $H^1(\mathbf{R}^m)$  has dual space BMO. Under the additional hypotheses

$$|\nabla K(x)| = O((1 + |x|)^{-m-1}) ,$$

the converse of Theorem 1 is true (and not difficult). It then follows that  $H^1(\mathbf{R}^m) = \{f \in L^1: f_K^* \in L^1\}$  where  $f_K^*$  is the maximal function  $\sup_{|t-x|<y} |f * K^y(t)|$ . See [5].

By the theorem of John and Nirenberg [7],  $\varphi \in \text{BMO}$  if and only if there is  $A > 0$  such that

$$(1.3) \quad \sup_Q \frac{1}{|Q|} \int_Q e^{A|\varphi - \varphi_Q|} dx < \infty .$$

In fact, (1.3) holds with  $A = c\|\varphi\|^{-1}$ ,  $c$  depending only on the dimension. Set

$$A(\varphi) = \sup\{A: (1.3) \text{ holds}\} .$$

**THEOREM 2** ([6]). *There are constants  $c_1(m)$  and  $c_2(m)$  such that if  $\varphi \in \text{BMO}$  then*

$$\frac{c_1(m)}{A(\varphi)} \leq \inf_{g \in L^\infty} \|\varphi - g\| \leq \frac{c_2(m)}{A(\varphi)} .$$

The left inequality is immediate since  $A(\varphi - g) \geq c\|\varphi - g\|^{-1}$ ,  $g \in L^\infty$ . We prove the other inequality.

Let  $w(x) > 0$  be a locally integrable function on  $\mathbf{R}^m$ , and let  $1 < p < \infty$ . We say  $w \in A_p$  if

$$\|w\|_{A_p} = \sup_Q \left( \frac{1}{|Q|} \int_Q w dx \right) \left( \frac{1}{|Q|} \int_Q \left( \frac{1}{w} \right)^{1/(p-1)} dx \right)^{p-1} < \infty ;$$

The Riesz transforms and the Hardy-Littlewood maximal functions are bounded on  $L^p(w dx)$  if and only if  $w \in A_p$  [2]. As  $p \rightarrow 1$  the limiting form of  $A_p$  is

$$\|w\|_{A_1} = \sup_Q \left( \frac{1}{|Q|} \int_Q w dx \right) \left( \left\| \frac{1}{w} \right\|_{L^\infty(Q)} \right),$$

and we say  $w \in A_1$  if  $\|w\|_{A_1} < \infty$ .

**THEOREM 3** ([8]). *If  $1 < p < \infty$ , then  $w \in A_p$  if and only if*

$$(1.4) \quad w = w_1(w_2)^{1-p}$$

where  $w_1, w_2 \in A_1$ .

Hölder's inequality shows that (1.4) is sufficient. Obtaining the factorization (1.4) for  $w \in A_p$  is more difficult.

Theorem 2 is a simple consequence of Theorem 3. Indeed, let  $\varphi \in \text{BMO}$  and take  $A(\varphi)/2 < A < A(\varphi)$ . Write  $w = e^{A\varphi}$ . Then for any  $Q$ ,

$$\begin{aligned} \left( \frac{1}{|Q|} \int_Q w dx \right) \left( \frac{1}{|Q|} \int_Q \frac{1}{w} dx \right) &= \left( \frac{1}{|Q|} \int_Q e^{A(\varphi-Q)} dx \right) \left( \frac{1}{|Q|} \int_Q e^{-A(\varphi-Q)} dx \right) \\ &\leq \left( \frac{1}{|Q|} \int_Q e^{A|\varphi-Q|} dx \right)^2, \end{aligned}$$

so that  $w \in A_2$ . By Theorem 3,

$$A\varphi = \log w = F_1 - F_2$$

where  $e^{F_1}, e^{F_2} \in A_1$ . From  $A_1$  it follows easily that

$$e^{F_j} \leq M(e^{F_j}) \leq ce^{F_j}$$

almost everywhere, where  $M(f)$  denotes the Hardy-Littlewood maximal function of  $f$ . Coifman and Rochberg [3] have shown  $\|\log M(f)\|_{\text{BMO}} \leq C(m)$  whenever  $f \in L^1_{\text{loc}}$ . Consequently

$$\begin{aligned} F_j &= \log M(e^{F_j}) + \log(e^{F_j}/M(e^{F_j})) \\ &= \psi_j + g_j \end{aligned}$$

where  $g_j \in L^\infty$  and  $\|\psi_j\| \leq C(m)$ . Hence

$$\varphi = \frac{g_1 - g_2}{A} + \frac{\psi_1 - \psi_2}{A} = g + \psi$$

with  $g \in L^\infty$  and  $\|\psi\| \leq 2C(m)/A$ .

The above reasoning also explains why Theorem 3 is a theorem about BMO. See [8] for further application of Theorem 3.

**THEOREM 4 (Uchiyama [11]).** *Let  $\lambda > 0$  and let  $E_1, E_2, \dots, E_N$  be measurable subsets of  $\mathbf{R}^m$  such that*

$$(1.5) \quad \text{Min}_{1 \leq i \leq N} \frac{|Q \cap E_i|}{|Q|} \leq 2^{-2m\lambda}$$

*for every cube  $Q$ . Then there exists functions  $f_1(x), f_2(x), \dots, f_N(x)$  such that almost everywhere*

$$(1.6) \quad f_i(x) = 0, \quad x \in E_i.$$

$$(1.7) \quad 0 \leq f_i(x) \leq 1.$$

$$(1.8) \quad \sum_{i=1}^N f_i(x) = 1$$

*and such that*

$$(1.9) \quad \|f_i\| \leq C(m, N)/\lambda, \quad 1 \leq i \leq N.$$

The converse (with  $\|f_i\| \leq C'(m, N)/\lambda$ ) of this theorem is not difficult. Theorem 4 for  $N = 2$  is roughly equivalent to Theorem 2. For  $N > 2$  it has interesting applications to function theory. See [8] and [11].

In §2 we prove the dyadic versions of Theorem 1 and Theorem 3. Although the arguments are well known (see [13] and [8]), they are included for completeness and because some of their by-products will be needed later. Theorem 3 is proved in §3 and Theorem 1 is proved in §4. In §5 we discuss Theorem 4 and its dyadic analogue.

We would like to acknowledge our indebtedness to Davis [4], who showed on the circle that  $T_\alpha f \in H^1_{\text{dyadic}}$  for almost every  $\alpha$  if  $f \in H^1$ , and to Varopoulos [12], who proved Theorem 2 by adapting the argument of the dyadic case to Brownian motion.

## 2. Two dyadic theorems.

**THEOREM 2.1.** *Let  $\varphi \in \text{BMO}_d$  and let  $Q_0$  be a fixed dyadic cube. Then there exists a sequence  $\{Q_k\}$  of dyadic cubes  $Q_k \subset Q_0$ , and a sequence  $\{a_k\}$  of real numbers such that*

$$(2.1) \quad \sum_{Q_k \subset Q} |a_k| |Q_k| \leq C \|\varphi\|_d |Q|,$$

for all dyadic cubes  $Q$ , and there exists  $g \in L^\infty$ ,

$$\|g\|_\infty \leq 2 \|\varphi\|_d,$$

such that

$$(2.2) \quad \varphi(x) - \varphi_{Q_0} = g(x) + \sum a_k \chi_{Q_k}(x)$$

almost everywhere on  $Q_0$ . The constant  $C$  depends only on the dimension.

To understand why Theorem 2.1 is the dyadic formulation of Theorem 1, replace  $\mathbf{R}_+^{m+1}$  by its discrete subset  $\mathcal{S} = \{p_Q = (c(Q), \ell(Q)), Q \text{ dyadic}\}$  where  $c(Q) \in \mathbf{R}^m$  is the center of  $Q$  and  $\ell(Q)$  is the sidelength of  $Q$ . The correspondence between  $p_Q$  and  $K_Q(x) = \chi_Q(x)/|Q|$  resembles the correspondence between  $(t, y) \in \mathbf{R}_+^{m+1}$  and  $K_y(x - t)$ . Let  $\sigma$  be the measure on  $\mathcal{S}$  having mass  $a_k |Q|$  at  $p_{Q_k}$ . Then (2.1) says that

$$\begin{aligned} |\sigma|(Q \times (0, \ell(Q))) &= \sum_{Q_k \subset Q} |a_k| |Q_k| \\ &\leq C \|\varphi\|_d |Q| \end{aligned}$$

and  $\sigma$  can be viewed as a dyadic Carleson measure. Since

$$\int K_Q(x) d\sigma(p_Q) = \sum a_k \chi_{Q_k}(x),$$

(2.2) is now the dyadic version of (1.2).

*Proof.* We suppose  $\varphi_{Q_0} = 0$ . Fix  $\lambda = 2 \|\varphi\|_d$  and set

$$G_1 = \{Q_k \subset Q_0: Q_k \text{ dyadic, } |\varphi_{Q_k}| > \lambda, \text{ and } Q_k \text{ maximal}\}.$$

Because  $Q_k \in G_1$  is maximal, we have

$$(2.3) \quad |\varphi_{Q_k}| \leq \lambda + 2^m \|\varphi\|_d \leq 2^{m+1} \|\varphi\|_d.$$

Indeed, if  $Q_k^*$  is that dyadic cube with  $Q_k^* \supset Q_k$  and  $|Q_k^*| = 2^m |Q_k|$ , then

$$|\varphi_{Q_k} - \varphi_{Q_k^*}| \leq \frac{1}{|Q_k|} \int_{Q_k^*} |\varphi - \varphi_{Q_k^*}| dx \leq 2^m \|\varphi\|_d$$

and  $|\varphi_{Q_k^*}| \leq \lambda$  as  $Q_k$  is maximal. The  $Q_k$  in  $G_1$  are pairwise disjoint, because they are maximal, so that

$$(2.4) \quad \sum_{G_1} |Q_k| \geq \frac{1}{\lambda} \sum_{G_1} \left| \int_{Q_k} \varphi dx \right| \leq \frac{1}{\lambda} \int_{Q_0} |\varphi| dx \leq \frac{\|\varphi\|_d |Q_0|}{\lambda} \leq |Q_0|/2.$$

Write  $a_k = \varphi_{Q_k}$ ,  $Q_k \in G_1$ . Then we have

$$\varphi(x) = g_1(x) + \sum_{G_1} a_k \chi_{Q_k}(x) + \varphi_1(x),$$

where  $g_1 = \varphi \chi_{E_1}$ ,  $E_1 = Q_0 \setminus \cup \{Q_k : Q_k \in G_1\}$ , satisfies  $|g_1| \leq \lambda$  by Lebesgue's theorem on differentiating the integral, where  $|a_k| \leq 2^{m+1} \|\varphi\|_d$  by (2.3), and where

$$\varphi_1 = \sum_{G_1} (\varphi(x) - \varphi_{Q_k}) \chi_{Q_k}(x).$$

Now because  $\varphi \in \text{BMO}_d$ ,  $\varphi_1 \chi_{Q_k} = (\varphi - \varphi_{Q_k}) \chi_{Q_k}$  has the same behavior on  $Q_k$  that  $\varphi$  has on  $Q_0$ , and we can repeat the construction with each  $\varphi_1 \chi_{Q_k}$ , and continue by induction. At stage  $n$  we have a family  $G_{n-1}$  of disjoint dyadic cubes and  $\varphi_{n-1} = \sum_{G_{n-1}} (\varphi(x) - \varphi_{Q_j}) \chi_{Q_j}(x)$ . For each  $Q_j \in G_{n-1}$  we set

$$G_1(Q_j) = \{Q_k \subset Q_j : Q_k \text{ dyadic, } |\varphi_{Q_k} - \varphi_{Q_j}| > \lambda, Q_k \text{ maximal}\}$$

and  $G_n = \cup \{G_1(Q_j) : Q_j \in G_{n-1}\}$ . Then

$$\varphi_{n-1}(x) = g_n(x) + \sum_{G_n} a_k \chi_{Q_k}(x) + \varphi_n(x),$$

where  $g_n = \varphi_{n-1} \chi_{E_n}$ ,  $E_n = \bigcup_{G_{n-1}} Q_j \setminus \bigcup_{G_n} Q_k$ , satisfies  $|g_n| \leq \lambda$  and where  $a_k = \varphi_{Q_k} - \varphi_{Q_j}$ ,  $Q_k \subset Q_j \in G_{n-1}$ , satisfies

$$(2.5) \quad |a_k| \leq 2^{m+1} \|\varphi\|_d$$

by the proof of (2.3). Moreover, the proof of (2.4) now gives

$$(2.6) \quad \sum_{\substack{Q_k \subset Q_j \\ Q_k \in G_n}} |Q_k| \leq |Q_j|/2$$

for all  $Q_j \in G_{n-1}$ . Consequently  $\varphi_n(x) \rightarrow 0$  almost everywhere, because  $\varphi_n$  has support  $\bigcup_{G_n} Q_k$  and this set has measure  $\leq 2^{-n} |Q_0|$ . Summing, we obtain

$$\varphi(x) = \sum_{n=1}^{\infty} g_n(x) + \sum_{n=1}^{\infty} \sum_{G_n} a_k \chi_{Q_k}(x).$$

Since  $|g_n| \leq \lambda$  and the  $g_n$  have pairwise disjoint supports,  $g = \sum g_n$  satisfies  $\|g\|_{\infty} \leq \lambda = 2 \|\varphi\|_d$  and we have the representation (2.2).

To prove (2.1) fix a dyadic cube and set  $G_1(Q) = \{Q_j \in \cup G_n : Q_j \subset Q, Q_j \text{ maximal}\}$ . The  $Q_j$  in  $G_1(Q)$  are disjoint and

$$\sum_{Q_k \subset Q} |a_k| |Q_k| = \sum_{Q_j \in G_1(Q)} \sum_{Q \subset Q_j} |a_k| |Q_k|.$$

Hence by (2.5), (2.6) and induction,



$$\begin{aligned} \sum_{Q_k \subset Q} |a_k| |Q_k| &\leq 2^{m+1} \|\varphi\|_d \sum_{Q_j \in G_1(Q)} \sum_{\substack{Q_k \subseteq Q_j \\ Q_k \in \cup G_n}} |Q_k| \\ &\leq 2^{m+1} \|\varphi\|_d \sum_{Q_j \in G_1(Q)} 2|Q_j| \\ &\leq 2^{m+2} \|\varphi\|_d |Q|. \end{aligned}$$

Theorem 2.1 is proved.

Notice that when applied to the translates  $T_\alpha \varphi$ ,  $\varphi \in \text{BMO}$ , the construction above produces functions  $g^{(\alpha)}(x)$  and coefficients  $a_k^{(\alpha)}$  which vary measurably in  $\alpha$ .

Now let  $w \geq 0$  and set  $\varphi = \log w$ . Then  $w \in A_p$ ,  $1 < p < \infty$ , if and only if

$$\sup_Q \left( \frac{1}{|Q|} \int_Q e^{\varphi - \varphi_Q} dx \right) \left( \frac{1}{|Q|} \int_Q e^{-(\varphi - \varphi_Q)/(p-1)} dx \right)^{p-1} < \infty .$$

By Jensen's inequality each factor is at least 1, and hence  $w \in A_p$  if and only if

$$(2.7) \quad \sup_Q \frac{1}{|Q|} \int_Q e^{\varphi - \varphi_Q} dx < \infty$$

and

$$(2.8) \quad \sup_Q \frac{1}{|Q|} \int_Q e^{-(\varphi - \varphi_Q)/(p-1)} dx < \infty .$$

For the dyadic form of Theorem 3, the suprema in (2.7) and (2.8) are taken over dyadic subcubes of  $Q_0$  only.

**THEOREM 2.2.** *Let  $\varphi(x)$  be a real function on a dyadic cube  $Q_0$  and let  $1 < p < \infty$ . Assume*

$$(2.7d) \quad \sup_{\substack{Q \subset Q_0 \\ Q \text{ dyadic}}} \frac{1}{|Q|} \int_Q e^{\varphi - \varphi_Q} dx < \infty ,$$

and

$$(2.8d) \quad \sup_{\substack{Q \subset Q_0 \\ Q \text{ dyadic}}} \frac{1}{|Q|} \int_Q e^{-(\varphi - \varphi_Q)/(p-1)} dx < \infty .$$

Then

$$\varphi - \varphi_Q = g + F - G ,$$

where  $g \in L^\infty$ ,  $\|g\|_\infty \leq C_1$ , where

$$(2.9) \quad \sup_{\substack{Q \subset Q_0 \\ Q \text{ dyadic}}} \left\{ \left( \frac{1}{|Q|} \int_Q e^F dx \right) \|e^{-F}\|_{L^\infty(Q)} \right\} < C_2$$

and where

$$(2.10) \quad \sup_{\substack{Q \subset Q_0 \\ Q \text{ dyadic}}} \left\{ \left( \frac{1}{|Q|} \int_Q e^{G/(p-1)} dx \right) \| e^{-G/(p-1)} \|_{L^\infty(Q)} \right\} < C_3 .$$

The constants  $C_1, C_2, C_3$  depend only on  $m$  and the bounds in (2.7d) and (2.8d).

Thus if  $w = e^\varphi$  satisfies the dyadic  $A_p$  condition (i.e., if 2.7d) and (2.8d), then  $w = w_1(w_2)^{1-p}$  where  $w_1 = e^{\varphi_{Q_0+g+f}}$  and  $w_2 = e^{G/(p-1)}$  satisfy  $A_1$  on dyadic subcubes of  $Q_0$ .

*Proof.* The construction is the same as in the proof of Theorem 2.1. By (2.7d) and (2.8d) and by Jensen's inequality,  $\varphi_{Q_0} \in \text{BMO}_d$ . Fix  $\lambda > 2 \|\varphi\|_d$  to be determined later and set  $G_1 = \{Q_k \subset Q_0: Q_k \text{ dyadic, } |\varphi_{Q_k} - \varphi_{Q_0}| > \lambda, Q_k \text{ maximal}\}$  and by induction

$$G_n = \bigcup_{Q_j \in G_{n-1}} \{Q_k \subset Q_j: Q_k \text{ dyadic, } |\varphi_{Q_k} - \varphi_{Q_j}| > \lambda, Q_k \text{ maximal}\} .$$

For  $Q_k \in G_n, Q_k \subset Q_j \in G_{n-1}$ , set  $a_k = (\varphi_{Q_k} - \varphi_{Q_j})$ . The proof of (2.3) gives

$$(2.11) \quad \lambda < |a_j| < \lambda + 2^m \|\varphi\|_d .$$

As in the proof of Theorem 2.1, we have

$$\varphi = \varphi_{Q_0} + g + \sum_{n=1}^{\infty} \sum_{G_n} a_k \chi_{Q_k}(x) ,$$

where  $\|g\|_\infty \leq \lambda$ . Write

$$(2.12) \quad F = \sum_{a_k > 0} a_k \chi_{Q_k} ,$$

$$(2.13) \quad G = - \sum_{a_k < 0} a_k \chi_{Q_k} .$$

Then  $\varphi = \varphi_{Q_0} + g + F - G$ .

To prove (2.9) and (2.10) we recall that there is  $\varepsilon > 0$ , depending only on the bounds in (2.7d) and (2.8d), such that

$$(2.14) \quad \sup_{\substack{Q \subset Q_0 \\ Q \text{ dyadic}}} \frac{1}{|Q|} \int_Q e^{(1+\varepsilon)(\varphi-\varphi_Q)} dx < \infty ,$$

and

$$(2.15) \quad \sup_{\substack{Q \subset Q_0 \\ Q \text{ dyadic}}} \frac{1}{|Q|} \int_Q e^{-(1+\varepsilon)(\varphi-\varphi_Q)/(p-1)} dx < \infty .$$

See [3] or [10].

We prove (2.9). Fix  $Q_j \in \cup G_n$  with  $a_j > 0$  and set

$$G_1^+(Q_j) = \{Q_k \in \cup G_n : Q_k \subseteq Q_j, a_k > 0, Q_k \text{ maximal}\},$$

and by induction  $G_{n+1}^+(Q) = \cup \{G_1^+(Q_k) : Q_k \in G_n^+(Q_j)\}$ . The critical inequality for the proof of (2.9) is

$$(2.16) \quad \sum_{G_n^+(Q_j)} \frac{|Q_k|}{|Q_j|} \leq (2C)^n e^{-n(1+\epsilon)\lambda},$$

where  $C$  is the suppremmum in (2.14). By induction we need only obtain (2.16) for  $n = 1$ . There are two case.

*Case 1.*  $Q_j \in G_n$  and  $Q_k \in G_{n+1}$ . Then  $\varphi_{Q_k} - \varphi_{Q_j} > \lambda$ , so by Jensen's inequality

$$e^{(1+\epsilon)\lambda} \leq \frac{1}{|Q_k|} \int_{Q_k} e^{(1+\epsilon)(\varphi - \varphi_{Q_j})} dx.$$

Since the  $Q_k$  are disjoint this gives

$$\begin{aligned} \sum_{\text{Case 1}} \frac{|Q_k|}{|Q_j|} &\leq e^{-(1+\epsilon)\lambda} \sum_{\text{Case 1}} \frac{1}{|Q_j|} \int_Q e^{(1+\epsilon)(\varphi - \varphi_{Q_j})} dx \\ &\leq e^{-(1+\epsilon)\lambda} \frac{1}{|Q_j|} \int_Q e^{(1+\epsilon)(\varphi - \varphi_{Q_j})} dx \\ &\leq C e^{-(1+\epsilon)\lambda}. \end{aligned}$$

*Case 2.*  $Q_j \in G_n$  and  $Q_k \in G_{n+p}$ ,  $p \geq 2$ . Then if  $Q_\ell \in G_{n+r}$ ,  $1 \leq r \leq p - 1$ , and if  $Q_k \subset Q_\ell$ , it must be that  $a_\ell < 0$ . Let  $D_1 = \{Q_\ell \in G_{n+1} : Q_\ell \subset Q_j, a_\ell < 0\}$  and by induction  $D_r = \{Q_\ell \in G_{n+r} : \exists Q_m \in D_{r-1}, Q_\ell \subset Q_m, a_\ell < 0\}$ . Then as in the proof of Case 1,  $|\cup D_1| \leq C e^{-(1+\epsilon)\lambda/p-1} |Q_j| \leq 1/2 |Q_j|$  if  $\lambda$  is large enough. Induction then shows  $|\cup D_r| \leq 2^{-r} |Q_j|$ . For  $Q_\ell \in D_r$ ,  $r \geq 1$ , let  $U(Q_\ell) = \{Q_k \in G_{n+r+1} : Q_k \subset Q_\ell, a_k > 0\}$ . By Case 1,  $|\cup U_\ell| \leq C |Q_\ell| e^{-(1+\epsilon)\lambda}$ . Consequently,

$$\begin{aligned} \sum_{\text{Case 2}} \frac{|Q_k|}{|Q_j|} &= \sum_{r=1}^\infty \frac{1}{|Q_j|} \sum_{Q_\ell \in D_r} |\cup U(Q_\ell)| \\ &\leq C e^{-(1+\epsilon)\lambda} \sum_{r=1}^\infty \frac{1}{|Q_j|} \sum_{Q_\ell \in D_r} |Q_\ell| \\ &\leq C e^{-(1+\epsilon)\lambda} \sum_{r=1}^\infty 2^{-r}, \end{aligned}$$

because the cubes  $Q_\ell \in D_r$  are disjoint for each  $r$ . Summing the two cases gives (2.16) for  $n = 1$ .

Now fix a dyadic cube  $Q \subset Q_0$  and set

$$F_1 = \sum_{\substack{Q_k \subseteq Q \\ a_k > 0}} a_k \chi_{Q_k}, \quad F_2 = \sum_{\substack{Q_k \supseteq Q \\ a_k > 0}} a_k \chi_{Q_k}.$$

On  $Q$ ,  $F = F_1 + F_2$ ,  $F_2$  is constant, and  $F_1 \geq 0$ . Hence

$$\begin{aligned} \left(\frac{1}{|Q|} \int_Q e^F dx\right) \|e^{-(F_1+F_2)}\|_{L^\infty(Q)} &= \left(\frac{1}{|Q|} \int_Q e^{F_1+F_2} dx\right) \|e^{-(F_1+F_2)}\|_{L^\infty(Q)} \\ &\leq \frac{1}{|Q|} \int_Q e^{F_1} dx, \end{aligned}$$

and it suffices to establish

$$(2.17) \quad \frac{1}{|Q|} \int_Q e^{F_1} dx \leq C_2.$$

If  $\{Q_j\}$  denotes the set of maximal cubes  $Q_j \subset Q$  having  $a_j > 0$ , then  $\sum |Q_j| \leq |Q|$  and

$$\frac{1}{|Q|} \int_Q (e^{F_1} - 1) dx \leq \sum \frac{|Q_j|}{|Q|} \left(\frac{1}{|Q_j|} \int_{Q_j} e^{F_1} dx\right) \leq \sup_j \left(\frac{1}{|Q_j|} \int_{Q_j} e^{F_1} dx\right).$$

Now by (2.11),

$$\{x \in Q: F_1(x) > (n+1)(\lambda + 2^m \|\varphi\|_d)\} \subset \bigcup_{G_n^+(Q_j)} Q_k,$$

so that by (2.16),

$$\frac{1}{|Q_j|} \int_{Q_j} e^{F_1} dx \leq \sum_{n=0}^\infty e^{(n+1)(\lambda+2^m\|\varphi\|_d)} (2C)^n e^{-n(1+\varepsilon)\lambda}.$$

If  $\lambda > \lambda(\varepsilon, C, \|\varphi\|_d)$  the series sums and we obtain (2.17) and therefore (2.9).

The proof of (2.11) is the same except that (2.15) is used in place of (2.14).

**3. The proof of Theorem 3.** Let  $w = e^\varphi \in A_p$  and let  $S_N$  be the cube  $\{|x_i| \leq 2^N, 1 \leq i \leq m\}$ .

**LEMMA 3.1.** *There exist  $g_N(x), F_N(x)$  and  $G_N(x), x \in S_N$ , such that  $\|g_N\|^\infty \leq C_1$ , and*

$$(3.1) \quad \sup_{Q \subset S_N} \left(\frac{1}{|Q|} \int_Q e^{F_N} dx\right) \|e^{-F_N}\|_{L^\infty(Q)} \leq C_2,$$

$$(3.2) \quad \sup_{Q \subset S_N} \left(\frac{1}{|Q|} \int_Q e^{G_N/p-1} dx\right) \|e^{-G_N/p-1}\|_{L^\infty(Q)} \leq C_3,$$

and such that

$$\varphi(x) - \varphi_{S_N} = g_N(x) + F_N(x) - G_N(x), \quad x \in S_N.$$

The constants  $C_1, C_2, C_3$  do not depend on  $N$ .

We first show how Lemma 3.1 easily implies Theorem 3. We suppose  $\varphi_{S_0} = 0$ . By Lemma 3.1,

$$\begin{aligned} \varphi &= R_N + (F_N - (F_N)_{S_0}) - (G_N - (G_N)_{S_0}) \\ &= R_N + \tilde{F}_N - \tilde{G}_N, \quad x \in S_N, \end{aligned}$$

where  $R_N = \varphi_{S_N} + g_N + (F_N)_{S_0} - (G_N)_{S_0}$  satisfies  $\|R_N\|_\infty \leq 2C_1$  since  $|\varphi_{S_N} + (F_N)_{S_0} - (G_N)_{S_0}| = |\varphi_{S_0} - (g_N)_{S_0}| \leq C_1$ . For  $N > M$ , (3.1) and (3.2) give

$$\begin{aligned} \frac{1}{|S_M|} \int_{S_M} |\tilde{F}_N - (\tilde{F}_N)_{S_M}|^2 dx &\leq C \\ \frac{1}{|S_M|} \int_{S_M} |\tilde{G}_N - (\tilde{G}_N)_{S_M}|^2 dx &\leq C, \end{aligned}$$

and hence as  $(\tilde{F}_N)_{S_0} = (\tilde{G}_N)_{S_0} = 0$ ,  $|(\tilde{F}_N)_{S_M}| \leq C_M$ ,  $|(\tilde{G}_N)_{S_M}| \leq C_M$ . Consequently  $\{(\tilde{F}_N: N \geq M)\}$  and  $\{(\tilde{G}_N: N \geq M)\}$  are bounded in  $L^2(S_M)$ . Choose  $N_j \rightarrow \infty$  so that  $\tilde{F}_{N_j} \rightarrow F$ ,  $\tilde{G}_{N_j} \rightarrow G$  weakly in  $L^2(S_M)$  for all  $M$  and so that  $R_{N_j} \rightarrow g$  weak-star in  $L^\infty$ . Then

$$\varphi = g + F - G$$

with  $\|g\|_\infty \leq C_1$ . For any cube  $Q$  there is a sequence of finite convex combinations

$$F^{(n)} = \sum_j t_{j,n} F_{N_{j,n}}, \quad t_{j,n} \geq 0, \quad \sum t_{j,n} = 1,$$

converging to  $F$  almost everywhere on  $S_M \subset Q$ . Then by Fatou's Lemma and Hölder's inequality

$$\left(\frac{1}{|Q|} \int_Q e^F dx\right) \|e^{-F}\|_{L^\infty(Q)} \leq \lim_{n \rightarrow \infty} \Pi \left(\frac{1}{|Q|} \int_Q e^{F_{N_{j,n}}} dx\right)^{t_{j,n}} \Pi \|e^{-F_{N_{j,n}}}\|^{t_{j,n}} \leq C_2,$$

and hence  $w_1 = e^{g+F} \in A_1$ . Using (3.2) we see  $w_2 = e^{G/p-1} \in A_1$  in the same way.

*Proof of Lemma 3.1.* We assume  $\varphi_{S_N} = 0$ . For  $\alpha \in S_N$  we use Theorem 2.2 on  $T_\alpha \varphi(x) = \varphi(x - \alpha)$  with  $Q_0 = S_{N+1}$  (which we pretend is a dyadic cube) to obtain

$$T_\alpha \varphi = g^{(\alpha)} + F^{(\alpha)} - G^{(\alpha)}$$

where  $F_\alpha$  and  $G_\alpha$  satisfy (2.9) and (2.10) respectively and where  $\|g\|_\infty \leq C_1$  (since  $\varphi \in \text{BMO}$  and  $\varphi_{S_N} = 0$ ,  $\sup_{\alpha \in S_N} (T_\alpha \varphi)_{S_{N+1}}$  is bounded). Almost everywhere on  $S_N$ ,

$$\begin{aligned} \varphi(x) &= \frac{1}{|S_N|} \int_{S_N} T_{-\alpha}(T_\alpha \varphi)(x) d\alpha \\ &= \frac{1}{|S_N|} \int_{S_N} T_{-\alpha}(g^{(\alpha)})(x) d\alpha + \frac{1}{|S_N|} \int_{S_N} T_{-\alpha}(F^{(\alpha)})(x) d\alpha \\ &\quad + \frac{1}{|S_N|} \int_{S_N} T_{-\alpha}(G^{(\alpha)})(x) d\alpha = g(x) + F(x) - G(x). \end{aligned}$$

Clearly  $\|g\|_\infty \leq C_1$ . By (2.12) there are  $\alpha_k^{(\alpha)} > 0$  such that

$$\begin{aligned}
 F^{(\alpha)}(x) &= \sum_{n=0}^{\infty} \sum_{\ell(Q_k)=2^{-n}\ell(S_N)} a_k^{(\alpha)} \chi_{Q_k}(x) \\
 &= \sum_{n=0}^{\infty} f_n^{(\alpha)}(x)
 \end{aligned}$$

and by (2.13),  $G^{(\alpha)}(x)$  has a similar representation. Write

$$f_n(x) = \frac{1}{|S_N|} \int_{S_N} (T_{-\alpha} f_n^{(\alpha)})(x) d\alpha$$

so that  $F = \sum_{n=0}^{\infty} f_n$ .

LEMMA 3.2. *If  $\sup_i |x_i - y_i| \leq 2^{-n}\ell(S_N)$  then*

$$(3.3) \quad |f_n(x) - f_n(y)| \leq \frac{C_4 2^n}{\ell(S_N)} |x - y|,$$

with  $C_4$  independent of  $n$ .

*Proof.* By (2.11),  $|a^{(\alpha)}| \leq C$ , and hence

$$|f_n(x) - f_n(y)| \leq \frac{C}{|S_N|} \int_{S_N} \sum_{\ell(Q_k)=2^{-n}\ell(S_N)} |\chi_{Q_k}(x + \alpha) - \chi_{Q_k}(y + \alpha)| d\alpha.$$

The integrand is twice the characteristic function of  $\{\alpha \in S_N: x + \alpha$  and  $y + \alpha$  fall in different  $Q_k, \ell(Q_k) = 2^{-n}\ell(S_N)\}$ , and this set has probability not exceeding

$$\frac{\sum_{i=1}^m |x_i - y_i|}{2^{-n}\ell(S_N)}$$

Returning to the proof of Lemma 3.1, we fix  $Q \subset S_N$  with  $2^{-k}\ell(S_N) < \ell(Q) \leq 2^{-k+1}\ell(S_N)$ . Then

$$\begin{aligned}
 F(x) &= \sum_{n>k} f_n(x) + \sum_{n\leq k} f_n(x) = F_1(x) + F_2(x) \\
 &= \frac{1}{|S_N|} \int_{S_N} F_1^{(\alpha)}(x + \alpha) d\alpha + \frac{1}{|S_N|} \int_{S_N} F_2^{(\alpha)}(x + \alpha) d\alpha.
 \end{aligned}$$

By Lemma 3.2,

$$\sup_Q F_2(x) - \inf_Q F_2(x) \leq \frac{C}{\ell(S_N)} \sum_{n=0}^k 2^n \ell(Q) \leq C.$$

Hence as  $F_1 \geq 0$ ,

$$\left( \frac{1}{|Q|} \int_Q e^{F_1+F_2} dx \right) \| e^{-(F_1+F_2)} \|_{L^\infty(Q)} \leq C \left( \frac{1}{|Q|} \int_Q e^{F_1} dx \right).$$

But by Jensen's inequality,

$$\begin{aligned} \frac{1}{|Q|} \int_Q e^{F_1(x)} dx &= \frac{1}{|Q|} \int_Q \exp\left(\frac{1}{|S_N|} \int_{S_N} F_1^{(\alpha)}(x + \alpha) d\alpha\right) dx \\ &\leq \frac{1}{|S_N|} \int_{S_N} \left(\frac{1}{|Q|} \int_Q e^{F_1^{(\alpha)}(x+\alpha)} dx\right) d\alpha \leq C_2. \end{aligned}$$

by (2.17). The proof of (3.2) is the same.

4. **The proof of Theorem 1.** We suppose  $\varphi \in \text{BMO}$  has support  $S_0 = \{|x_i| \leq 1, 1 \leq i \leq m\}$  and  $\int \varphi dx = 0$ . For each  $\alpha \in S_0$  we have, by Theorem 2.1,

$$T_\alpha \varphi(x) = \varphi(x - \alpha) = g^{(\alpha)}(x) + \sum_{Q_k \subset Q_0} a_k^{(\alpha)} \chi_{Q_k}(x)$$

where  $Q_0 = \{|x_i| \leq 2, 1 \leq i \leq m\}$ , where  $\|g^{(\alpha)}\|_\infty \leq C\|\varphi\|$ , and where

$$(4.1) \quad \sum_{Q_k \subset Q} |a_k^{(\alpha)}| |Q_k| \leq C\|\varphi\| |Q|.$$

Write

$$f_n^{(\alpha)}(x) = \sum_{\mathcal{L}(Q) = 2^{-n}} a_k^{(\alpha)} \chi_{Q_k}(x),$$

so that  $T_\alpha \varphi(x) = g^{(\alpha)}(x) + \sum_{n=0}^\infty f_n^{(\alpha)}(x)$ . Then as before

$$\begin{aligned} \varphi(x) &= \frac{1}{|S_0|} \int_{S_0} g^{(\alpha)}(x + \alpha) d\alpha + \sum_{n=0}^\infty \frac{1}{|S_0|} \int_{S_0} f_n^{(\alpha)}(x + \alpha) d\alpha \\ &= g(x) + \sum_{n=0}^\infty f_n(x) \end{aligned}$$

where  $\|g\|_\infty \leq C\|\varphi\|$ . For any cube  $Q$  we have

$$(4.2) \quad \begin{aligned} \frac{1}{|Q|} \int_{Q_{2^{-n} \leq \mathcal{L}(Q)}} |f_n(x)| dx &\leq \sup_{\alpha \in S_0} \left( \frac{1}{|Q|} \int_{Q_{2^{-n} < \mathcal{L}(Q)}} |f_n^{(\alpha)}(x)| dx \right) \\ &\leq \sup_{\alpha \in S_0} \left( \frac{1}{|Q|} \sum_{Q_k \subset Q} |a_k^{(\alpha)}| |Q_k| \right) \leq C\|\varphi\|, \end{aligned}$$

where  $\tilde{Q}$  is concentric with  $Q$  and  $\mathcal{L}(\tilde{Q}) = 3\mathcal{L}(Q)$ . Thus for any  $\delta > 0$ ,  $d\sigma = \sum f_n(x) d\sigma_n$ , where  $d\sigma_n$  is surface measure on  $\mathbf{R}^m \times \{y = \delta 2^{-n}\}$ , is a Carleson measure and  $N(\sigma) \leq C\delta^{-m} \|\varphi\|$ , and

$$\begin{aligned} \int K_y(x - z) d\sigma(z, y) &= \sum_{n=0}^\infty f_n * K_{\delta 2^{-n}}(x) \\ &= \sum h_n(x). \end{aligned}$$

We will show that when  $\delta$  is small,

$$(4.3) \quad \left\| \sum (f_n - h_n) \right\| \leq \frac{1}{2} \|\varphi\|.$$

With an iteration, that will prove Theorem 1.

To prove (4.3) fix a cube  $Q$  and a point  $x_0 \in Q$ . We have

$$\begin{aligned} & \frac{1}{|Q|} \int_Q |\sum (f_n(x) - h_n(x) - (f_n - h_n)_Q)| dx \\ & \leq 2 \sum_{2^{-n} \geq A \ell(Q)} \frac{1}{|Q|} \int_Q |(f_n - h_n)(x) - (f_n - h_n)(x_0)| dx \\ & \quad + 2 \sum_{\ell(Q) \leq 2^{-n} < A \ell(Q)} \frac{1}{|Q|} \int_Q |f_n(x) - h_n(x)| dx \\ & \quad + 2 \sum \frac{1}{|Q|} \int_Q |f_n(x) - h_n(x)| dx \\ & = 2 \sum_1 + 2 \sum_2 + 2 \sum_3, \end{aligned}$$

where  $A \geq 2$  is a constant to be determined.

To estimate  $\sum_1$ , recall that

$$(4.4) \quad |f_n(x) - f_n(y)| \leq C2^n \|\varphi\| |x - y|$$

by the proof of Lemma 3.2. The convolution  $h_n = K_{\delta 2^{-n}} * f_n$  has the same continuity as  $f_n$ , since  $\int K dx = 1$ , and we have

$$\begin{aligned} \sum_1 & \leq \sum_{2^{-n} > A \ell(Q)} \frac{1}{|Q|} \int_Q 2C2^n \|\varphi\| |x - x_0| dx \leq C' \|\varphi\| \ell(Q) \sum_{2^n \leq (A \ell(Q))^{-1}} 2^n \\ & \leq C' \|\varphi\| / A. \end{aligned}$$

Hence  $2 \sum_1 \leq \|\varphi\| / 6$  if  $A$  is large.

To estimate  $\sum_2$ , note that by (4.4) and the bound  $\|f_n\|_\infty \leq C \|\varphi\|$  (because  $|a_k^{(\alpha)}| \leq C \|\varphi\|$ ), we have

$$\|f_n - f_n * K_{\delta 2^{-n}}\|_\infty \leq \varepsilon \|\varphi\|$$

if  $\delta$  is small, independent of  $n$ . Therefore

$$2 \sum_2 \leq 2\varepsilon \|\varphi\| \sum_{\ell(Q) \leq 2^{-n} \leq A \ell(Q)} \leq C\varepsilon \|\varphi\| \log A \leq \|\varphi\| / 6$$

if  $\varepsilon \log A$  is small.

Finally, we have

$$\sum_3 \leq \sup_\alpha \frac{1}{|Q|} \int_{Q_{2^{-n} < \ell(Q)}} |f_n^{(\alpha)} - f_n^{(\alpha)} * K_{\delta 2^{-n}}^{(\alpha)}| dx$$

by the definition of  $f_n$ . After a translation it is enough to consider  $\alpha = 0$ . Let  $Q^{(0)} = Q$  and pave  $R^m$  with cubes  $Q^{(j)}$  congruent to  $\tilde{Q}$ . Then

$$\sum_3 \leq \sum_j \sum_{Q_k \subset Q^{(j)}} |a_k^{(0)}| |Q_k| \frac{1}{|Q|} \int_Q \left| \frac{\chi_{Q_k}(x)}{|Q_k|} - \frac{\chi_{Q_k} * K_{\delta \ell(Q_k)}(x)}{|Q_k|} \right| dx$$



By a change of scale,

$$\frac{1}{|Q|} \int_{\mathbf{R}^m} \left| \frac{\chi_{Q_k}(x)}{|Q_k|} - \frac{\chi_{Q_k} * K_{\delta \ell(Q_k)}(x)}{|Q_k|} \right| dx$$

does not depend on  $\ell(Q_k)$ . Thus for  $\varepsilon > 0$  we can choose  $\delta$  so that

$$\frac{1}{|Q|} \int_Q \left| \frac{\chi_{Q_k}(x)}{|Q_k|} - \frac{\chi_{Q_k} * K_{\delta \ell(Q_k)}(x)}{|Q_k|} \right| dx < \varepsilon / |Q|.$$

Moreover, if  $Q_k \subset Q^{(j)}$ ,  $Q^{(j)} \neq Q^{(0)}$ , then  $\chi_{Q_k}(x) = 0$  on  $Q$  and by (1.1),

$$\begin{aligned} \frac{1}{|Q|} \int_Q \frac{\chi_{Q_k} * K_{\delta \ell(Q_k)}(x)}{|Q_k|} dx &\leq \sup_{\substack{x \in Q \\ x \in Q_k}} K_{\delta \ell(Q_k)}(x - t) \leq \frac{C\delta \ell(Q_k)}{(\text{dist}(Q_k, Q))^{m+1}} \\ &\leq \frac{C\delta \ell(Q)}{(\text{dist}(Q^{(j)}, Q))^{m+1}} \end{aligned}$$

since  $\text{dist}(Q_k, Q) \geq \text{dist}(Q^{(j)}, Q)$ . Therefore

$$\begin{aligned} \sum_3 &\leq \varepsilon \sum_{Q_k \subset Q^{(0)}} \frac{|a_k^{(0)}| |Q_k|}{|Q|} \\ &\quad + C\delta \ell(Q) \sum_{j \neq 0} \frac{1}{(\text{dist}(Q^{(j)}, Q))^{m+1}} \sum_{Q_k \subset Q^{(j)}} |a_k^{(0)}| |Q_k|, \end{aligned}$$

and by (4.1),

$$\begin{aligned} 2 \sum_3 &\leq C\varepsilon \|\varphi\| + C\delta \|\varphi\| \ell(Q) \sum_{j \neq 0} \frac{|Q^{(j)}|}{(\text{dist}(Q^{(j)}, Q))^{m+1}} \\ &\leq C\varepsilon \|\varphi\| + C\delta \|\varphi\| \ell(Q) \int_{\mathbf{R}^m / Q^{(0)}} \frac{dx}{|x - x_0|^{m+1}} \\ &\leq C(\varepsilon + \delta) \|\varphi\| \leq \|\varphi\| / 6 \end{aligned}$$

if  $\varepsilon$  and  $\delta$  are small.

**5. The proof of Theorem 4.** We begin with the dyadic form of the theorem, which is also due to Uchiyama.

**THEOREM 5.1.** *Let  $\lambda > 0$ , let  $Q_0$  be a dyadic cube in  $\mathbf{R}^m$ , and let  $E_1, E_2, \dots, E_N$  be measurable subsets of  $Q_0$  such that*

$$(5.1) \quad \text{Min}_{1 \leq i \leq N} \frac{|Q \cap E_i|}{|Q|} \leq 2^{-2m\lambda}$$

for all dyadic  $Q \subset Q_0$ . Then there exist  $f_1(x), f_2(x), \dots, f_N(x)$  such that almost everywhere on  $Q_0$ ,

$$(5.2) \quad f_i(x) = 0, x \in E_i,$$

$$(5.3) \quad 0 \leq f_i(x) \leq 1 ;$$

$$(5.4) \quad \sum_{i=1}^N f_0(x) = 1 ;$$

and

$$(5.5) \quad \sup_{\substack{Q \subset Q_0 \\ Q \text{ dyadic}}} \frac{1}{|Q|} \int_Q |f_i - (f_i)_Q| dx \leq C_1(m, N)/\lambda .$$

*Proof.* By (5.1),  $|\cap E_i| = 0$  and the bounded solutions  $f_i(x) = (1 - \chi_{E_i}(x))/\sum_j(1 - \chi_{E_j}(x))$  satisfy (5.5) if  $\lambda$  is not large. Thus we assume  $\lambda > N$ .

We shall inductively choose families  $G_n$  of dyadic cubes  $Q_k \subset Q_0$  and functions  $\psi_i^{(n)}, 1 \leq i \leq N$  such that

$$(5.6) \quad \psi_i^{(n)} = \psi_i^{(n-1)} + \sum_{Q_k \in G_n} a_{i,k} \chi_{Q_k}(x) ,$$

$$0 \leq \psi_i^{(n)} \leq \lambda ,$$

$$(5.7) \quad \sum_i \psi_i^{(n)} = \lambda ,$$

$$(5.8) \quad (\psi_i^{(n)})_{Q_k} \leq \text{Max}\left(0, -N + \frac{1}{m} \log_2\left(\frac{|Q_k|}{|Q_k \cap E_i|}\right)\right)$$

if  $Q_k \in G_n$ , and

$$(5.9) \quad |a_{i,k}| \leq N^2 - 1, Q_k \in G_n, n \geq 1 .$$

For each dyadic cube  $Q \subset Q_0$ , (5.1) ensures there exists an index  $i(Q), 1 \leq i(Q) \leq N$ , such that

$$(5.10) \quad 2\lambda \leq \frac{1}{m} \log_2\left(\frac{|Q_k|}{|Q \cap E_{i(Q)}|}\right)$$

To start the induction take  $G_0 = \{G_0\}$ , and  $\psi_i^{(0)}(x) = a_{i,0} \chi_{Q_0}(x)$ , where

$$a_{i,0} = \begin{cases} 0, & i \neq i(Q_0) \\ \lambda, & i = i(Q_0) . \end{cases}$$

Then (5.6) and (5.7) are trivial and (5.8) follows from (5.10) and our choice  $\lambda > N$ . At  $n = 0$ , (5.9) is not required.

Let  $G_n$  be the set of maximal dyadic cubes satisfying  $Q_k \subset Q_j \in G_{n-1}$  and

$$(5.11) \quad (\psi_i^{(n-1)})_{Q_k} > \frac{1}{m} \log_2\left(\frac{|Q_k|}{|Q_k \cap E_i|}\right) ,$$

for some  $i$ ,  $1 \leq i \leq N$ . Define

$$a_{i,k} = \begin{cases} -\text{Min}(N + 1, (\psi_i^{(n-1)})_{Q_k}), & i \neq i(Q_k) \\ -\sum_{j \neq i(Q_k)} a_{j,k}, & i = i(Q_k). \end{cases}$$

Then by definition  $\psi_i^{(n)} = \psi_i^{(n-1)} + \sum_{Q_k \in G_n} a_{i,k} \chi_{Q_k}$  clearly satisfies  $\psi_i^{(n)} \geq 0$  and  $\sum_i \psi_i^{(n)} = \lambda$ . Thus (5.6) and (5.7) hold. Since  $|a_{i,k}| \leq N + 1$  for  $i \neq i(Q_k)$ , and since  $|a_{i(Q_k),k}| \leq (N - 1)(N + 1)$ , (5.9) holds.

We now verify inequality (5.8). If  $i = i(Q_k)$ , then by (5.6) and (5.10),

$$(\psi_i^{(n)})_{Q_k} \leq \lambda \leq -N + \frac{1}{m} \log_2 \left( \frac{|Q_k|}{|Q_k \cap E_i|} \right).$$

Suppose  $Q_k \in G_n$  and  $i \neq i(Q_k)$ . If  $Q_k^* \supset Q_k$  is that dyadic cube with  $|Q_k^*| = 2^m |Q_k|$ , then  $(\psi_i^{(n-1)})_{Q_k^*} = (\psi_i^{(n-1)})_{Q_k}$  and

$$\log_2 \left( \frac{|Q_k^*|}{|Q_k^* \cap E_i|} \right) \leq m + \log_2 \left( \frac{|Q_k|}{|Q_k \cap E_i|} \right).$$

Since  $Q_k$  is maximal, (5.11) fails for  $Q_k^*$ , and so we have

$$(5.12) \quad 1 = \frac{1}{m} \log_2 \left( \frac{|Q_k|}{|Q_k \cap E_i|} \right) \geq (\psi_i^{(n-1)})_{Q_k}.$$

If  $a_{i,k} = -(\psi_i^{(n-1)})_{Q_k}$ , then  $(\psi_i^{(n)})_{Q_k} = 0$  and (5.9) is clear. If  $a_{i,k} = -(N + 1)$ , then (5.9) follows from (5.12). Thus the induction is completed.

We thank J. Michael Wilson for this argument.

To obtain convergence and ultimately (5.5) we observe that if  $Q_j \in G_{n-1}$ , then

$$(5.13) \quad \sum_{\substack{Q_k \in G_n^{(i)} \\ Q_k \subset Q_j}} |Q_k| \leq 2^{-mN} |Q_j|.$$

Indeed, if the left side of (5.13) is nonzero, we have  $(\psi_i^{(n-1)})_{Q_j} > 0$ , and then (5.11) and (5.8) yield

$$\begin{aligned} \sum_{\substack{Q_k \in G_n^{(i)} \\ Q_k \subset Q_j}} |Q_k| &\leq 2^{m(\psi_i^{(n-1)})_{Q_j}} \sum_{\substack{Q_k \in G_n^{(i)} \\ Q_k \subset Q_j}} |Q_k \cap E_i| \\ &\leq 2^{m(\psi_i^{(n-1)})_{Q_j}} |Q_j \cap E_i| \\ &\leq 2^{-mN} |Q_j|. \end{aligned}$$

Since  $N2^{-mN} < 1$ , (5.9), (5.13) and induction show  $\sum \|\psi_i^{(n)} - \psi_i^{(n-1)}\|_1 < \infty$ , so that

$$\psi_i(x) = \lim_n \psi_i^{(n)}(x)$$

exists almost everywhere. Moreover, if  $Q \subset Q_0$  is a dyadic cube, then by (5.9) and (5.13)

$$\begin{aligned}
 (5.19) \quad & \frac{1}{|Q|} \int_Q |\psi_i - (\psi_i)_Q| dx \\
 & \leq \frac{2}{|Q|} \int_Q \sum_{Q_k \equiv Q} |a_{i,k}| \chi_{Q_k}(x) dx \\
 & \leq 2(N^2 - 1) \sum_{\substack{Q_k \equiv Q \\ Q_k \in \cup G_n}} |Q_k|/|Q| \leq 2(N^2 - 1) \sum_{\ell=1}^{\infty} (N2^{-mN})^\ell \\
 & = C_1(m, N).
 \end{aligned}$$

Write  $f_i = \psi_i/\lambda$ . Then (5.5) follows from (5.14) and (5.6) and (5.7) give (5.3) and (5.4).

To conclude the proof we establish (5.2). Almost every point  $x \in Q_0$  lies in a unique dyadic cube  $Q_k(x)$ ,  $|Q_k| = 2^{-mk}$ ,  $k = 0, 1, 2, \dots$ . For almost such  $x$ ,  $Q_k(x) \in \cup G_n$  for only finitely  $k$ , because by (5.13),  $\sum_n |\cup \{Q_k: Q_k \in G_n\}| < \infty$ . Hence for almost every  $x$  there exist  $k_x < \infty$  and  $n_x < 8$  such that for  $k > k_x$  and  $n > n_x$

$$Q_k(x) \notin G_n$$

and

$$\psi_i(x) = \psi_i^{(n)}(x).$$

So by the definition of  $G_n$ ,

$$\psi_i(x) = (\psi_i^{(n-1)})_{Q_k(x)} \leq \frac{1}{m} \log_2 \left( \frac{|Q_k(x)|}{|Q_k(x) \cap E_i|} \right)$$

$k > k_x, n > n_x$ , almost all  $x$ . On the other hand,

$$\log_2 \left( \frac{|Q_k(x)|}{|Q_k(x) \cap E_i|} \right) \longrightarrow 0 \quad (k \longrightarrow \infty),$$

almost everywhere on  $E_i$ . Therefore  $f_i(x) = \psi_i(x) = 0$  almost everywhere on  $E_i$ .

*Proof of Theorem 4.* The argument is much like the proof of Theorem 3. Let  $S_M$  be the cube  $\{x: |x_i| \leq 2^M\}$ . It is enough to produce  $f_{1,M}(x), \dots, f_{N,M}(x)$  which satisfy (1.6), (1.7) and (1.8) for  $x \in S_M$  and also

$$(5.15) \quad \sup_{Q \subset S_M} \frac{1}{|Q|} \int_Q |f_{i,M} - (f_{i,M})_Q| dx \leq C(m, N)/\lambda,$$

by then taking  $f_i(x)$  an  $L^\infty$  weak-star limit of  $\{f_{1,M}(x)\}_{M=1}^\infty$ .

So fix  $S_M$ . For  $\alpha \in S_M$  we set  $E_i^{(\alpha)} = \{x + \alpha : x \in E_i \cap S_M\} \subset S_{M+1}$ . With  $Q_0 = S_{M+1}$ , (5.1) holds for  $E_1^{(\alpha)}, \dots, E_N^{(\alpha)}$ , and Theorem 5.1 gives us  $f_1^{(\alpha)}(x), \dots, f_N^{(\alpha)}(x)$  satisfying (5.2), (5.3), (5.4) and (5.5) on  $S_{M+1}$ . Define, as before,

$$f_{i,M}(x) = \frac{1}{|S_M|} \int f_i^{(\alpha)}(x + \alpha) d\alpha, \quad x \in S_M.$$

Then (1.6), (1.7) and (1.8) hold on  $S_M$ . To prove (5.15) write

$$\begin{aligned} f_i^{(\alpha)}(x) &= \frac{1}{\lambda} \sum_{n=0}^{\infty} \sum_{\ell(Q_k)=2^{-n}\ell(S_{M+1})} a_{i,k}^{(\alpha)} \chi_{Q_k}(x) \\ &= \sum_{n=0}^{\infty} f_{i,n}^{(\alpha)}(x) \end{aligned}$$

and

$$\begin{aligned} f_{i,M}(x) &= \sum_{n=0}^{\infty} \frac{1}{|S_M|} \int f_{i,n}^{(\alpha)}(x + \alpha) d\alpha \\ &= \sum_{n=0}^{\infty} f_{i,n}(x). \end{aligned}$$

If  $Q \subset S_M$  and  $2^{-k}\ell(S_{M+1}) < \ell(Q) \leq 2^{-k+1}\ell(S_{M+1})$ , then

$$\begin{aligned} \frac{1}{|Q|} \int_Q |f_{i,M}(x) - (f_{i,M})_Q| dx &\leq \sum_{n < k} \frac{1}{|Q|} \int_Q |f_{i,n}(x) - (f_{i,n})_Q| dx \\ &\quad + 2 \sum_{n \geq k} \frac{1}{|Q|} \int_Q |f_{i,n}(x)| dx = \sum_1 + 2 \sum_2. \end{aligned}$$

By the proof of Lemma 3.2

$$|f_{i,n}(x) - f_{i,n}(y)| \leq \frac{C(N)2^n}{\lambda\ell(S_{M+1})} |x - y|,$$

so that  $\sum_1 \leq C(N)/\lambda$ , and by (5.13) and (5.9),

$$\begin{aligned} \sum_2 &\leq \sup_{\alpha \in S_M} \sum_{Q_k \subset Q + \alpha} |a_{i,k}^{(\alpha)}| |Q| \\ &\leq \frac{C(N^2 - 1)}{\lambda} |Q|. \end{aligned}$$

Hence (5.15) holds and Theorem 4 is proved.

REFERENCES

1. L. Carleson *Two remarks on  $H^1$  and BMO*, Advances in Math., **22** (1976) 269-277.
2. R. R. Coifman and C. Fefferman, *Weighted norm inequalities for maximal functions and singular integrals*, Studia Math. **LI** (1974), 241-250.
3. R. R. Coifman and R. Rochberg, *Another characterization of BMO*, Proc. Amer. Math. Soc., **79** (1980), 249-254.

4. B. Davis, *Hardy spaces and rearrangements*, Trans. Amer. Math. Soc., **261** (1980), 211-233.
5. C. Fefferman and E. M. Stein,  *$H^p$  spaces of several real variables*, Acta Math., **129** (1972), 137-193.
6. J. B. Garnett and P. W. Jones *The distance in BMO to  $L^\infty$* , Annals of Math., **108** (1978), 373-393.
7. F. John and L. Nirenberg, *On function of bounded mean oscillation*, Comm. Pure Appl. Math., **14** (1961), 415-426.
8. P. Jones, *Estimates for the corona problem*, to appear in J. Functional Analysis.
9. ———, *Factorization of  $A_p$  weights*, Annals of Math., **111** (1980), 511-530.
10. B. Muckenhoupt, *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc., **165** (1972), 207-226.
11. A. Uchiyama, *The construction of certain BMO functions and the corona problem*, to appear.
12. N. Th. Varopoulos, *A probabilistic proof of the Garnett-Jones theorem on BMO*, to appear in Pacific J. Math.
13. N. Th. Varopoulos, *BMO functions and the  $\bar{\partial}$  equation*, Pacific J. Math., **71** (1977), 221-273.

Received November 19, 1980. The first author was partially supported by NSF Grant MCS80-02955 and the second by NSF Grant MCS810-2631.

UNIVERSITY OF CALIFORNIA  
LOS ANGELES, CA 90024  
AND  
UNIVERSITY OF CHICAGO  
CHICAGO, IL 60637



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California  
Los Angeles, CA 90024

HUGO ROSSI

University of Utah  
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR AGUS

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University  
Stanford, CA 94305

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$102.00 a year (6 Vols., 12 issues). Special rate: \$51.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1982 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan



<b>Thomas E. Armstrong and Karel Libor Prikry</b> , On the semimetric on a Boolean algebra induced by a finitely additive probability measure . . . . .	249
<b>Walter Russell Bloom</b> , Strict local inclusion results between spaces of Fourier transforms . . . . .	265
<b>Richard Clark Brown</b> , Notes on generalized boundary value problems in Banach spaces. II. Infinite-dimensional extension theory . . . . .	271
<b>Sui Sun Cheng</b> , Isoperimetric eigenvalue problem of even order differential equations . . . . .	303
<b>Lung O. Chung and Jiang Luh</b> , Derivations of higher order and commutativity of rings . . . . .	317
<b>Ali Ahmad Fora</b> , A fixed point theorem for product spaces . . . . .	327
<b>Barry J. Gardner</b> , Radical classes of regular rings with Artinian primitive images . . . . .	337
<b>John Brady Garnett and Peter Wilcox Jones</b> , BMO from dyadic BMO . . . . .	351
<b>Allen E. Hatcher</b> , On the boundary curves of incompressible surfaces . . . . .	373
<b>Richard Howard Hudson and Kenneth S. Williams</b> , Resolution of ambiguities in the evaluation of cubic and quartic Jacobsthal sums . . . . .	379
<b>Viktor Losert</b> , Counter-examples to some conjectures about doubly stochastic measures . . . . .	387
<b>Kenneth Derwood Magill, Jr., P. R. Misra and Udai Bhan Tewari</b> , Structure spaces for sandwich semigroups . . . . .	399
<b>Mark Mandelker</b> , Continuity of monotone functions . . . . .	413
<b>Kenneth Guy Miller</b> , An index theorem and hypoellipticity on nilpotent Lie groups . . . . .	419
<b>Evelyn M. Nelson</b> , Homomorphisms of mono-unary algebras . . . . .	427
<b>Marvin E. Ortel</b> , The support of an extremal dilatation . . . . .	431
<b>R. S. Pathak and O. P. Singh</b> , Finite Hankel transforms of distributions . . . . .	439
<b>Richard Cole Penney</b> , The theory of ad-associative Lie algebras . . . . .	459
<b>Linda Ruth Sons</b> , Zero distribution of functions with slow or moderate growth in the unit disc . . . . .	473
<b>Russell Bruce Walker</b> , Transversals to laminations . . . . .	483