

Pacific Journal of Mathematics

**ON THE BEHAVIOR NEAR A TORUS OF FUNCTIONS
HOLOMORPHIC IN THE BALL**

PATRICK ROBERT AHERN

ON THE BEHAVIOR NEAR A TORUS OF FUNCTIONS HOLOMORPHIC IN THE BALL

PATRICK AHERN

If f is bounded and holomorphic in the unit ball in \mathbb{C}^n then it has radial limits at almost all points of the boundary of the ball. More is true; for example, f will have limits almost everywhere with respect to arclength on any arc that forms part of the boundary of an analytic disc. Motivated by these considerations we consider an n -dimensional torus in the boundary of the ball and ask if there are growth conditions less restrictive than boundedness that imply the existence of radial limits on this torus. We show that the answer is no for some of the standard function classes. For example, we show that there is holomorphic function of bounded mean oscillation in the ball that has a finite radial limit at no point of the torus.

Let B_n denote the unit ball in \mathbb{C}^n and let σ_n be Lebesgue measure on its boundary, ∂B_n , normalized so that $\sigma_n(\partial B_n) = 1$. If f is a holomorphic in B_n , we say that $f \in H^p(B_n)$, $0 < p < \infty$, if

$$\|f\|_p^p = \sup_{0 < r < 1} \int_{\partial B_n} |f(r\xi)|^p d\sigma_n(\xi) < \infty;$$

we say $f \in H^\infty(B_n)$ if $\|f\|_\infty = \sup_{\xi \in B_n} |f(\xi)| < \infty$. If $f \in H^2(B_n)$ we say that $f \in \text{BMO}(B_n)$ if \exists a constant C such that for all $F \in H^2(B_n)$ we have $|\int_{\partial B_n} F \bar{f} d\sigma_n| \leq C \|F\|_1$. Then $\text{BMO}(B_n)$ serves as the dual of $H^1(B_n)$ and we have $H^\infty(B_n) \subseteq \text{BMO}(B_n) \subseteq H^p(B_n)$, $0 < p < \infty$. For a more intrinsic description $\text{BMO}(B_n)$, see [1].

Next we describe some function spaces in the open unit disc U in the complex plane. If μ is a positive measure on U then $A^p(d\mu)$ will denote the space of holomorphic functions in $L^p(d\mu)$, $0 < p \leq \infty$. When $d\mu(r, \theta) = (1-r)^\alpha dr d\theta$, $\alpha > -1$, we use the notation $A^p(d\mu) = A_\alpha^p$. Finally we say that g is a Bloch function, $g \in \mathfrak{B}(U)$, if

$$\|g\|_{\mathfrak{B}} = \sup_{|z| < 1} (1 - |z|) |g'(z)| < \infty.$$

We have a mapping $\pi: \mathbb{C}^n \rightarrow \mathbb{C}$ given by $\pi(z_1, \dots, z_n) = n^{n/2} \prod_{j=1}^n z_j$. It is easily checked that $\pi(B_n) = U$, $\pi(\bar{B}_n) = \bar{U}$, and that $\pi^{-1}(\partial U) = T_n = \{(z_1, \dots, z_n): |z_j| = n^{-1/2}, j = 1, \dots, n\}$. In this paper it is shown that if $g \in A_{(n-3)/2}^p$ then $g \circ \pi \in H^p(B_n)$, and that if $g \in \mathfrak{B}(U)$ then $g \circ \pi \in \text{BMO}(B_n)$.

The following remarks are intended to motivate the results of this paper. It is known, see [8], for instance, that if $f \in H^p(B_n)$ then f has radial limits almost everywhere ($d\sigma_n$). If $f \in H^\infty(B_n)$, then more can be said; for example, f has radial limits almost everywhere with respect to arclength on the curve $\Gamma = \{(0, \dots, 0, e^{i\theta}) : 0 \leq \theta \leq 2\pi\}$. Of course $\sigma_n(\Gamma) = 0$. A. Nagel and W. Rudin [5] have generalized this to certain other curves in ∂B_n . The following question arises: are there growth conditions less restrictive than boundedness that still imply the existence of radial limits on sets of zero Lebesgue measure? It is natural to look for the existence of radial limits on submanifolds of ∂B_n that are nowhere complex tangential. One such submanifold with the largest possible dimension is the torus T_n , described above. This torus is the distinguished boundary of the poly disc $D_n = \{(z_1, \dots, z_n) : |z_j| < n^{-1/2}, j = 1, \dots, n\} \subseteq B_n$. It is known that if f is holomorphic and of bounded characteristic in D_n , then f has radial limits almost everywhere on T_n , see [7]. It has already been noted that $\pi^{-1}(\partial U) = T_n$; so that if g manifests a certain property near ∂U , $g \circ \pi$ will manifest that same property near T_n . So the results of this paper show that a function in $H^p(B_n)$ can be expected to behave no better near T_n than a function in $A^p_{(n-3)/2}$ can be expected to behave near ∂U . A similar statement can be made about the spaces $\text{BMO}(B_n)$ and $\mathfrak{B}(U)$. It is known [6] that there is a function $g \in \mathfrak{B}(U)$ that has a finite radial limit at no point of ∂U . It follows that the function $G = g \circ \pi$ is in $\text{BMO}(B_n)$ but has a finite radial limit at no point of T_n , in particular the restriction of G to D_n cannot have bounded characteristic.

If we let $D = \{(z, z, \dots, z) : |z| < n^{-1/2}\} \subseteq D_n \subseteq B_n$, then it follows from [8] and [4] that $H^p(B_n)$ and $H^p(D_n)$ have the same restriction to D . That is to say, even though the restriction of $H^p(B_n)$ to D_n contains functions of unbounded characteristic, $H^p(B_n)$ and $H^p(D_n)$ have the same restriction to the diagonal D .

1. The first result is a calculation upon which the results of this paper are based. We point out that the case $n = 2$ of Theorem 1 is computationally much simpler than the general case. Moreover, the proof shows that $w_2(r) = 2r(1 - r^2)^{-1/2}$. A formula for $w_n(r)$, $n \geq 3$, is not obvious.

THEOREM 1. For each integer $n \geq 2$, $\exists w_n : (0, 1) \rightarrow [0, \infty) \ni$

- (i) $\int_0^1 w_n(r) dr < \infty$,
- (ii) $0 < \lim_{r \rightarrow 1} w_n(r)(1 - r)^{(3-n)/2} < \infty$.

(iii) If g is a continuous complex valued function defined on \bar{U} then

$$\int_{\partial B_n} g \circ \pi \, d\sigma_n = \int_0^{2\pi} \int_0^1 g(re^{i\theta}) w_n(r) \, dr \, d\theta.$$

Proof. The proof proceeds by induction. We assume the result is true for $n - 1$, $n \geq 3$. If G is a continuous function defined on ∂B_n , then we have, [8],

$$(1.1) \quad \int_{\partial B_n} G \, d\sigma_n = \int_{B_{n-1}} \int_0^{2\pi} G\left(\xi, (1 - |\xi|^2)^{1/2} e^{i\theta}\right) \frac{d\theta}{2\pi} \, d\nu_{n-1}(\xi),$$

where ν_{n-1} denotes Lebesgue measure on \mathbf{C}^{n-1} , normalized so that $\nu_{n-1}(B_{n-1}) = 1$. Next we introduce polar coordinates, see [8], then the right hand side of (1.1) becomes,

$$(1.2) \quad 2(n - 1) \int_0^1 r^{2n-3} \int_{\partial B_{n-1}} \int_0^{2\pi} G\left((r\xi, 1 - r^2)^{1/2} e^{i\theta}\right) \frac{d\theta}{2\pi} \, d\sigma_{n-1}(\xi) \, dr.$$

Now we fix r and θ and look at the integral over ∂B_{n-1} in (1.2) in the case $G = g \circ \pi$; we obtain

$$(1.3) \quad \int_{\partial B_{n-1}} g\left(a_n a_{n-1}^{-1} (1 - r^2)^{1/2} r^{n-1} e^{i\theta} a_{n-1} \prod_{j=1}^{n-1} z_j\right) \, d\sigma_{n-1}(z),$$

where $a_n = n^{n/2}$.

By the induction hypothesis, (1.3) is equal to

$$(1.4) \quad \int_0^{2\pi} \int_0^1 g\left(a_n a_{n-1}^{-1} r^{n-1} (1 - r^2)^{1/2} e^{i\theta} \rho e^{i\psi}\right) w_{n-1}(\rho) \, d\rho \, d\psi.$$

Inserting (1.4) into (1.2) and changing the order of integration we arrive at,

$$(1.5) \quad \int_{\partial B_n} g \circ \pi \, d\sigma_n \\ = 2(n - 1) \int_0^1 \int_0^1 \int_0^{2\pi} \int_0^{2\pi} g\left(a_n a_{n-1}^{-1} r^{n-1} (1 - r^2)^{1/2} \rho e^{i(\theta+\psi)}\right) \\ \cdot \frac{d\theta}{2\pi} \, d\psi \, w_{n-1}(\rho) \, d\rho \, r^{2n-3} \, dr.$$

Next replace $\theta + \psi$ by θ , integrate out ψ , and interchange the order of integration again and we have

$$(1.6) \quad \int_{\partial B_n} g \circ \pi d\sigma_n \\ = 2(n-1) \int_0^{2\pi} \int_0^1 w_{n-1}(\rho) \int_0^1 g(a_n a_{n-1}^{-1} r^{n-1} (1-r^2)^{1/2} \rho e^{i\theta}) \\ \cdot r^{2n-3} dr d\rho d\theta.$$

Let $h(r) = a_n a_{n-1}^{-1} r^{n-1} (1-r^2)^{1/2}$; from elementary calculus it follows that h is increasing on the interval $[0, b_n]$ and decreasing on the interval $[b_n, 1]$, where $b_n = [(n-1)/n]^{1/2}$. Moreover, $h(0) = h(1) = 0$ and $h(b_n) = 1$. We also see that h' vanishes to order $n-2$ at 0, to order $-1/2$ at 1, and to order 1 at b_n . We break the innermost integral in (1.6) into two pieces, the integral from 0 to b_n , and the integral from b_n to 1. In each of these we make the substitution $t = h(r)$. If λ denotes the inverse function of h (in either case) then each of the two integrals takes the form

$$(1.7) \quad \int_0^1 g(t\rho e^{i\theta}) \lambda(t)^{-(n-1)} [h'(\lambda(t))]^{-1} dt.$$

We are interested in the behaviour of the “weight” $w(t) = \lambda(t)^{-(n-1)} [h'(\lambda(t))]^{-1}$, when t is near 1. First note that $\lambda(1) = b_n$. Next we may calculate that $h'(\lambda(t))$ vanishes like $(1-t)^{1/2}$ when t approaches 1. Now if we substitute (1.7) back in for the inner integral in (1.6) we get,

$$(1.8) \quad \int_{\partial B_n} g \circ \pi d\sigma_n = \int_0^{2\pi} \int_0^1 w_{n-1}(\rho) \int_0^1 g(t\rho e^{i\theta}) w(t) dt d\rho d\theta,$$

where $w(t) = (c_1 + o(1))(1-t)^{-1/2}$ as $t \rightarrow 1$, $c_1 > 0$. (We have absorbed the constant $2(n-1)$ into w .) Finally, in the inner integral in (1.8) we make the substitution $r = t\rho$, and then interchange the order of integration to obtain

$$(1.9) \quad \int_{\partial B_n} g \circ \pi d\sigma_n = \int_0^{2\pi} \int_0^1 g(re^{i\theta}) w_n(r) dr d\theta,$$

where

$$w_n(r) = \int_r^1 w\left(\frac{r}{\rho}\right) w_{n-1}(\rho) \frac{d\rho}{\rho}.$$

We now check that w_n has the right properties. First of all

$$\int_0^1 w_n(r) dr = \int_0^1 w(r) dr \int_0^1 w_{n-1}(r) dr < \infty,$$

so (i) is satisfied. Now we know that $w(r/\rho) = (c_1 + o(1))(1 - r/\rho)^{1/2}$, as $r/\rho \rightarrow 1$, and $w_{n-1}(\rho) = (c_2 + o(1))(1 - \rho)^{(n-4)/2}$, as $\rho \rightarrow 1$. It follows that

$$\begin{aligned} w_n(r) &= \int_r^1 w\left(\frac{r}{\rho}\right) w_{n-1}(\rho) \frac{d\rho}{\rho} \\ &= (c + o(1)) \int_r^1 (\rho - r)^{-1/2} (1 - \rho)^{(n-4)/2} d\rho \quad \text{as } r \rightarrow 0, \end{aligned}$$

for some $c > 0$. In this integral we make the substitution

$$s = (\rho - r) / (1 - \rho)$$

we arrive at

$$\begin{aligned} w_n(r) &= (c + o(1))(1 - r)^{(n-3)/2} \int_0^1 s^{-1/2} (1 - s)^{(n-4)/2} ds \\ &= (c + o(1))(1 - r)^{(n-3)/2} \beta\left(\frac{1}{2}, (n - 2)/2\right). \end{aligned}$$

To complete the proof we should check that the theorem is true for $n = 2$. This is done by the same method as the induction step given above. Indeed, it is somewhat simpler and the details will be omitted.

Now if g is continuous on \bar{U} we may apply Theorem 1 to the function $|g|^p$ and conclude that

$$\int_{\partial B_n} |Tg|^p d\sigma_n = \int_V |g|^p d\mu_n,$$

where $Tg = g \circ \pi$, and $d\mu_n(r, \theta) = w_n(r) dr d\theta$. It is now clear that T extends uniquely to be an isometry of $L^p(d\mu_n)$ into $L^p(d\sigma_n)$. If g is holomorphic, then it is obvious from Theorem 1 that $g \in L^p(d\mu_n)$ if and only if $g \in A^p_{(n-3)/2}$. Also if g is holomorphic, then so is Tg . We may conclude with

COROLLARY 1. *T is a bounded, linear, one-to-one map of $A^p_{(n-3)/2}$ into $H^p(B_n)$.*

2. In this section we show that if $g \in \mathfrak{B}(U)$ then $Tg = g \circ \pi \in \text{BMO}(B_n)$. To do this it is sufficient to show that if $g \in \mathfrak{B}(U) \exists$ constant $C \ni$ if F is a holomorphic polynomial in n variables then

$$\left| \int F \overline{g \circ \pi} d\sigma_n \right| \leq C \int |F| d\sigma_n.$$

To do this we proceed as follows. Since $F \in L^2(d\sigma_n)$ we have

$$\int F\overline{Tg} \, d\sigma_n = \int T^*F\overline{g} \, d\mu_n$$

where T^* is the adjoint of the isometry $T: L^2(d\mu_n) \rightarrow L^2(d\sigma_n)$. The proof is then accomplished in two steps. The first is to show that if F is a holomorphic polynomial in \mathbb{C}^n then T^*F is a holomorphic polynomial in \mathbb{C} and $\int |T^*F| \, d\mu_n \leq \int |F| \, d\sigma_n$. The second step is to show that if $g \in \mathfrak{B}(U)$ then \exists constant $C \ni$ for any holomorphic polynomial h of one variable we have

$$\left| \int h\overline{g} \, d\mu_n \right| \leq C \int |h| \, d\mu_n.$$

The first step is quite easy. The second is slightly trickier than it may appear.

LEMMA 2.1. *If $F(z) = \sum F_\alpha z^\alpha$ is a holomorphic polynomial then $(T^*F)(z) = \sum_{k \geq 0} F_{(k, \dots, k)} n^{-nk/2} z^k$. Moreover $\int |T^*F| \, d\mu_n \leq \int |F| \, d\sigma_n$.*

Proof. To prove the first part we show that $T^*z^\alpha = 0$ unless $\alpha_1 = \alpha_2 = \dots = \alpha_n$, and then we show $T^*z_1^k \dots z_n^k = n^{-nk/2} z^k$. Because of the rotational invariance of σ_n , the integral $I_\alpha = \int z^\alpha \overline{g(\pi(z))} \, d\sigma_n(z)$ is unchanged if z_k is replaced by $z_k e^{i\theta}$ and z_l by $z_l e^{-i\theta}$. Since $g \circ \pi$ is also unchanged by these substitutions we see that $I_\alpha = e^{i(\alpha_k - \alpha_l)\theta} I_\alpha$, for any θ . It follows that $I_\alpha = 0$ unless $\alpha_k = \alpha_l$ for all $k, l = 1, \dots, n$. If $\alpha = (k, \dots, k)$ then

$$\begin{aligned} \int z^\alpha \overline{g(\pi(z))} \, d\sigma_n(z) &= \int (z_1 \dots z_n)^k \overline{g(n^{n/2} z_1 \dots z_n)} \, d\sigma_n(z) \\ &= n^{-nk/2} \int h(\pi(z)) \, d\sigma_n(z), \end{aligned}$$

where $h(z) = z^k \overline{g(z)}$. By Theorem 1 (iii) we see that

$$\int z^\alpha \overline{g(\pi(z))} \, d\sigma_n(z) = n^{-nk/2} \int z^k \overline{g(z)} \, d\mu_n(z).$$

Since $T: L^p(d\mu_n) \rightarrow L^p(d\sigma_n)$, $1 \leq p < \infty$, is an isometry it follows that $T^*: L^q(d\sigma_n) \rightarrow L^q(d\mu_n)$, $1 < q \leq \infty$, has norm at most 1. So if F is a holomorphic polynomial in n variables

$$\int |T^*F|^q \, d\mu_n \leq \int |F|^q \, d\sigma_n$$

for each $q > 1$. Now we let $q \rightarrow 1$, by the bounded convergence theorem

$$\int |F|^q d\sigma_n \rightarrow \int |F| d\sigma_n \quad \text{and} \quad \int |T^*F|^q d\mu_n \rightarrow \int |T^*F| d\mu_n.$$

Next we want to show that if $g \in \mathfrak{B}(U) \exists$ a constant $C \ni$ for every holomorphic polynomial h in one variable we have

$$\left| \int h\bar{g} d\mu_n \right| \leq c \int |h| d\mu_n.$$

In other words, the mapping $h \mapsto \int h\bar{g} d\mu_n$ is continuous on $A^1(d\mu_n)$, the closure in $L^1(d\mu_n)$ of the holomorphic polynomials. We have already observed that since $w_n(r)$ behaves like $(1 - r)^{(n-3)/2}$ as $r \rightarrow 1$, $A^1(d\mu_n)$ and $A^1_{(n-3)/2}$ have the same elements. Now A^1_α has another name in the literature, it is called B^p where $p = 1/(2 + \alpha)$. The dual of B^p (and hence of A^1_α) is known to be a certain space of Lipschitz function depending on p , see [3]. However, the duality is effected by an integral over the “boundary” of U rather than by an integral over U itself. We seem to be saying that the dual of B^p is $\mathfrak{B}(U)$, for all p , if we use the area pairing. This can be seen as follows: by results from [3] any two B^p spaces are isomorphic by means of fractional derivatives. In particular B^p is isomorphic to $B^{1/2}$. From [3] we know that the dual of $B^{1/2}$ is the Zygmund class Λ^* . By a theorem of Zygmund, [2] Λ^* is the set of indefinite integrals of $\mathfrak{B}(U)$. However, rather than following this tortuous path we prefer to give a simple direct proof of what we need.

LEMMA 2.2. *If $\alpha > -1$ and $g \in \mathfrak{B}(U)$ then \exists a constant $C \ni$ for every holomorphic polynomial h we have*

$$\left| \int_0^{2\pi} \int_0^1 h(re^{i\theta}) \overline{g(re^{i\theta})} (1 - r)^\alpha dr d\theta \right| \leq C \int |h(re^{i\theta})| (1 - r)^\alpha dr.$$

Proof. If f is holomorphic in U , f_k will denote its k th Taylor coefficient. We then calculate that

$$\begin{aligned} \int_0^1 \int_0^{2\pi} h(re^{i\theta}) \overline{g(re^{i\theta})} d\theta (1 - r)^\alpha dr &= 2\pi \sum h_k \bar{g}_k \int_0^1 r^{2k} (1 - r)^\alpha dr \\ &= 2\pi \sum h_k \bar{g}_k \frac{\Gamma(2k + 1)\Gamma(\alpha + 1)}{\Gamma(2k + \alpha + 2)}. \end{aligned}$$

Also we see that

$$\begin{aligned}
 & \int_0^1 \int_0^{2\pi} h(re^{i\theta}) \overline{re^{i\theta} g'(re^{i\theta})} d\theta (1-r)^{\alpha+1} dr \\
 &= 2\pi \sum h_k \bar{g}_k k \int_0^1 r^{2k} (1-r)^{\alpha+1} dr \\
 &= 2\pi \sum h_k \bar{g}_k k \frac{\Gamma(2k+1)\Gamma(\alpha+2)}{\Gamma(2k+\alpha+3)} \\
 &= 2\pi \sum h_k \bar{g}_k k (\alpha+1) \frac{\Gamma(2k+1)\Gamma(\alpha+1)}{(2k+\alpha+2)\Gamma(2k+\alpha+2)} \\
 &= 2\pi(\alpha+1) \sum h_k \bar{g}_k \frac{\Gamma(2k+1)\Gamma(\alpha+1)}{\Gamma(2k+\alpha+2)} \cdot \frac{k}{2k+\alpha+2} \\
 &= 2\pi(\alpha+1) \sum h_k \bar{g}_k \frac{\Gamma(2k+1)\Gamma(\alpha+1)}{\Gamma(2k+\alpha+2)} \left\{ \frac{1}{2} - \frac{\alpha/2+1}{2k+\alpha+2} \right\} \\
 &= \frac{\alpha+1}{2} \int_0^1 \int_0^{2\pi} h(re^{i\theta}) \overline{g(re^{i\theta})} d\theta (1-r)^\alpha dr \\
 &\quad - 2\pi \frac{(\alpha+1)(\alpha+2)}{2} \sum \frac{h_k \bar{g}_k}{2k+\alpha+2} \int_0^1 r^{2k} (1-r)^\alpha dr.
 \end{aligned}$$

In other words

$$\begin{aligned}
 & \int_0^1 \int_0^{2\pi} h(re^{i\theta}) \overline{g(re^{i\theta})} d\theta (1-r)^\alpha dr \\
 &= \frac{2}{(\alpha+1)} \int_0^1 \int_0^{2\pi} h(re^{i\theta}) \overline{re^{i\theta} g'(re^{i\theta})} d\theta (1-r)^{\alpha+1} dr \\
 &\quad + 2\pi(\alpha+2) \sum \frac{h_k \bar{g}_k}{2k+\alpha+2} \int_0^1 r^{2k} (1-r)^\alpha dr.
 \end{aligned}$$

Using the fact that $g \in \mathfrak{B}(\mathcal{U})$, the first term above has modulus at most

$$\frac{2}{\alpha+1} \|g\|_{\mathfrak{B}} \int |h|(1-r)^\alpha dr d\theta.$$

Using the fact that $|g_k| \leq C \|g\|_{\mathfrak{B}}$ for some constant C , the modulus of the second term is at most

$$\sum 2\pi(\alpha+2) C \|g\|_{\mathfrak{B}} \frac{|h_k|}{k+1} \int_0^1 r^{2k} (1-r)^\alpha dr.$$

The proof will be finished if we show that there is a constant C such that

$$\sum \frac{|h_k|}{k+1} \int_0^1 r^{2k} (1-r)^\alpha dr \leq C \int_0^1 \int_0^{2\pi} |h(re^{i\theta})| (1-r)^\alpha d\theta dr,$$

for any holomorphic polynomial h . To see this we apply the well known inequality of Hardy and Littlewood [2] on a circle of radius $r < 1$ to obtain

$$\sum \frac{|h_k| r^k}{k + 1} \leq C \int_0^{2\pi} |h(re^{i\theta})| d\theta,$$

and hence

$$\sum \frac{|h_k| r^{2k}}{k + 1} (1 - r)^\alpha \leq C \int_0^{2\pi} |h(re^{i\theta})| d\theta (1 - r)^\alpha.$$

Now we just integrate on r from 0 to 1.

Next we want to show that Lemma 2.2 remains true if the measure $(1 - r)^{(n-3)/2} dr d\theta$ is replaced by the measure $d\mu_n$. Since $A^1_{(n-3)/2}$ and $A^1(d\mu_n)$ are the “same” it may seem obvious that they have the “same” dual. A little reflection shows that we need more information to draw this conclusion. It turns out that we do know enough about μ_n to get the result we want.

LEMMA 2.3. *If $n \geq 2$ and $g \in \mathfrak{B}(U)$ are given, \exists a constant $C \ni$ for every holomorphic polynomial h we have*

$$\left| \int h \bar{g} d\mu_n \right| \leq C \int |h| d\mu_n.$$

Proof. The idea is to show that $\int h \bar{g} d\mu_n$ and $\int h \bar{g} (1 - r)^{(n-3)/2} dr d\theta$ differ by a manageable amount and then to apply Lemma 2.2. As before we calculate that

$$\begin{aligned} & \int_0^1 \int_0^{2\pi} h(re^{i\theta}) \overline{g(re^{i\theta})} (1 - r)^{(n-3)/2} d\theta dr \\ &= 2\pi \sum h_k \bar{g}_k \int_0^1 r^{2k} (1 - r)^{(n-3)/2} dr \\ &= 2\pi \sum h_k \bar{g}_k \frac{\Gamma(2k + 1) \Gamma((n - 1)/2)}{\Gamma(2k + 1 + (n - 1)/2)}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \int h \bar{g} d\mu_n &= \int_0^1 \int_0^{2\pi} h(re^{i\theta}) \overline{g(re^{i\theta})} d\theta w_n(r) dr \\ &= 2\pi \sum h_k \bar{g}_k \int_0^1 r^{2k} w_n(r) dr. \end{aligned}$$

To calculate $\int_0^1 r^{2k} w_n(r) dr$ we use Theorem 1, (iii);

$$\begin{aligned} \int_0^1 r^{2k} w_n(r) dr &= \frac{1}{2} \int_0^{2\pi} \int_0^1 |re^{i\theta}|^{2k} w_n(r) dr d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} \int_0^1 g(re^{i\theta}) w_n(r) dr d\theta, \quad \text{where } g(z) = |z|^{2k}. \end{aligned}$$

So, by Theorem 1 (iii) we see that

$$\begin{aligned} \int_0^1 r^{2k} w_n(r) dr &= \frac{1}{2\pi} \int g \circ \pi d\sigma_n = \frac{1}{2\pi} \int |n^{n/2} z_1 \cdots z_n|^{2k} d\sigma_n(z) \\ &= \frac{n^{nk}}{2\pi} \int |z_1 \cdots z_n|^{2k} d\sigma_n(z) = \frac{n^{nk}}{2\pi} (n-1)! \frac{(k!)^n}{(n-1+(nk))!} \\ &= \frac{(n-1)!}{2\pi} n^{nk} \frac{\Gamma(k+1)^n}{\Gamma(nk+n)}; \end{aligned}$$

here we have used the formula found in [8]. So we conclude that

$$\int h\bar{g} d\mu_n = \sum h_k \bar{g}_k (n-1)! n^{nk} \frac{\Gamma(k+1)^n}{\Gamma(nk+n)}.$$

We will use Stirling’s formula:

$$\Gamma(x+1) = (2\pi e)^{1/2} \left(\frac{x}{e}\right)^{x+1/2} \left[1 + O\left(\frac{1}{x}\right)\right], \quad \text{as } x \rightarrow \infty.$$

We have

$$\begin{aligned} n^{nk} \frac{\Gamma(k+1)^n}{\Gamma(nk+n)} &= n^{nk} \frac{\left[(2\pi e)^{1/2} (ke^{-1})^{k+1/2}\right]^n}{(2\pi e)^{1/2} ((nk+n+1)e^{-1})^{nk+n-1/2}} \left[1 + O\left(\frac{1}{k}\right)\right] \\ &= C_n n^{nk} \frac{k^{nk+n/2}}{(nk)^{nk+n-1/2} (1+(n-1)/nk)^{nk+n-1/2}} \left[1 + O\left(\frac{1}{k}\right)\right] \\ &= C'_n \frac{1}{k^{(n-1)/2}} \frac{1}{(1+(n-1)/nk)^{nk+n-1/2}} \cdot \left[1 + O\left(\frac{1}{k}\right)\right], \end{aligned}$$

where C_n, C'_n depend only on n , not on k . Next note that

$$(1+(n-1)/nk)^{nk} = e^{n-1} [1 + O(1/k)].$$

We may conclude that

$$n^{nk} \frac{\Gamma(k + 1)^n}{\Gamma(nk + n)} = C_0 \frac{1}{k^{(n-1)/2}} \left[1 + O\left(\frac{1}{k}\right) \right],$$

where C_0 depends only on n , not on k . A similar calculation shows us that

$$\frac{\Gamma(2k + 1)}{\Gamma(2k + 1 + (n - 1)/2)} = C_1 \frac{1}{k^{(n-1)/2}} \left[1 + O\left(\frac{1}{k}\right) \right]$$

for some constant C_1 . We may conclude that

$$\begin{aligned} \int h\bar{g} \, d\mu_n &= C \int h\bar{g}(1 - r)^{(n-3)/2} \, dr \, d\theta \\ &+ O\left[\sum \frac{h_k \bar{g}_k}{k^{(n+1)/2}} \right] \end{aligned}$$

for some constant C . By Lemma 2.2 the first term above has its modulus at most a constant times

$$\int |h|(1 - r)^{(n-3)/2} \, dr \, d\theta$$

which is in turn bounded by a constant times $\int |h| \, d\mu_n$. The second term is bounded by a constant times

$$\sum \frac{|h_k|}{k} \frac{1}{k^{(n-1)/2}}$$

which is again bounded by a constant times

$$\sum \frac{|h_k|}{k} \int_0^1 r^{2k}(1 - r)^{(n-3)/2} \, dr,$$

by Stirling's formula. This last expression, as was seen at the end of the proof of Lemma 2.2 is at most a constant times $\int |h|(1 - r)^{(n-3)/2} \, dr \, d\theta$, which is, finally, majorized by $\int |h| \, d\mu_n$. This completes the proof.

REFERENCES

[1] R. Coifman, R. Rochberg and G. Weiss, *Factorization theorems for Hardy spaces in several variables*, Ann. Math., **103** (1976), 611–635.
 [2] P. Duren, *Theory of H^p Spaces*, Academic Press, New York and London, 1970.
 [3] P. Duren, B. Romberg and A. Shields, *Linear functionals on H^p with $0 < p < 1$* , J. Reine Angew. Math., **238** (1969), 32–60.
 [4] C. Horowitz and D. Oberlin, *Restrictions of H^p functions to the diagonal of U^n* , Indiana Math. J., **24**, (1975), 767–772.

- [5] A. Nagel and W. Rudin, *Local boundary behavior of bounded holomorphic functions*, *Canad. J. Math.*, **30** (1978), 583–592.
- [6] C. Pommerenke, *On Bloch functions*, *J. London Math. Soc.*, **2**(2), (1970), 689–695.
- [7] W. Rudin, *Function Theory in Polydiscs*, Benjamin, New York and Amsterdam, 1969.
- [8] _____, *Function Theory in The Unit Ball in \mathbf{C}^n* , Springer, New York, Heidelberg and Berlin, 1980.

Received September 22, 1981 and in revised form May 5, 1982.

UNIVERSITY OF WISCONSIN
MADISON, WI 53706

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR OGUS

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90089-1113

R. FINN and H. SAMELSON

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

(1906–1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$132.00 a year (6 Vol., 12 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics ISSN 0030-8730 is published monthly by the Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1983 by Pacific Journal of Mathematics

Driss Abouabdillah , <i>Topologies de corps A linéaires</i>	257
Patrick Robert Ahern , <i>On the behavior near a torus of functions holomorphic in the ball</i>	267
Donald Werner Anderson , <i>There are no phantom cohomology operations in K-theory</i>	279
Peter Bloomfield , Nicolas P. Jewell and Eric Hayashi , <i>Characterizations of completely nondeterministic stochastic processes</i>	307
Sydney Dennis Bulman-Fleming and K. McDowell , <i>Absolutely flat semigroups</i>	319
C. Debiève , <i>On a Radon-Nikodým problem for vector-valued measures</i>	335
Dragomir Z. Djokovic , <i>Products of positive reflections in real orthogonal groups</i>	341
Thomas Farmer , <i>The dual of the nilradical of the parabolic subgroups of symplectic groups</i>	349
Gary R. Greenfield , <i>Uniform distribution in subgroups of the Brauer group of an algebraic number field</i>	369
Paul Daniel Hill , <i>When $\text{Tor}(A, B)$ is a direct sum of cyclic groups</i>	383
Hiroshi Maehara , <i>Regular embeddings of a graph</i>	393
Nikolaos S. Papageorgiou , <i>Nonsmooth analysis on partially ordered vector spaces. I. Convex case</i>	403
Louis Jackson Ratliff, Jr. , <i>Powers of ideals in locally unmixed Noetherian rings</i>	459
F. Dennis Sentiilles and Robert Francis Wheeler , <i>Pettis integration via the Stonian transform</i>	473