

# Pacific Journal of Mathematics

**A PROOF OF THE BENDER-KNUTH CONJECTURE**

BASIL GORDON

# A PROOF OF THE BENDER-KNUTH CONJECTURE

BASIL GORDON

Let  $b_r(n | I_m)$  denote the number of  $r$ -rowed partitions of  $n$  whose parts lie in the set  $I_m = \{1, 2, \dots, m\}$  and decrease strictly along each row. It is shown that

$$\sum_{n=0}^{\infty} b_r(n | I_m) x^n = \prod_{i=1}^m \prod_{j=1}^i (1 - x^{r+i+j-1}) / (1 - x^{i+j-1}).$$

**1. Introduction.** For any given set  $S$  of positive integers, let  $b_r(n | S)$  denote the number of  $r$ -rowed partitions of  $n$  whose parts lie in  $S$  and decrease strictly along each row. Put  $b(n | S) = \lim_{r \rightarrow \infty} b_r(n | S)$ . Bender and Knuth [3] have proved the remarkable formula

$$\begin{aligned} B(x | S) &= \sum_{n=0}^{\infty} b(n | S) x^n \\ &= \prod_{i \in S} (1 - x^i)^{-1} \prod_{\substack{j, k \in S \\ j < k}} (1 - x^{j+k})^{-1}, \end{aligned}$$

valid for  $|x| < 1$ . As yet no such simple expression has been found for  $B_r(x | S) = \sum_{n=0}^{\infty} b_r(n | S) x^n$ . However there are two situations where a "product" formula for  $B_r(x | S)$  can indeed be given, namely when  $S = I_m = \{1, 2, \dots, m\}$  or  $S = J_m = \{1, 3, 5, \dots, 2m - 1\}$ . The formulas are:

$$(1) \quad B_r(x | I_m) = \prod_{i=1}^m \prod_{j=1}^i \frac{1 - x^{r+i+j-1}}{1 - x^{i+j-1}},$$

$$(2) \quad B_r(x | J_m) = \prod_{i=1}^m \frac{1 - x^{r+2i-1}}{1 - x^{2i-1}} \prod_{j=i+1}^m \frac{1 - x^{2(r+i+j-1)}}{1 - x^{2(i+j-1)}}.$$

Equation (2) was conjectured by MacMahon [5], while (1) was conjectured by Bender and Knuth [2]. Some years earlier the author had already found a proof of (1), but published only the limiting case  $m \rightarrow \infty$ . Then Andrews [1] proved (2), and also showed in [3] that (1) and (2) are equivalent. This of course gave another proof of (1). Over the years, a number of people have expressed a desire to see the original direct proof of (1) in print. It will therefore be belatedly presented here.

**2. Notation.** If  $n$  is a positive integer, while  $a$  and  $y$  are indeterminates, we write

$$(a; y)_n = (1 - a)(1 - ay)(1 - ay^2) \cdots (1 - ay^{n-1}).$$

By convention,  $(a; y)_0 = 1$ . If  $0 \leq k \leq n$ , put

$$\begin{bmatrix} n \\ k \end{bmatrix} = \frac{(x; x)_n}{(x; x)_k (x; x)_{n-k}}.$$

If  $k < 0$  or  $k > n$ , define  $\begin{bmatrix} n \\ k \end{bmatrix} = 0$ .

**3. Proof of (1).** Let  $\lambda_1, \lambda_2, \dots, \lambda_r$  be a sequence of integers satisfying  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r \geq 0$ . We consider  $r$ -rowed partitions of the type enumerated by  $b_r(n | S)$ , but where there are exactly  $\lambda_i$  non-zero parts in the  $i$ th row ( $i = 1, \dots, r$ ). Let  $b_r(n; \lambda_1, \dots, \lambda_r | S)$  be the number of such partitions of  $n$ , and put

$$B_r(x; \lambda_1, \dots, \lambda_r | S) = \sum_{n=0}^{\infty} b_r(n; \lambda_1, \dots, \lambda_r | S) x^n.$$

Clearly  $B_r(x | S) = \sum_{(\lambda_i)} B_r(x; \lambda_1, \dots, \lambda_r | S)$ , where the sum is extended over all sequences  $(\lambda_i)$  with  $\lambda_1 \geq \dots \geq \lambda_r \geq 0$ . We now obtain an expression for  $B_r(x; \lambda_1, \dots, \lambda_r | I_m)$  as a determinant.

**THEOREM 1.**

$$B_r(x; \lambda_1, \dots, \lambda_r | I_m) = \det_{1 \leq i, j \leq r} \left( x^{\binom{m}{2} + \lambda_j + 1} \begin{bmatrix} m \\ i - j + \lambda_j \end{bmatrix} \right).$$

*Proof.* Given a partition  $\pi$  of  $n$  of the type enumerated by  $b_r(n; \lambda_1, \dots, \lambda_r | I_m)$ , we subtract 1 from each of its parts. If  $\lambda = \lambda_1 + \lambda_2 + \dots + \lambda_r$ , this gives a partition of  $n - \lambda$  of the type enumerated by  $b_r(n - \lambda; \lambda_1 - \varepsilon_1, \dots, \lambda_r - \varepsilon_r | I_{m-1})$ , where  $\varepsilon_j = 0$  or 1 according as the last part of the  $j$ th row of  $\pi$  is greater than 1 or equal to 1. Moreover, every such partition of  $n - \lambda$  is the image of exactly one  $\pi$  under this map. It follows readily that

$$B_r(x; \lambda_1, \dots, \lambda_r | I_m) = x^\lambda \sum_{\varepsilon_j=0}^1 B_r(x; \lambda_1 - \varepsilon_1, \dots, \lambda_r - \varepsilon_r | I_{m-1}),$$

where we make the convention that  $B_r(x; \lambda_1 - \varepsilon_1, \dots, \lambda_r - \varepsilon_r | I_{m-1}) = 0$  if the inequalities  $\lambda_1 - \varepsilon_1 \geq \lambda_2 - \varepsilon_2 \geq \dots \geq \lambda_r - \varepsilon_r \geq 0$  are not satisfied. If  $\lambda_1 = \dots = \lambda_r = 0$ , we have  $B_r(x; 0, \dots, 0 | I_m) = 1$ .

Now let  $C_r(x; \lambda_1, \dots, \lambda_r | I_m)$  be the determinant in the statement of the theorem. Then

$$C_r(x; 0, \dots, 0 | I_m) = \det_{1 \leq i, j \leq r} \left( x^{\binom{i-j+1}{2}} \begin{bmatrix} m \\ i-j \end{bmatrix} \right) = 1,$$

since the diagonal entries are all equal to 1, while those above the diagonal vanish. Thus  $B_r(x; 0, \dots, 0 | I_m) = C_r(x; 0, \dots, 0 | I_m)$ . The proof can now be completed by induction by showing that

$$C_r(x; \lambda_1, \dots, \lambda_r | I_m) = x^\lambda \sum_{\epsilon_j=0}^1 C_r(x; \lambda_1 - \epsilon_1, \dots, \lambda_r - \epsilon_r | I_{m-1}).$$

To do this, we recall the well-known identity

$$\begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} a-1 \\ b-1 \end{bmatrix} + x^b \begin{bmatrix} a-1 \\ b \end{bmatrix}.$$

Using this identity, we can write the general term of the determinant  $C_r(x; \lambda_1, \dots, \lambda_r | I_m)$  in the form

$$x^{\binom{i-j+\lambda_j+1}{2}} \begin{bmatrix} m-1 \\ i-j+\lambda_j-1 \end{bmatrix} + x^{\binom{i-j+\lambda_j+1}{2}+i-j+\lambda_j} \begin{bmatrix} m-1 \\ i-j+\lambda_j \end{bmatrix}.$$

Now

$$\binom{i-j+\lambda_j+1}{2} = \binom{i-j+\lambda_j}{2} + i-j+\lambda_j,$$

so the above term can be written as

$$x^{i-j+\lambda_j} \left( x^{\binom{i-j+\lambda_j}{2}} \begin{bmatrix} m-1 \\ i-j+\lambda_j-1 \end{bmatrix} + x^{\binom{i-j+\lambda_j+1}{2}} \begin{bmatrix} m-1 \\ i-j+\lambda_j \end{bmatrix} \right).$$

We can now remove a common factor of  $x^i$  from the  $i$ th row, and a common factor of  $x^{-j+\lambda_j}$  from the  $j$ th column. The product of these factors is  $x^{\lambda_1+\dots+\lambda_r} = x^\lambda$ . Hence

$$\begin{aligned} (3) \quad C_r(x; \lambda_1, \dots, \lambda_r | I_m) &= x^\lambda \det \left( x^{\binom{i-j+\lambda_j}{2}} \begin{bmatrix} m-1 \\ i-j+\lambda_j-1 \end{bmatrix} + x^{\binom{i-j+\lambda_j+1}{2}} \begin{bmatrix} m-1 \\ i-j+\lambda_j \end{bmatrix} \right). \end{aligned}$$

The general term of the determinant on the right side of (3) can be written in the form

$$\sum_{\epsilon_j=0}^1 x^{\binom{i-j+\lambda_j+1-\epsilon_j}{2}} \begin{bmatrix} m-1 \\ i-j+\lambda_j-\epsilon_j \end{bmatrix}.$$

Hence, since the determinant is a linear function of each column vector, we have

$$\begin{aligned} C_r(x; \lambda_1, \dots, \lambda_r | I_m) &= x^\lambda \sum_{\epsilon_j=0}^1 \det \left( x^{(\binom{r-j+\lambda_j}{2} + 1 - \epsilon_j)} \begin{bmatrix} m-1 \\ i-j+\lambda_j-\epsilon_j \end{bmatrix} \right) \\ &= x^\lambda \sum_{\epsilon_j=0}^1 C_r(x; \lambda_1 - \epsilon_1, \dots, \lambda_r - \epsilon_r | I_m). \end{aligned}$$

This completes the proof of Theorem 1.

Now let

$$a_\nu = x^{\binom{r+1}{2}} \begin{bmatrix} m \\ \nu \end{bmatrix}, \quad g_i = i, \quad h_j = \lambda_j - j.$$

Then the result of Theorem 1 can be written in the form

$$B_r(x; \lambda_1, \dots, \lambda_r | I_m) = \det(a_{g_i+h_j}).$$

The requirement  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r$  is equivalent to  $h_1 > h_2 > \dots > h_r$ , and therefore

$$B_r(x | I_m) = \sum_{h_1 > \dots > h_r} \det(a_{g_i+h_j}).$$

Set  $s = \sum_{n=0}^\infty a_n$ ,  $c_\nu = \sum_{n=0}^\infty a_n a_{n+\nu}$ ,  $d_\nu = c_0 + 2(c_1 + \dots + c_{\nu-1}) + c_\nu$  ( $\nu > 0$ ),  $d_0 = 0$ , and  $d_{-\nu} = -d_\nu$ . Then by Lemma 1 of [4], we see that for even  $r$ ,  $B_r(x | I_m)$  is the Pfaffian of the skew-symmetric  $r \times r$  matrix  $D_r = (d_{j-i})$ , while for odd  $r$ ,  $B_r(x | I_m)$  is the Pfaffian of the  $(r+1) \times (r+1)$  matrix

$$D'_r = \begin{bmatrix} D_r & -s \\ s & 0 \end{bmatrix}$$

obtained by bordering  $D_r$  with a row of  $s$ 's, a column of  $-s$ 's, and a zero.

We next proceed to evaluate the quantities  $s$  and  $c_{\nu-1} + c_\nu$ .

**THEOREM 2.**  $s = \prod_{i=1}^m (1 + x^i)$ , and

$$c_{\nu-1} + c_\nu = x^{\binom{\nu}{2}} \begin{bmatrix} 2m+1 \\ m+\nu \end{bmatrix}.$$

*Proof.* The quantity  $a_\nu = x^{\binom{r+1}{2}} \begin{bmatrix} m \\ \nu \end{bmatrix}$  can be interpreted combinatorially as the generating function of ordinary (i.e. one-rowed) partitions into exactly  $\nu$  distinct parts, all  $\leq m$ . This observation follows from Theorem 1 with  $r = 1$ , but is also easy to see directly. On the other hand, if the

product  $\prod_{i=1}^m(1 + x^i y)$  is expanded as a power series in  $x$  and  $y$ , the coefficient of  $y^\nu$  is also the generating function of such partitions. Therefore

$$(4) \quad \prod_{i=1}^m (1 + x^i y) = \sum_{\nu=0}^{\infty} a_\nu y^\nu.$$

Putting  $y = 1$  in (4), we get  $\prod_{i=1}^m(1 + x^i) = \sum_{\nu=0}^{\infty} a_\nu = s$ . Now let  $f_m(x, y) = \prod_{i=1}^m(1 + x^i y)(1 + x^i y^{-1})$ . Then from (4) it follows that

$$f_m(x, y) = \sum_{\lambda=0}^m \sum_{\mu=0}^m a_\lambda a_\mu y^{\lambda-\mu} = \sum_{\nu=-m}^m \left( \sum_{\lambda-\mu=\nu} a_\lambda a_\mu \right) y^\nu.$$

For  $\nu \geq 0$ , we have  $\sum_{\lambda-\mu=\nu} a_\lambda a_\mu = \sum_{n=0}^m a_n a_{n+\nu} = c_\nu$ . Hence if we put  $c_{-\nu} = c_\nu$ , we have  $f_m(x, y) = \sum_{\nu=-\infty}^{\infty} c_\nu(m) y^\nu$ , where we have written  $c_\nu(m)$  instead of  $c_\nu$  to emphasize the dependence on  $m$ . Now

$$f(x, y) = f_{m-1}(x, y)(1 + x^m y)(1 + x^m y^{-1}),$$

and hence

$$\sum_{-\infty}^{\infty} c_\nu(m) y^\nu = (1 + x^m y)(1 + x^m y^{-1}) \sum_{-\infty}^{\infty} c_\nu(m-1) y^\nu.$$

Equating powers of  $y^\nu$  in these two Laurent series, we find that

$$(5) \quad c_\nu(m) = (1 + x^{2m})c_\nu(m-1) + x^m c_{\nu-1}(m-1) + x^m c_{\nu+1}(m-1).$$

For convenience of notation, put  $\gamma_\nu(m) = c_{\nu-1}(m) + c_\nu(m)$ . If in equation (5) we replace  $\nu$  by  $\nu - 1$ , and then add the result to (5), we obtain

$$(6) \quad \gamma_\nu(m) = (1 + x^{2m})\gamma_\nu(m-1) + x^m \gamma_{\nu-1}(m-1) + x^m \gamma_{\nu+1}(m-1).$$

When  $m = 0$ , we have  $a_0 = 1$  and  $a_\nu = 0$  for all  $\nu > 0$ . Hence  $c_0(0) = 1$ , while  $c_\nu(0) = 0$  for all  $\nu \neq 1$ . This gives  $\gamma_0(0) = \gamma_1(0) = 1$ , and  $\gamma_\nu(0) = 0$  for all  $\nu \neq 0, 1$ . On the other hand, when  $m = 0$  we have

$$x^{\binom{2}{2}} \left[ \begin{matrix} 2m + 1 \\ m + 1 - \nu \end{matrix} \right] = x^{\binom{2}{2}} \left[ \begin{matrix} 1 \\ 1 - \nu \end{matrix} \right],$$

which is also equal to 1 when  $\nu = 0$  or 1, and is equal to 0 otherwise. This proves the theorem in the case  $m = 0$ .

Now suppose that  $m > 0$ , and that the theorem has already been proved for  $m - 1$ . Then from equation (6) we have

$$(7) \quad \gamma_\nu(m) = (1 + x^{2m})x^{\binom{2}{2}} \left[ \begin{matrix} 2m - 1 \\ m - 1 + \nu \end{matrix} \right] + x^m x^{\binom{r-1}{2}} \left[ \begin{matrix} 2m - 1 \\ m - 2 + \nu \end{matrix} \right] \\ + x^m x^{\binom{r+1}{2}} \left[ \begin{matrix} 2m - 1 \\ m + \nu \end{matrix} \right].$$

It is a straightforward matter to check that the expression on the right side of (7) is equal to

$$x^{\binom{2}{2}} \left[ \begin{matrix} 2m + 1 \\ m + \nu \end{matrix} \right].$$

This proves Theorem 2 by induction on  $m$ .

If  $\nu > 0$ , we have  $d_\nu = \sum_{\mu=1}^{\nu} (c_{\mu-1} + c_\mu)$ , and therefore by Theorem 2,

$$d_\nu = \sum_{\mu=1}^{\nu} x^{\binom{2}{\mu}} \left[ \begin{matrix} 2m + 1 \\ m + \mu \end{matrix} \right].$$

Consider now the determinant  $\det D_r$ , where  $r$  is even. Subtracting each row from the previous row, we obtain

$$\det D_r = \begin{vmatrix} \gamma_1 & \gamma_1 & \gamma_2 & \gamma_3 & \cdots & \gamma_{r-1} \\ \gamma_2 & \gamma_1 & \gamma_1 & \gamma_2 & \cdots & \gamma_{r-2} \\ \vdots & & & & & \\ \gamma_{r-1} & \gamma_{r-2} & \gamma_{r-3} & \gamma_{r-4} & \cdots & \gamma_1 \\ -d_{r-1} & -d_{r-2} & -d_{r-3} & -d_{r-4} & \cdots & 0 \end{vmatrix}.$$

Adding all the rows to the last row, we get

$$\det D_r = \begin{vmatrix} \gamma_1 & \gamma_1 & \gamma_2 & \gamma_3 & \cdots & \gamma_{r-1} \\ \gamma_2 & \gamma_1 & \gamma_1 & \gamma_2 & \cdots & \gamma_{r-2} \\ \vdots & & & & & \\ \gamma_{r-1} & \gamma_{r-2} & \gamma_{r-3} & \gamma_{r-4} & \cdots & \gamma_1 \\ 0 & d_1 & d_2 & d_3 & \cdots & d_{r-1} \end{vmatrix}.$$

It is now convenient to extend the definition of  $\gamma_\nu$  to  $\nu \leq 0$ , by putting  $\gamma_\nu = \gamma_{1-\nu}$  in that case. Thus for  $\nu \leq 0$ , we have

$$\gamma_\nu = x^{\binom{1-\nu}{2}} \left[ \begin{matrix} 2m + 1 \\ m + 1 - \nu \end{matrix} \right] = x^{\frac{\nu(\nu-1)}{2}} \left[ \begin{matrix} 2m + 1 \\ m + \nu \end{matrix} \right].$$

If we make the convention that  $\binom{\nu}{2} = \nu(\nu - 1)/2$  for all  $\nu \in \mathbf{Z}$ , then we have

$$\gamma_\nu = x^{\binom{\nu}{2}} \left[ \begin{matrix} 2m + 1 \\ m + \nu \end{matrix} \right]$$

for all  $\nu$ . The above expression for  $\det D_r$  can be written as

$$(8) \quad \det D_r = \begin{vmatrix} \gamma_0 & \gamma_1 & \gamma_2 & \cdots & \gamma_{r-1} \\ \gamma_{-1} & \gamma_0 & \gamma_1 & \cdots & \gamma_{r-2} \\ \vdots & & & & \\ \gamma_{2-r} & \gamma_{3-r} & \gamma_{4-r} & \cdots & \gamma_1 \\ 0 & d_1 & d_2 & \cdots & d_{r-1} \end{vmatrix}.$$

In exactly the same way we find that for  $r$  odd,

$$\det D'_r = \begin{vmatrix} \gamma_0 & \gamma_1 & \gamma_2 & \cdots & \gamma_{r-1} & 0 \\ \gamma_{-1} & \gamma_0 & \gamma_1 & \cdots & \gamma_{r-2} & 0 \\ \vdots & & & & & \\ \gamma_{2-r} & \gamma_{3-r} & \gamma_{4-r} & \cdots & \gamma_1 & 0 \\ 0 & d_1 & d_2 & \cdots & d_{r-1} & -s \\ s & s & s & \cdots & s & 0 \end{vmatrix},$$

and hence

$$(9) \quad \det D'_r = s^2 \begin{vmatrix} \gamma_0 & \gamma_1 & \cdots & \gamma_{r-1} \\ \gamma_{-1} & \gamma_0 & \cdots & \gamma_{r-2} \\ \vdots & & & \\ \gamma_{2-r} & \gamma_{3-r} & \cdots & \gamma_1 \\ 1 & 1 & \cdots & 1 \end{vmatrix}.$$

The idea is now to put the determinants (8) and (9) into superdiagonal form by elementary row operations. All but the last of these operations are the same for both determinants; we describe these in terms of an arbitrary matrix  $(a_{ij})$ ,  $1 \leq i, j \leq r$ . For the moment we leave the bottom row unchanged. For each  $i$  in the range  $2 \leq i \leq r - 1$ , we multiply the  $(i - 1)$ th row by  $a_{i,1}/a_{i-1,1}$  and subtract the result from the  $i$ th row. This gives a matrix  $(b_{ij})$  with  $b_{i1} = 0$  for  $2 \leq i \leq r - 1$ . Next, for each  $i$  in the range  $3 \leq i \leq r - 1$ , we multiply the  $(i - 1)$ th row by  $b_{i,2}/b_{i-1,2}$  and subtract the result from the  $i$ th row. This gives a matrix  $(c_{ij})$  with  $c_{i1} = 0$  for  $2 \leq i \leq r - 1$  and  $c_{i2} = 0$  for  $3 \leq i \leq r - 1$ . We proceed in this manner until we obtain a matrix which, except for its bottom row, is in supertriangular form.



In the present case, if we temporarily ignore the factor  $s^2$  in (9) we have, for both (8) and (9).

$$a_{i,j} = \gamma_{j-i} = \left[ \begin{array}{c} 2m+1 \\ m+j-i \end{array} \right] x^{\binom{j-i}{2}} \quad \text{for } 1 \leq i \leq r-1.$$

When the above procedure is applied, the first step yields

$$b_{ij} = \frac{1-x^{j-1}}{1-x^{m+i}} \left[ \begin{array}{c} 2m+2 \\ m+j-i+1 \end{array} \right] x^{\binom{j-i}{2}} \quad \text{for } 2 \leq i \leq r-1,$$

the second step yields

$$c_{ij} = \frac{(1-x^{j-1})(1-x^{j-2})}{(1-x^{m+i})(1-x^{m+i+1})} \left[ \begin{array}{c} 2m+3 \\ m+j-i+2 \end{array} \right] x^{\binom{j-i}{2}} \quad \text{for } 3 \leq i \leq r-1,$$

etc. In general if  $a_{ij}^{(p)}$  is the matrix obtained after  $p$  steps of the procedure, we have

$$(10) \quad a_{ij}^{(p)} = \frac{\left[ \begin{array}{c} j-1 \\ p \end{array} \right]}{\left[ \begin{array}{c} m+i \\ p \end{array} \right]} \left[ \begin{array}{c} 2m+p+1 \\ m+j-i+p \end{array} \right] x^{\binom{j-i}{2}} \quad \text{for } p+1 \leq i \leq r-1.$$

The proof is by a straightforward induction on  $p$ , which we omit here. Since the  $i$ th row ( $1 \leq i \leq r-1$ ) remains constant after  $i-1$  steps, the final determinant obtained from (8) is

$$(11) \quad \begin{vmatrix} A_{11} & A_{12} & A_{13} & \cdots & A_{1,r-1} & A_{1,r} \\ 0 & A_{22} & A_{23} & \cdots & A_{2,r-1} & A_{2,r} \\ 0 & 0 & A_{33} & \cdots & A_{3,r-1} & A_{3,r} \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & A_{r-1,r-1} & A_{r-1,r} \\ 0 & d_1 & d_2 & \cdots & d_{r-2} & d_{r-1} \end{vmatrix},$$

where

$$(12) \quad A_{ij} = \frac{\left[ \begin{array}{c} j-1 \\ i-1 \end{array} \right]}{\left[ \begin{array}{c} m+i \\ i-1 \end{array} \right]} \left[ \begin{array}{c} 2m+i \\ m+j-1 \end{array} \right] x^{\binom{j-i}{2}}.$$

(This is obtained by putting  $p = i - 1$  in (10).) Similarly, the result of performing these operations on (9) is the determinant

$$(13) \quad s^2 \begin{vmatrix} A_{11} & A_{12} & A_{13} & \cdots & A_{1,r-1} & A_{1,r} \\ 0 & A_{22} & A_{23} & \cdots & A_{2,r-1} & A_{2,r} \\ 0 & 0 & A_{33} & \cdots & A_{3,r-1} & A_{3,r} \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & A_{r-1,r-1} & A_{r-1,r} \\ 1 & 1 & 1 & \cdots & 1 & 1 \end{vmatrix}.$$

The next task is to clear out the bottom rows of (11) and (13), except for their rightmost entries. To do this for (13), we multiply the  $(2k + 1)$ th row ( $0 \leq k \leq (r - 4)/2$ ) by

$$(14) \quad B_k = \frac{1}{\begin{bmatrix} 2m+1 \\ m \end{bmatrix}} \frac{(x; x^2)_k}{(x^{2m+3}; x^2)_k},$$

and subtract the result from the bottom row. To clear out (11), we multiply its  $(2k + 2)$ th row ( $0 \leq k \leq (r - 3)/2$ ) by

$$(15) \quad C_k = \frac{(x^2; x^2)_k}{(x^{2m+4}; x^2)_k (1 + x^{m+1})},$$

and subtract the result from the bottom row. To show that these operations do indeed clear out the bottom rows of (13) and (11), we must evidently prove that

$$\sum_{k=0}^{(r-4)/2} B_k A_{2k,j} = 0 \quad (1 \leq j \leq r - 1)$$

and

$$\sum_{k=0}^{(r-3)/2} C_k A_{2k+1,j} = 0 \quad (1 \leq j \leq r - 1).$$

In view of (12), (14) and (15), this is tantamount to showing that

$$(16) \quad \sum_{k=0}^{(r-4)/2} \frac{\begin{bmatrix} j-1 \\ 2k \end{bmatrix} \begin{bmatrix} 2m+2k+1 \\ m+j-1 \end{bmatrix}}{\begin{bmatrix} 2m+1 \\ m \end{bmatrix}} \frac{(x^2; x^2)_k}{(x^{2m+3}; x^2)_k} x^{\binom{j-1}{2}-2k} = 1$$

( $1 \leq j \leq r - 1$ ),

and

$$(17) \quad \sum_{k=0}^{(r-3)/2} \begin{bmatrix} j-1 \\ 2k+1 \end{bmatrix} \begin{bmatrix} 2m+2k+2 \\ m+j-1 \end{bmatrix} \frac{(x^2; x^2)_k}{(x^{2m+4}; x^2)_k} \frac{x^{(j-2k-2)}}{1+x^{m+1}} = d_{j-1}$$

$$(1 \leq j \leq r-1).$$

To simplify the notation a little, we put  $j - 1 = n$ . Moreover we note that because of the inequality  $j \leq r - 1$ , the summations in (16) and (17) can be extended from  $k = 0$  to  $\infty$  without affecting the left sides. Indeed outside the indicated  $k$ -ranges, we have  $[j-1-2k] = 0$  in (16), and  $[j-1-2k+1] = 0$  in (17). The restriction  $j \leq r - 1$  then becomes irrelevant. Thus we wish to prove that

$$(18) \quad \sum_{k \geq 0} \frac{\begin{bmatrix} n \\ 2k \end{bmatrix} \begin{bmatrix} 2m+2k+1 \\ m+n \end{bmatrix}}{\begin{bmatrix} 2m+1 \\ n \end{bmatrix}} \frac{(x; x^2)_k}{(x^{2m+3}; x^2)_k} x^{(n-\frac{3}{2}k)} = 1$$

and

$$(19) \quad \sum_{k \geq 0} \begin{bmatrix} n \\ 2k+1 \end{bmatrix} \begin{bmatrix} 2m+2k+2 \\ m+n \end{bmatrix} \frac{(x^2; x^2)_k x^{(n-\frac{3}{2}k-1)}}{(x^{2m+4}; x^2)_k (1+x^{m+1})}$$

$$= d_n = \sum_{\nu=1}^n \begin{bmatrix} 2m+1 \\ m+\nu \end{bmatrix} x^{\binom{\nu}{2}}$$

for all  $n \geq 0$ . Professor Andrews has pointed out to me that (18) and (19) can be derived from Saalschütz's summation of basic hypergeometric series [5, p. 247]. To keep the presentation self-contained, however, we will give direct proofs. Let  $F(m, n)$  and  $G(m, n)$  denote the left sides of (18) and (19) respectively. When  $n = 0$  or  $1$ , the only non-vanishing term of the series in (18) is the one with  $k = 0$ . Hence

$$F(m, 0) = \begin{bmatrix} 2m+1 \\ m \end{bmatrix} / \begin{bmatrix} 2m+1 \\ m \end{bmatrix} = 1$$

and

$$F(m, 1) = \begin{bmatrix} 2m+1 \\ m+1 \end{bmatrix} / \begin{bmatrix} 2m+1 \\ m \end{bmatrix} = 1.$$

Thus to complete the proof of (18) it suffices to show that  $F(m, n) = F(m, n + 2)$  for all  $n \geq 0$ . It is easy to verify that

$$(20) \quad \begin{bmatrix} 2m+2k+1 \\ m+n \end{bmatrix} = x^{2n-2k+1} \begin{bmatrix} 2m+2k+1 \\ m+n+2 \end{bmatrix}$$

$$+ \frac{1-x^{2n-2k+1}}{1-x^{2m+2k+3}} \begin{bmatrix} 2m+2k+3 \\ m+n+2 \end{bmatrix}.$$

Since

$$\binom{n-2k}{2} + 2n - 2k + 1 = \binom{n+2-2k}{2} + 2k,$$

it follows from (20) that

$$\begin{aligned} \left[ \begin{matrix} 2m+1 \\ m \end{matrix} \right] F(m, n) &= \sum_{k \geq 0} \left[ \begin{matrix} n \\ 2k \end{matrix} \right] \left[ \begin{matrix} 2m+2k+1 \\ m+n+2 \end{matrix} \right] \frac{(x; x^2)_k}{(x^{2m+3}; x^2)_k} x^{(n+\frac{n-2k}{2})+2k} \\ &+ \sum_{k \geq 0} \left[ \begin{matrix} n \\ 2k \end{matrix} \right] \left[ \begin{matrix} 2m+2k+3 \\ m+n+2 \end{matrix} \right] \frac{(x; x^2)_k (1-x^{2n-2k+1})}{(x^{2m+3}; x^2)_{k+1}} x^{(\frac{n-2k}{2})}. \end{aligned}$$

In the second sum we replace  $k$  by  $k-1$ , thus obtaining

$$\begin{aligned} \left[ \begin{matrix} 2m+1 \\ m \end{matrix} \right] F(m, n) &= \sum_{k \geq 0} \left[ \begin{matrix} n \\ 2k \end{matrix} \right] \left[ \begin{matrix} 2m+2k+1 \\ m+n+2 \end{matrix} \right] \frac{(x; x^2)_k}{(x^{2m+3}; x^2)_k} x^{(n+\frac{n-2k}{2})} \\ &+ \sum_{k \geq 1} \left[ \begin{matrix} n \\ 2k-2 \end{matrix} \right] \left[ \begin{matrix} 2m+2k+1 \\ m+n+2 \end{matrix} \right] \frac{(x; x^2)_{k-1} (1-x^{2n-2k+3})}{(x^{2m+3}; x^2)_k} x^{(n+\frac{n-2k}{2})} \\ &= \sum_{k \geq 0} \left\{ \left[ \begin{matrix} n \\ 2k \end{matrix} \right] x^{2k} + \left[ \begin{matrix} n \\ 2k-2 \end{matrix} \right] \frac{1-x^{2n-2k+3}}{1-x^{2k-1}} \right\} \\ &\cdot \left[ \begin{matrix} 2m+2k+1 \\ m+n+2 \end{matrix} \right] \frac{(x; x^2)_k}{(x^{2m+3}; x^2)_k} x^{(n+\frac{n-2k}{2})}. \end{aligned}$$

Since

$$(21) \quad \left[ \begin{matrix} n \\ 2k \end{matrix} \right] x^{2k} + \left[ \begin{matrix} n \\ 2k-2 \end{matrix} \right] \frac{1-x^{2n-2k+3}}{1-x^{2k}} = \left[ \begin{matrix} n+2 \\ 2k \end{matrix} \right],$$

we have

$$\begin{aligned} \left[ \begin{matrix} 2m+1 \\ m \end{matrix} \right] F(m, n) &= \sum_{k \geq 0} \left[ \begin{matrix} n+2 \\ 2k \end{matrix} \right] \left[ \begin{matrix} 2m+2k+1 \\ m+n+2 \end{matrix} \right] \frac{(x; x^2)_k}{(x^{2m+3}; x^2)_k} x^{(n+\frac{n-2k}{2})} \\ &= \left[ \begin{matrix} 2m+1 \\ m \end{matrix} \right] F(m, n+2), \end{aligned}$$

completing the proof.

The treatment of (19) is similar. First of all, when  $n = 0$  the terms of (19) all vanish, so  $G(m, 0) = 0$ . When  $n = 1$ , only the term  $k = 0$  of (19) is non-zero, so

$$(1+x^m)G(m, 1) = \left[ \begin{matrix} 2m+2 \\ m+1 \end{matrix} \right] = \frac{1-x^{2m+2}}{1-x^{m+1}} \left[ \begin{matrix} 2m+1 \\ m \end{matrix} \right] = (1+x^{m+1})d_1.$$

Hence it suffices to show that

$$G(m, n+2) = G(m, n) + \begin{bmatrix} 2m+1 \\ m+n+1 \end{bmatrix} x^{\binom{n+1}{2}} + \begin{bmatrix} 2m+1 \\ m+n+2 \end{bmatrix} x^{\binom{n+2}{2}}$$

for all  $n > 0$ . The relevant analogue of (20) is

$$\begin{aligned} \begin{bmatrix} 2m+2k+2 \\ m+n \end{bmatrix} &= x^{2n-2k} \begin{bmatrix} 2m+2k+2 \\ m+n+2 \end{bmatrix} \\ &+ \frac{1-x^{2n-2k}}{1-x^{2m+2k+4}} \begin{bmatrix} 2m+2k+4 \\ m+n+2 \end{bmatrix}. \end{aligned}$$

Using this, we split the series for  $G(m, n)$  into a sum of two series, and replace  $k$  by  $k-1$  in the second of these. This yields

$$\begin{aligned} G(m, n) &= \sum_{k \geq 0} \left\{ \begin{bmatrix} n \\ 2k+1 \end{bmatrix} x^{2k+1} + \begin{bmatrix} n \\ 2k-1 \end{bmatrix} \frac{1-x^{2n-2k+2}}{1-x^{2k}} \right\} \\ &\cdot \begin{bmatrix} 2m+2k+2 \\ m+n+2 \end{bmatrix} \frac{(x^2; x^2)_k x^{\binom{n+1}{2}-2k}}{(x^{2m+4}; x^2)_k (1+x^{m+1})}, \end{aligned}$$

where the second term in the curly bracket is to be interpreted as 0 when  $k=0$ . In analogy with (21) we have

$$\begin{bmatrix} n \\ 2k+1 \end{bmatrix} x^{2k+1} + \begin{bmatrix} n \\ 2k-1 \end{bmatrix} \frac{1-x^{2n-2k+2}}{1-x^{2k}} = \begin{cases} \begin{bmatrix} n+2 \\ 2k+1 \end{bmatrix} & \text{if } k > 0, \\ \begin{bmatrix} n \\ 1 \end{bmatrix} & \text{if } k = 0. \end{cases}$$

Therefore

$$\begin{aligned} (22) \quad G(m, n) &= \sum_{k \geq 0} \begin{bmatrix} n+2 \\ 2k+1 \end{bmatrix} \begin{bmatrix} 2m+2k+2 \\ m+n+2 \end{bmatrix} \frac{(x^2; x^2)_k x^{\binom{n+1}{2}-2k}}{(x^{2m+3}; x^2)_k (1+x^{m+1})} \\ &+ \left( \begin{bmatrix} n \\ 1 \end{bmatrix} x - \begin{bmatrix} n+2 \\ 1 \end{bmatrix} \right) \begin{bmatrix} 2m+2 \\ m+n+2 \end{bmatrix} \frac{x^{\binom{n+1}{2}}}{1+x^{m+1}}. \end{aligned}$$

It is easily checked that

$$\begin{aligned} &\left( \begin{bmatrix} n \\ 1 \end{bmatrix} x - \begin{bmatrix} n+2 \\ 1 \end{bmatrix} \right) \begin{bmatrix} 2m+2 \\ m+n+2 \end{bmatrix} \frac{x^{\binom{n+1}{2}}}{1+x^{m+1}} \\ &= - \begin{bmatrix} 2m+1 \\ m+n+1 \end{bmatrix} x^{\binom{n+1}{2}} - \begin{bmatrix} 2m+1 \\ m+n+2 \end{bmatrix} x^{\binom{n+2}{2}}. \end{aligned}$$

Hence from (22) we get

$$G(m, n) = G(m, n + 2) - \begin{bmatrix} 2m + 1 \\ m + n + 1 \end{bmatrix} x^{\binom{m+1}{2}} - \begin{bmatrix} 2m + 1 \\ m + n + 2 \end{bmatrix} x^{\binom{n+2}{2}},$$

completing the proof of (19).

Identities (18) and (19) also enable us to determine the entries which appear in the lower right corners when the clearing process is applied to (11) and (13). In the case of (11), the process subtracts from  $d_{r-1}$  all the non-zero terms of the series on the left of (19) with  $n = r - 1$ , except for the term with  $k = (r - 2)/2$ . In view of (19), the resulting entry in the lower right corner is the term with  $n = r - 1, k = (r - 2)/2$ , viz.

$$\begin{aligned} & q \begin{bmatrix} r - 1 \\ r - 1 \end{bmatrix} \begin{bmatrix} 2m + r \\ m + r - 1 \end{bmatrix} \frac{(x^2; x^2)_{(r-2)/2} x^{\binom{r}{2}}}{(x^{2m+4}; x^2)_{(r-2)/2} (1 + x^{m+1})} \\ &= \begin{bmatrix} 2m + r \\ m + r - 1 \end{bmatrix} \frac{(x^2; x^2)_{(r-2)/2}}{(x^{2m+4}; x^2)_{(r-2)/2} (1 + x^{m+1})}. \end{aligned}$$

Hence for even  $r$  we have

$$\begin{aligned} \det D_r &= \prod_{i=1}^{r-1} A_{ii} \begin{bmatrix} 2m + r \\ m + r - 1 \end{bmatrix} \frac{(x^2; x^2)_{(r-2)/2}}{(x^{2m+4}; x^2)_{(r-2)/2} (1 + x^{m+1})} \\ &= \prod_{i=1}^{r-1} \frac{\begin{bmatrix} 2m + i \\ m + i - 1 \end{bmatrix}}{\begin{bmatrix} m + i \\ i - 1 \end{bmatrix}} \begin{bmatrix} 2m + r \\ m + r - 1 \end{bmatrix} \frac{(x^2; x^2)_{(r-2)/2}}{(x^{2m+4}; x^2)_{(r-2)/2} (1 + x^{m+1})} \end{aligned}$$

by (12). Denote this last expression by  $f(r)$ . For even  $r, B_r(x | I_m)$  is the Pfaffian of  $D_r$ , and hence  $B_r(x | I_m) = \sqrt{f(r)}$ . Now

$$f(2) = \begin{bmatrix} 2m + 1 \\ m \end{bmatrix} \begin{bmatrix} 2m + 2 \\ m + 1 \end{bmatrix} \frac{1 - x^{m+1}}{1 - x^{2m+2}} = \begin{bmatrix} 2m + 1 \\ m \end{bmatrix}^2.$$

Hence  $B_2(x | I_m) = \sqrt{f(2)} = \begin{bmatrix} 2m + 1 \\ m \end{bmatrix}$ . On the other hand, when  $r = 2$  the right hand side of equation (1) telescopes to

$$\frac{(1 - x^{m+2})(1 - x^{m+3}) \cdots (1 - x^{2m+1})}{(1 - x)(1 - x^2) \cdots (1 - x^m)} = \begin{bmatrix} 2m + 1 \\ m \end{bmatrix}.$$

This proves the Bender-Knuth formula for  $r = 2$ . We proceed by induction to prove it for all even  $r$ . Clearly

$$\frac{f(r+2)}{f(r)} = \frac{\begin{bmatrix} 2m+r+1 \\ m+r \end{bmatrix} \begin{bmatrix} 2m+r+2 \\ m+r+1 \end{bmatrix}}{\begin{bmatrix} m+r \\ r-1 \end{bmatrix} \begin{bmatrix} m+r+1 \\ r \end{bmatrix}} \frac{1-x^r}{1-x^{2m+r+2}}.$$

After some straightforward cancellation, the right side reduces to

$$\prod_{\nu=1}^m \frac{(1-x^{m+r+\nu+1})^2}{(1-x^{r+\nu})^2},$$

from which we conclude that

$$(23) \quad \frac{B_{r+2}(x | I_m)}{B_r(x | I_m)} = \prod_{\nu=1}^m \frac{1-x^{m+r+\nu+1}}{1-x^{r+\nu}}.$$

On the other hand, if the right side of (1) is denoted by  $h(r)$ , then

$$\frac{h(r+2)}{h(r)} = \prod_{i=1}^m \prod_{j=1}^m \frac{1-x^{r+i+j+1}}{1-x^{r+i+j-1}}.$$

Here we have essentially the same telescope as the one mentioned above. The surviving factors are

$$\frac{(1-x^{r+m+2})(1-x^{r+m+3}) \cdots (1-x^{r+2m+1})}{(1-x^{r+1})(1-x^{r+2}) \cdots (1-x^{r+m})},$$

which is the right side of (23). This completes the induction (through even values of  $r$ ).

We can deal similarly with the case of odd  $r$ . When the clearing process is applied to (13), it subtracts from the 1 in the lower right corner all the non-zero terms of the series (18) with  $n = r - 1$  except the term with  $k = (r - 1)/2$ . Hence the resulting entry in the lower right corner is just this missing term, viz.

$$\frac{\begin{bmatrix} 2m+r \\ m+r-1 \end{bmatrix} (x; x^2)_{(r-1)/2}}{\begin{bmatrix} 2m+1 \\ m \end{bmatrix} (x^{2m+3}; x^2)_{(r-1)/2}}.$$

Thus for odd  $r$  we have

$$\det D'_r = s^2 \prod_{i=1}^{r-1} \frac{\begin{bmatrix} 2m+i \\ m+i-1 \end{bmatrix} \begin{bmatrix} 2m+r \\ m+r-1 \end{bmatrix} (x; x^2)_{(r-1)/2}}{\begin{bmatrix} m+i \\ i-1 \end{bmatrix} \begin{bmatrix} 2m+r \\ m \end{bmatrix} (x^{2m+3}; x^2)_{(r-1)/2}}.$$

If we temporarily denote the right hand side of this equation by  $g(r)$ , we have

$$B_r(x | I_m) = \sqrt{g(r)} \quad \text{for odd } r.$$

The proof that  $\sqrt{g(r)}$  is equal to the right side of (1) is completely analogous to the one given above for  $\sqrt{f(r)}$ , so can be omitted here.

#### REFERENCES

- [1] G. E. Andrews, *Plane partitions I. The MacMahon conjecture*, *Advances in Math. Suppl. Studies*, **1** (1978), 131–150.
- [2] G. E. Andrews, *Plane partitions II. The equivalence of the Bender-Knuth and MacMahon conjectures*, *Pacific J. Math.*, **72** (1977), 283–291.
- [3] E. A. Bender and D. E. Knuth, *Enumeration of plane partitions*, *J. Combinatorial Theory*, **13** (1972), 40–54.
- [4] B. Gordon and L. Houten, *Notes on plane partitions II*, *J. Combinatorial Theory*, **4** (1968), 81–99.
- [5] P. A. MacMahon, *Partitions of numbers whose graphs possess symmetry*, *Trans. Cambridge Phil. Soc.*, **17** (1898–99), 149–170.
- [6] L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge, 1961.

Received October 30, 1981

UNIVERSITY OF CALIFORNIA  
LOS ANGELES, CA 90024





# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California  
Los Angeles, CA 90024

HUGO ROSSI

University of Utah  
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR OGUS

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90089-1113

R. FINN and H. SAMELSON

Stanford University  
Stanford, CA 94305

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH  
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

Waleed A. Al-Salam and A. Verma, $q$ -Konhauser polynomials	1
Alfred David Andrew, The Banach space $JT$ is primary	9
Thomas E. Bengtson, Bessel functions on $P_n$	19
Joaquim Bruna Floris and Francesc Tugores, Free interpolation for holomorphic functions regular to the boundary	31
Peter Dierolf and Susanne Dierolf, Topological properties of the dual pair $\langle \mathfrak{B}(\Omega)', \mathfrak{B}(\Omega)'' \rangle$	51
Gerald Arthur Edgar, An ordering for the Banach spaces	83
Basil Gordon, A proof of the Bender-Knuth conjecture	99
Harold T. Hodes, A minimal upper bound on a sequence of Turing degrees which represents that sequence	115
Kuo Shih Kao, A note on $M_1$ -spaces	121
Frank Kost, Topological extensions of product spaces	129
Eva Lowen-Colebunders, On the convergence of closed and compact sets	133
Doron Lubinsky, Divergence of complex rational approximations	141
Warren May and Elias Hanna Toubassi, Endomorphisms of rank one mixed modules over discrete valuation rings	155
Richard Patrick Morton, The quadratic number fields with cyclic 2-classgroups	165
Roderic Murufas, Rank of positive matrix measures	177
Helga Schirmer, Fixed point sets of homotopies	191
E. Taflin, Analytic linearization of the Korteweg-de Vries equation	203
James Thomas Vance, Jr., $L^p$ -boundedness of the multiple Hilbert transform along a surface	221
Hiroshi Yamaguchi, A property of some Fourier-Stieltjes transforms	243