

Pacific Journal of Mathematics

ON UNITS OF PURE QUARTIC NUMBER FIELDS

AKIRA ENDÔ

ON UNITS OF PURE QUARTIC NUMBER FIELDS

AKIRA ENDÔ

Let $K = Q(\sqrt[4]{D^4 \pm d})$ be a pure quartic number field, where D and d are natural numbers such that d divides D^3 and d is fourth power free.

Then $\varepsilon = \pm(\sqrt[4]{D^4 \pm d} + D)/(\sqrt[4]{D^4 \pm d} - D)$ is a unit of K whose relative norm to the quadratic subfield of K is 1. We consider the condition for ε to be a member of a system of fundamental units of K .

1. Introduction. There have been many investigations concerning units of pure extensions of the rational number field of degree $n > 2$ generated by $\sqrt[n]{D^n \pm d}$, where D and d are natural numbers satisfying certain conditions ([2], [4], [7], [9], etc.). In general, suppose d divides D^{n-1} or, if n is a power of a prime number p , d divides pD^{n-1} . Then the numbers

$$\varepsilon_k = \frac{\omega^k - D^k}{(\omega - D)^k}, \quad \omega = \sqrt[n]{D^n \pm d},$$

where k runs over all the divisors of n except 1, are units and, moreover, independent in the real algebraic number field generated by ω [1], [2], [4]. (The proof of independence of the ε_k 's given by Halter-Koch and Stender [4] is incomplete. But the proof can be corrected by a slight modification.) When $n = 3, 4$ or 6 , the number of such divisors is equal to the rank of the unit group of the field $Q(\omega)$, where Q denotes the rational number field. In this paper we shall treat these units in the case $n = 4$.

The following result is established by Stender [8], [9]:

Let D and d be two natural numbers such that $d|2D^3$, and put $A = D^4 \pm d$ and $\omega = \sqrt[4]{A}$. Suppose that d is fourth power free and A/d or $2A/d$ is square free, according as $d|D^3$ or $d|2D^3$. Then

$$\varepsilon_2 = \pm \frac{\omega + D}{\omega - D} \quad \text{and} \quad \varepsilon_4 = \begin{cases} \frac{d}{(\omega - D)^4} & \text{if } d \text{ is not a square,} \\ \frac{\sqrt{d}}{(\omega - D)^2} & \text{if } d \text{ is a square and } d \neq 1, \\ \pm \frac{1}{\omega - D} & \text{if } d = 1 \end{cases}$$

form a system of fundamental units of $Q(\omega)$, except for the three cases $\omega^k = 8 = 2^4 - 8$, $\omega^4 = 12 = 2^4 - 4$ and $\omega^4 = 20 = 2^4 + 4$.

In this paper we shall remove the above assumption on A/d , and study the properties of ε_2 .

2. Known facts. First, we state a few known facts on units of a pure quartic number field. Let A be a natural number which is fourth power free; then we can write $A = fg^2h^3$ with natural numbers f, g, h such that fgh is square free. We suppose $fh \neq 1$. Then the pure quartic number field $K = Q(\sqrt[4]{A})$ generated by $\sqrt[4]{A}$ contains a unique quadratic subfield, namely $Q(\sqrt{fh})$. Any integer α of K is of the form

$$\alpha = \frac{1}{k} \left(x_0 + x_1 \sqrt[4]{fg^2h^3} + x_2 \sqrt{fh} + x_3 \sqrt[4]{f^3g^2h} \right)$$

with rational integers x_0, x_1, x_2, x_3 and $k = 1, 2$ or 4 , and its conjugate relative to $Q(\sqrt{fh})$ is

$$\alpha' = \frac{1}{k} \left(x_0 - x_1 \sqrt[4]{fg^2h^3} + x_2 \sqrt{fh} - x_3 \sqrt[4]{f^3g^2h} \right).$$

Now let $\varepsilon_0 > 1$ be the smallest unit of K such that $\varepsilon_0 \varepsilon'_0 = 1$, and $\varepsilon^* > 0$ the fundamental unit of $Q(\sqrt{fh})$.

LEMMA 1 ([5], [6]). ε_0 and ε^* or ε_0 and $\sqrt{\varepsilon^* \varepsilon_0}$ form a system of fundamental units of K ; the former case occurs if and only if neither ε^* nor $-\varepsilon^*$ is the norm of a unit of K to $Q(\sqrt{fh})$.

In any case, ε_0 appears as a member of a system of fundamental units of K . The following result will aid in determining ε_0 :

LEMMA 2 ([6]). Let A_1 and A_2 be two positive rational integers such that $Q(\sqrt[4]{A_1 A_2^3}) = Q(\sqrt[4]{A})$. Then the indeterminate equation

$$A_1 x^4 - A_2 y^4 = \pm C \quad \text{with } C = 1, 2, 4$$

has at most one positive integer solution. If (a, b) is a positive integer solution of this equation, then $\pm(a\sqrt[4]{A_1} + b\sqrt[4]{A_2})/(a\sqrt[4]{A_1} - b\sqrt[4]{A_2})$ is a unit of $Q(\sqrt[4]{A})$ whose relative norm to $Q(\sqrt{A})$ is 1, and furthermore is equal to ε_0 or ε_0^2 with the only two exceptions $x^4 - 5y^4 = 1$ and $4x^4 - 3y^4 = 1$.

3. Theorems. From now on, we take A so that $K = Q(\sqrt[4]{A}) = Q(\sqrt[4]{D^4 \pm d})$, and suppose that $d \mid D^3$ and d is fourth power free. Then there is a natural number u satisfying

$$u^4 A = D^4 \pm d.$$

We write $d = d_1 d_2^2 d_3^3$ with natural numbers d_1, d_2, d_3 such that $d_1 d_2 d_3$ is square free. It is easy to see that $d_1 \mid f, d_2 \mid g, d_3 \mid h$.

Now we write $\sqrt[4]{A} = \omega$ and put

$$\epsilon_2 = \pm \frac{u\omega + D}{u\omega - D},$$

which is a unit of K . In the special case where $u = 1$ and A/d is square free, i.e. $g = d_2, h = d_3$, as already mentioned in the introduction, Stender's result [8], [9] states that ϵ_2 is contained in a system of fundamental units of K with the exception of three cases. Moreover [3],

$$\epsilon^* = \begin{cases} \frac{d}{(\omega^2 - D^2)^2}, & d_1 d_3 \neq 1, \\ \pm \frac{\sqrt{d}}{\omega^2 - D^2}, & d = d_2^2, \end{cases}$$

and $\pm \epsilon^*$ are the norms of no unit of K to $Q(\sqrt{fh})$.

Since

$$Q(\sqrt[4]{D^4 \pm d}) = Q(\sqrt[4]{D'^4 \mp d'}),$$

where $D' = u(f/d_1)(g/d_2)(h/d_3)$, $d' = (f/d_1)^3(g/d_2)^2(h/d_3)$, $d' \mid D'^3$ and d' is fourth power free, we treat below exclusively the plus case, i.e. $u^4 A = D^4 + d$. We write simply $\epsilon_2 = \epsilon$:

$$\epsilon = \frac{u\omega + D}{u\omega - D} = \frac{1}{d} (2D^4 + d + 2D^3 u\omega + 2D^2 u^2 \omega^2 + 2Du^3 \omega^3).$$

Obviously $\epsilon\epsilon' = 1$. We then consider whether there exists a unit η of K such that $\eta\eta' = 1$ and $\epsilon = \eta^2$.

Let

$$\eta = \frac{1}{k} \left(x_0 + x_1 \sqrt[4]{fg^2h^3} + x_2 \sqrt{fh} + x_3 \sqrt[4]{f^3g^2h} \right)$$

be a unit of K with $\eta\eta' = 1$. Then

$$(1) \quad x_0^2 + x_2^2 fh - 2x_1 x_3 fgh = k^2,$$

$$(2) \quad x_1^2gh + x_3^2fg - 2x_0x_2 = 0,$$

and

$$\eta^2 = \frac{1}{k^2} \left(x_0^2 + x_1^2fh + 2x_1x_3fgh + 2(x_0x_1 + x_2x_3f)\sqrt[4]{fg^2h^3} \right. \\ \left. + (x_1^2gh + x_3^2fg + 2x_0x_2)\sqrt{fh} + 2(x_0x_3 + x_1x_2h)\sqrt[4]{f^3g^2h} \right).$$

Hence (1) and (2) imply that $\varepsilon = \eta^2$ if and only if

$$(3) \quad \frac{D^4}{d} = \frac{2}{k^2}x_1x_3fgh = \frac{1}{k^2}(x_0^2 + x_2^2fh) - 1,$$

$$(4) \quad \frac{D^3}{d}u = \frac{1}{k^2}(x_0x_1 + x_2x_3f),$$

$$(5) \quad \frac{D^2}{d}u^2gh = \frac{2}{k^2}x_0x_2 = \frac{1}{k^2}(x_1^2gh + x_3^2fg),$$

$$(6) \quad \frac{D}{d}u^3gh^2 = \frac{1}{k^2}(x_0x_3 + x_1x_2h).$$

From (3)–(6) we have

$$2x_0x_2(x_0x_1 + x_2x_3f)h = 2x_1x_3fgh(x_0x_3 + x_1x_2h).$$

It easily follows from this, together with (2), that

$$(x_0x_1 - x_2x_3f)(x_1^2h - x_3^2f) = 0,$$

from which, as $fh \neq 1$ is not a square,

$$(7) \quad x_0x_1 = x_2x_3f.$$

REMARK 1. It is easily shown that in the above situation the following facts hold:

$$k = 1 \quad \text{if } 4 \mid d, 2 \mid fh \text{ or } 2 \nmid fgh, \\ k = 2 \quad \text{if } 2 \nmid d \text{ and } 2 \mid g.$$

We prove here the following:

THEOREM 1. *Notations being as above, suppose that $u^4A = D^4 + d$ and $A \neq 5^3, 2^23^3$. Then $\varepsilon = \varepsilon_0$ or ε_0^2 , and moreover $\varepsilon = \varepsilon_0^2$ if and only if $A = d$ or $4d$ and either $2(u^2 + \sqrt{d/A})$ or $2(u^2 - \sqrt{d/A})$ is a square.*

Proof. It follows from Lemma 2 that $\varepsilon = \varepsilon_0$ or ε_0^2 . Suppose that $\varepsilon = \eta^2$ with $\eta\eta' = 1$ as above. Then from (3) we have

$$u^4A = u^4fg^2h^3 = D^4 + d = \frac{2}{k^2}x_1x_3fghd + d.$$

This implies $A = d$ or $4d$ because $d_1|f, d_2|g, d_3|h$, and if $2|fh, k = 1$ by Remark 1. Furthermore, from (3)–(7) we obtain

$$x_1 = \frac{k^2D^3u}{2d} \frac{1}{x_0}, \quad x_2 = \frac{k^2D^2u^2gh}{2d} \frac{1}{x_0}, \quad x_3 = \frac{D}{ufgh} x_0,$$

and

$$\frac{D}{d} u^3gh^2 = \frac{1}{k^2} \left(\frac{D}{ufgh} x_0^2 + \frac{k^4D^5u^3gh}{4d^2} \frac{1}{x_0} \right).$$

From the last equation we have

$$\begin{aligned} 0 &= x_0^4 - \frac{k^2u^4A}{d} x_0^2 + \frac{k^4u^4D^4A}{4d^2} \\ &= \begin{cases} \left(x_0^2 - \frac{k^2u^2(u^2 + 1)}{2} \right) \left(x_0^2 - \frac{k^2u^2(u^2 - 1)}{2} \right), & A = d, \\ \left(x_0^2 - k^2u^2(2u^2 + 1) \right) \left(x_0^2 - k^2u^2(2u^2 - 1) \right), & A = 4d. \end{cases} \end{aligned}$$

Since x_0 is a rational integer, $(u^2 \pm 1)/2$ or $2u^2 \pm 1$ must be a square, according as $A = d$ or $A = 4d$. Conversely, if these conditions are satisfied, then

$$x_0 = kuw, \quad x_1 = \frac{kD^3}{2d} \frac{1}{v}, \quad x_2 = \frac{kD^2ugh}{2d} \frac{1}{v}, \quad x_3 = \frac{kD}{fgh} v,$$

where

$$v = \begin{cases} \sqrt{\frac{u^2 \pm 1}{2}}, & A = d, \\ \sqrt{2u^2 \pm 1}, & A = 4d, \end{cases}$$

satisfy conditions (1)–(6). Thus the theorem follows.

REMARK 2. In the above theorem, $u \neq 1$ if $A = d$. Since the fundamental unit of the real quadratic number field $Q(\sqrt{2})$ is $1 + \sqrt{2}$, the natural numbers u such that $(u^2 \pm 1)/2$ is a square are given by $u + \sqrt{u^2 + 1} = (1 + \sqrt{2})^{2l+1}$ or $u + \sqrt{u^2 - 1} = (1 + \sqrt{2})^{2l}$ for some $l \geq 1$.

Moreover, the natural numbers u such that $2u^2 \pm 1$ is a square are given by $\sqrt{2u^2 + 1} + u\sqrt{2} = (1 + \sqrt{2})^{2l}$ or $\sqrt{2u^2 - 1} + u\sqrt{2} = (1 + \sqrt{2})^{2l-1}$ for some $l \geq 1$.

In the minus case we have the following:

THEOREM 2. *Suppose $u^4A = D^4 - d$ and $A \neq 5, 2^2 \cdot 3$. Then $\varepsilon = \varepsilon_0$ or ε_0^2 , and $\varepsilon = \varepsilon_0^2$ if and only if $d = 1$ or 4 and either $2(D^2 + \sqrt{d})$ or $2(D^2 - \sqrt{d})$ is a square.*

Proof. Immediate from Theorem 1 and the remark at the beginning of this section.

REMARK 3. In the above theorem, $D \neq 1$ if $d = 1$. The natural numbers D such that $D^2/2 \pm 1$ is a square are given by $\sqrt{2(D^2 + 2)} + D\sqrt{2} = 2(1 + \sqrt{2})^{2l}$ or $\sqrt{2(D^2 - 2)} + D\sqrt{2} = 2(1 + \sqrt{2})^{2l-1}$ for some $l \geq 1$.

COROLLARY ([9]). *If $A = D^4 \pm d$ and A/d is square free, there exists no unit η of K such that $\eta\eta' = 1$ and $\varepsilon = \eta^2$, with the single exception of $A = 12 = 2^4 - 4$.*

Proof. By Theorem 1 such a unit cannot exist in the plus case, and hence Theorem 2 shows that $d = 1$ or 4 . Then, from the assumption, we have $A = 4f = D^4 - d = 12$, namely $D = 2, d = 4$, which gives the only exception stated above.

REMARK 4. Stender [10] has obtained some sufficient conditions for $\varepsilon = \varepsilon_0$, which can also be deduced from Theorems 1 and 2.

REFERENCES

- [1] L. Bernstein, *Der Hasse-Bernsteinsche Einheitensatz für den verallgemeinerten Jacobi-Perronschen Algorithmus*, Abh. Math. Sem. Univ. Hamburg, **43** (1975), 11–20.
- [2] L. Bernstein und H. Hasse, *Einheitenberechnung mittels des Jacobi-Perronschen Algorithmus*, J. Reine Angew. Math., **218** (1965), 51–69.
- [3] G. Degert, *Über die Bestimmung der Grundeinheit gewisser reell-quadratischer Zahlkörper*, Abh. Math. Sem. Univ. Hamburg, **22** (1958), 92–97.
- [4] F. Halter-Koch und H.-J. Stender, *Unabhängige Einheiten für die Körper $K = \mathbb{Q}(\sqrt[n]{D^n \pm d})$ mit $d \mid D^n$* , Abh. Math. Sem. Univ. Hamburg, **42** (1974), 33–40.
- [5] W. Ljunggren, *Über die Lösung einiger unbestimmten Gleichungen vierten Grades*, Avh. Norske Vid.-Akad. Oslo (I), 1934 No. 14, 1–35.

- [6] ———, *Einige Eigenschaften der Einheiten reeller quadratischer und rein-biquadratischer Zahlkörper mit Anwendung auf die Lösung einer Klasse unbestimmter Gleichungen vierten Grades*, Skr. Norske Vid.-Akad. Oslo (I), 1936 No. 12, 1–73.
- [7] R. J. Rudman, *On the fundamental unit of a purely cubic field*, Pacific J. Math., **46** (1973), 253–256.
- [8] H.-J. Stenderr, *Grundeinheiten für einige unendliche Klassen reiner biquadratischer Zahlkörper mit einer Anwendung auf die diophantische Gleichung $x^4 - ay^4 = \pm c$ ($c = 1, 2, 4$ oder 8)*, J. Reine Angew. Math., **264** (1973), 207–220.
- [9] ———, *Eine Formel für Grundeinheiten in reinen algebraischen Zahlkörpern dritten, vierten und sechsten Grades*, J. Number Theory, **7** (1975), 235–250.
- [10] ———, *Lösbare Gleichungen $ax^n - by^n = c$ und Grundeinheiten für einige algebraische Zahlkörper vom Grade $n = 3, 4, 6$* , J. Reine Angew. Math., **290** (1977), 24–62.

Received April 2, 1981 and in revised form April 28, 1982.

KUMAMOTO UNIVERSITY
KUMAMOTO, JAPAN

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR OGUS

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90089-1113

R. FINN and H. SAMELSON

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$132.00 a year (6 Vol., 12 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics ISSN 0030-8730 is published monthly by the Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P. O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1983 by Pacific Journal of Mathematics

Pacific Journal of Mathematics

Vol. 109, No. 2

June, 1983

Tibor Bisztriczky , On the singularities of almost-simple plane curves	257
Peter B. Borwein , On Sylvester's problem and Haar spaces	275
Emilio Bujalance , Cyclic groups of automorphisms of compact nonorientable Klein surfaces without boundary	279
Robert Jay Daverman and John J. Walsh , Acyclic decompositions of manifolds	291
Lester Eli Dubins , Bernstein-like polynomial approximation in higher dimensions	305
Allan L. Edelson and Jerry Dee Schuur , Nonoscillatory solutions of $(rx^n)^n \pm f(t, x)x = 0$	313
Akira Endô , On units of pure quartic number fields	327
Hector O. Fattorini , A note on fractional derivatives of semigroups and cosine functions	335
Ronald Fintushel and Peter Sie Pao , Circle actions on homotopy spheres with codimension 4 fixed point set	349
Stephen Michael Gagola, Jr. , Characters vanishing on all but two conjugacy classes	363
Saverio Giulini , Singular characters and their L^p norms on classical Lie groups	387
Willy Govaerts , Locally convex spaces of non-Archimedean valued continuous functions	399
Wu-Chung Hsiang and Bjørn Jahren , A remark on the isotopy classes of diffeomorphisms of lens spaces	411
Hae Soo Oh , Compact connected Lie groups acting on simply connected 4-manifolds	425
Frank Okoh and Frank A. Zorzitto , Subsystems of the polynomial system	437
Knut Øyma , An interpolation theorem for H_E^∞	457
Nikolaos S. Papageorgiou , Nonsmooth analysis on partially ordered vector spaces. II. Nonconvex case, Clarke's theory	463