

Pacific Journal of Mathematics

**AN INTERPOLATION THEOREM FOR ANALYTIC FAMILIES
OF OPERATORS ACTING ON CERTAIN H^p SPACES**

EUGENIO HERNANDEZ

AN INTERPOLATION THEOREM FOR ANALYTIC FAMILIES OF OPERATORS ACTING ON CERTAIN H^p SPACES

EUGENIO HERNÁNDEZ

The main objective of this paper is to obtain an interpolation theorem for families of operators acting on atomic H^p spaces, $0 < p \leq 1$. We prove that if $0 < p_0 < p_1 \leq 1$ and $\{T_z\}$, $z \in \bar{S} = \{z \in \mathbb{C} / 0 \leq \text{Real } z \leq 1\}$, is an analytic and admissible family of linear transformations such that $T_{j+i\gamma}$ maps H^{p_j} into L^{p_j} , where $-\infty < \gamma < \infty$, with norm not exceeding A_j , $j = 0, 1$, then for all θ , $0 \leq \theta \leq 1$, T_θ maps H^r into L^r with norm not exceeding $cA_0^{1-\theta}A_1^\theta$, where $1/r = (1-\theta)/p_0 + \theta/p_1$.

1. Introduction. For $x \in \mathbb{R}^n$ and $\alpha = (\alpha_1, \dots, \alpha_n)$ a multi-index, with α_j a natural number, $j = 1, \dots, n$, we write $x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ and $|\alpha| = \alpha_1 + \cdots + \alpha_n$. For $0 < p \leq 1$ and an integer $s \geq [n(1/p - 1)]^1$ we say that a function $a(x)$ is a (p, s) -atom if:

- (i) the support of a is contained in a ball $B \subset \mathbb{R}^n$.
- (ii) $|a(x)| \leq |B|^{-1/p}$ for all $x \in \mathbb{R}^n$ ($|B|$ denotes the measure of B).
- (iii) $\int_{\mathbb{R}^n} a(x)x^\alpha dx = 0$ for all multi-indexes α with $|\alpha| \leq s$.

The space $H^{p,s}(\mathbb{R}^n)$ consists of those linear functionals f defined on an appropriate Lipschitz class of test functions (see [6] for details, where more general atoms are also considered) that can be expressed in the form $f = \sum_{i=1}^\infty \lambda_i a_i$, where the a_i are (p, s) -atoms and $\sum_{i=1}^\infty |\lambda_i|^p < \infty$. The quasi-norm of f is defined to be the infimum of the numbers $(\sum_{i=1}^\infty |\lambda_i|^p)^{1/p}$ taken over all the above representations of f ; it is denoted by $\|f\|_{H^{p,s}}$. For fixed p , the spaces $H^{p,s}$ are independent of s and their quasi-norms are equivalent (for this and the relation of these spaces to the classical H^p spaces see [6]). For this reason we shall denote each of these equivalent spaces by H^p and $\|\cdot\|_{H^p}$ will denote the quasi-norm.

Another aspect of this theorem that we must make precise is the type of transformations we are dealing with.

A function Φ defined on \bar{S} , the closure of S , is said to be of *admissible growth* if there exist positive constants B and b , $b < \pi$, such that

$$\Phi(z) \leq B e^{b|\text{Im } z|}$$

for all $z \in \bar{S}$.

¹ $[t]$ denotes the integer part of the non-negative real number t .

A family of linear transformations $\{T_z\}$, $z \in \bar{S}$, each mapping simple functions to measurable functions on \mathbf{R}^n , is called *analytic* if the function

$$z \rightarrow \int_{\mathbf{R}^n} (T_z \varphi) \psi \, dx$$

is analytic for each pair of simple functions φ and ψ on \mathbf{R}^n .

This family is said to be *admissible* if the function $\log \|T_z \varphi\|_1$ is of admissible growth for each simple φ on \mathbf{R}^n .

We are now ready to state the main theorem.

THEOREM I. *Let $0 < p_0 < p_1 \leq 1$ and $\{T_z\}$, $z \in \bar{S}$, be an analytic and admissible family of linear transformations. We suppose*

$$(1.1) \quad \|T_{j+iy} f\|_{L^{p_j}} \leq A_j \|f\|_{H^{p_j}}, \quad j = 0, 1,$$

for all $f \in H^{p_j} \cap L^{p_j}$, where $-\infty < y < \infty$ and A_j , $j = 0, 1$, are positive constants.

Then, for each θ , $0 \leq \theta \leq 1$, we have

$$(1.2) \quad \|T_\theta f\|_{L^r} \leq c A_0^{1-\theta} A_1^\theta \|f\|_{H^r}$$

for all $f \in H^r$, where $1/r = (1 - \theta)/p_0 + \theta/p_1$ and c depends only on the dimension n and the quasi-norm used in H^r .

A similar theorem for transformations acting on H^p of the unit disk was proved by E. M. Stein and G. Weiss [5].²

2. Proof of Theorem I. Due to the equivalence of the spaces $H^{p,s}$ for p fixed, Theorem I follows from the theorem below.

THEOREM II. *Let $0 < p_0 < p_1 \leq 1$ and s an integer satisfying $s \geq [n(1/p_0 - 1)]$. Let $\{T_z\}$, $z \in \bar{S}$, be an analytic and admissible family of linear transformations such that*

$$(2.1) \quad \|T_{j+iy} f\|_{L^{p_j}} \leq A_j \|f\|_{H^{p_j, s}}, \quad j = 0, 1,$$

for all $f \in H^{p_j, s} \cap L^{p_j}$, where $-\infty < y < \infty$ and A_j , $j = 0, 1$, are positive constants.

² In [4] E. M. Stein proved a similar theorem for transformations acting on L^p spaces.

Then, for each $\theta, 0 \leq \theta \leq 1$, we have

$$(2.2) \quad \|T_\theta f\|_{L^r} \leq A_0^{1-\theta} A_1^\theta \|f\|_{H^{r,s}}$$

for all $f \in H^{r,s}$, where $1/r = (1 - \theta)/p_0 + \theta/p_1$.

Three lemmas are used to prove this theorem.

LEMMA 1. Let $\Phi: S \rightarrow \mathbf{R}$ be an upper semi-continuous function of admissible growth and subharmonic in S . Then for $z_0 = x_0 + iy_0 \in S$ we have

$$\begin{aligned} \Phi(z_0) &\leq \int_{-\infty}^{\infty} \Phi(i[y_0 + y])\omega(1 - x_0, y) dy \\ &\quad + \int_{-\infty}^{\infty} \Phi(1 + i[y_0 + y])\omega(x_0, y) dy, \end{aligned}$$

where

$$\omega(x, y) = \frac{1}{2} \frac{\sin \pi x}{\cos \pi x + \cosh \pi y}.$$

For a proof see [5].

LEMMA 2. If $V(z)$ is an analytic function mapping S into L^1 , then, for any positive $c \leq 1$, $\Phi(z) = \int_{\mathbf{R}^n} |V(z)|^c dx$ is continuous and $\log \Phi(z)$ is subharmonic.

The proof can be found in [5].

LEMMA 3. Let $0 < p_0 < p_1 \leq 1$ and s an integer satisfying $s \geq [n(1/p_0 - 1)]$. Let a be an (r, s) -atom, where $1/r = (1 - \theta)/p_0 + \theta/p_1$ and $0 \leq \theta \leq 1$.

Then there exists an analytic function $h(z)$ mapping S into L^1 such that

$$\|h(iy)\|_{H^{p_0,s}} \leq 1, \quad \|h(1 + iy)\|_{H^{p_1,s}} \leq 1$$

for any real y and $h(\theta) = a$.

Proof. Let B be the support of a and define $h(z) = |B|^{\alpha(z)} a$ where $\alpha(z) = (1/r - 1/p_0)(1 - z) + (1/r - 1/p_1)z$. Observe that

$$(2.2) \quad \alpha(\theta) = 0.$$

$$(2.3) \quad \text{Real}\{\alpha(iy)\} = 1/r - 1/p_0 \quad \text{for all real } y.$$

$$(2.4) \quad \text{Real}\{\alpha(1 + iy)\} = 1/r - 1/p_1 \quad \text{for all real } y.$$

Using the fact (easily verified) that a belongs to $H^{p_0, s}$ with quasi-norm not exceeding $|B|^{1/p_0 - 1/r}$ and (2.3), we obtain

$$\|h(iy)\|_{H^{p_0, s}} = |B|^{1/r - 1/p_0} \|a\|_{H^{p_0, s}} \leq 1.$$

Similarly, $\|h(1 + iy)\|_{H^{p_1, s}} \leq 1$. Finally, (2.2) implies $h(\theta) = a$. □

Proof of Theorem II. Let a be an (r, s) -atom and denote by $h(z)$ the analytic function associated with a as in Lemma 3. Choose an integer k such that $kp_0 > r$ and g a positive simple function with $\|g\|_{k'} \leq 1$ where $1/k + 1/k' = 1$. Consider

$$\Phi(z) = \int_{\mathbf{R}^n} [|g(x)]^{\beta(z)} |(T_z h(z))(x)|^{r/k} dx$$

where

$$\beta(z) = k' - \frac{k'r}{k} \left(\frac{1-z}{p_0} + \frac{z}{p_1} \right).$$

Note that

$$(2.5) \quad \beta(\theta) = 1.$$

$$(2.6) \quad \text{Real } \beta(iy) = \frac{k'(kp_0 - r)}{kp_0} \quad \text{for all real } y.$$

$$(2.7) \quad \text{Real } \beta(1 + iy) = \frac{k'(kp_1 - r)}{kp_1} \quad \text{for all real } y.$$

Using Hölder's inequality with index $kp_0/r > 1$, (2.6) and the fact that the $L^{k'}$ norm of g is bounded by 1, we obtain

$$|\Phi(iy)| \leq \left\{ \int_{\mathbf{R}^n} |(T_{iy} h(iy))(x)|^{p_0} dx \right\}^{r/kp_0}.$$

Using this inequality, hypothesis (2.1) and Lemma 3, we obtain

$$(2.8) \quad |\Phi(iy)| \leq A_0^{r/k} \|h(iy)\|_{H^{p_0, s}} \leq A_0^{r/k}.$$

Similarly, using Hölder's inequality with index $kp_1/r > 1$, we obtain

$$(2.9) \quad |\Phi(1 + iy)| \leq A_1^{r/k}.$$

Lemma 2 implies that $\log \Phi(z)$ is subharmonic. Consequently, Lemma 1 and inequalities (2.8) and (2.9) imply³

$$\log \Phi(\theta) \leq \log A_0^{r(1-\theta)/k} A_1^{r\theta/k}.$$

³ Observe that $\int_{-\infty}^{\infty} \omega(1 - \theta, y) dy = 1 - \theta$ and $\int_{-\infty}^{\infty} \omega(\theta, y) dy = \theta$.

Noticing that $\Phi(\theta) = \int_{\mathbf{R}^n} g(x) |(T_\theta a)(x)|^{r/k} dx$, the above inequality, together with

$$\left\{ \int_{\mathbf{R}^n} |(T_\theta a)(x)|^r dx \right\}^{1/k} = \sup \left\{ \int_{\mathbf{R}^n} |(T_\theta a)(x)|^{r/k} g(x) dx \mid g \geq 0, g \text{ simple}, \|g\|_k \leq 1 \right\},$$

implies

$$\left\{ \int_{\mathbf{R}^n} |(T_\theta a)(x)|^r dx \right\}^{1/k} \leq A_0^{r(1-\theta)/k} A_1^{r\theta/k},$$

which proves Theorem II for atoms. From here the theorem follows from the definition of $H^{r,s}$ in terms of atoms. \square

3. Remarks. We start with some historical remarks. H^p spaces can be defined on spaces of homogeneous type (see [1]). The main difference with the H^p spaces that we consider here is that (iii) in the definition of an atom is replaced by $\int a(x) dx = 0$. The reader will have no trouble verifying that Theorem I can be extended to include the homogeneous type case. In this context R. Macías proved in his dissertation (1973–1974) a particular case of Theorem I (see [3]). An interpolation theorem for linear operators acting on $H^p(\mathbf{R}^n)$ was published in 1974 by C. Fefferman, N. Rivière and Y. Sagher ([2]). Their proof uses the real method of interpolation and is quite different from Macías’ proof. For this reason, it is unfortunate that Macías’ dissertation was never published. Our work fills the gap left by the unpublished work of R. Macías.

It is worthwhile to notice that the fact proved by M. Taibleson and G. Weiss ([6]) concerning the equivalence of $H^{p,s}$ spaces for fixed p plays an important role in the proof of Theorem I.

Theorem I has several extensions. First, \mathbf{R}^n can be replaced by any other space for which an atomic theory similar to the one described above can be defined. Second, condition (1.1) can be replaced by

$$\|T_{j+iy} f\|_{L^{q_j}} \leq A_j(y) \|f\|_{H^{p_j}}, \quad j = 0, 1,$$

where $0 < q_j < \infty$, $\log A_j(y) \leq C_j e^{d_j|y|}$, $0 < d_j < \pi$ and $C_j > 0$, $j = 0, 1$. Then, it is easy to check that the conclusion of the theorem becomes

$$\|T_\theta f\|_{L^q} \leq C \|f\|_{H^r}$$

where C depends on θ , p_j , q_j , C_j , d_j ($j = 0, 1$) and the dimension n and $1/q = (1 - \theta)/q_0 + \theta/q_1$.

REFERENCES

- [1] R. Coifman and G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bull. Amer. Math. Soc., **83**, No. 4 (1977), 569–645.
- [2] C. Fefferman, N. Rivière and Y. Sagher, *Interpolation between H^p spaces, the real method*, Trans. Amer. Math. Soc., **19** (1974), 75–82.
- [3] R. A. Macias, *Interpolation theorems on generalized Hardy spaces*, Dissertation, Washington Univ., St. Louis, Missouri (1974).
- [4] E. M. Stein, *Interpolation of linear operators*, Trans. Amer. Math. Soc., **83** (1956), 482–492.
- [5] E. M. Stein and G. Weiss, *On the interpolation of analytic families of operators acting on H^p spaces*, Tôhoku Math. J., **8** (1957), 318–339.
- [6] M. H. Taibleson and G. Weiss, *The molecular characterization of certain Hardy spaces*, Astérisque, **77** (1980), 67–149.

Received March 8, 1982 and in revised form December 16, 1982. This work is part of the author's dissertation which was completed at Washington University in St. Louis. The author expresses deep appreciation to his advisors, Richard Rochberg and Guido Weiss, for their support.

WASHINGTON UNIVERSITY
ST. LOUIS, MO 63130

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)
University of California
Los Angeles, CA 90024

HUGO ROSSI
University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR OGUS
University of California
Berkeley, CA 94720

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, CA 90089-1113

R. FINN and H. SAMELSON
Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Wojciech Abramczuk , A class of surjective convolution operators	1
K. Adachi , Extending bounded holomorphic functions from certain subvarieties of a weakly pseudoconvex domain	9
Malvina Florica Baica , An algorithm in a complex field and its application to the calculation of units	21
Giuliana Bianchi and Robert Cori , Colorings of hypermaps and a conjecture of Brenner and Lyndon	41
Ronald James Evans , Determinations of Jacobsthal sums	49
Leslie Foged , Characterizations of \aleph -spaces	59
Nassif A. Ghoussoub and Paulette Saab , Weak compactness in spaces of Bochner integrable functions and the Radon-Nikodým property	65
J. Gómez Gil , On local convexity of bounded weak topologies on Banach spaces	71
Masaru Hara , On Gamelin constants	77
Wilfried Hauenschild, Eberhard Kaniuth and Ajay Kumar , Harmonic analysis on central hypergroups and induced representations	83
Eugenio Hernandez , An interpolation theorem for analytic families of operators acting on certain H^p spaces	113
Thomas Alan Keagy , On “Tauberian theorems via block-dominated matrices”	119
Thomas Landes , Permanence properties of normal structure	125
Daniel Henry Luecking , Closed ranged restriction operators on weighted Bergman spaces	145
Albert Thomas Lundell , The p -equivalence of $SO(2n + 1)$ and $Sp(n)$	161
Mark D. Meyerson , Remarks on Fenn’s “the table theorem” and Zaks’ “the chair theorem”	167
Marvin Victor Mielke , Homotopically trivial toposes	171
Gerard J. Murphy , Hyperinvariant subspaces and the topology on $\text{Lat } A$...	183
Subhashis Nag , On the holomorphy of maps from a complex to a real manifold	191
Edgar Milan Palmer and Robert William Robinson , Enumeration of self-dual configurations	203
John J. Walsh and David Clifford Wilson , Continuous decompositions into cells of different dimensions	223
Walter John Whiteley , Infinitesimal motions of a bipartite framework	233