

Pacific Journal of Mathematics

THE JORDAN DECOMPOSITION AND HALF-NORMS

DEREK W. ROBINSON AND SADAYUKI YAMAMURO

THE JORDAN DECOMPOSITION AND HALF-NORMS

DEREK W. ROBINSON AND SADAYUKI YAMAMURO

Let \mathfrak{B} be a Banach space, with norm $\|\cdot\|$, ordered by a positive cone \mathfrak{B}_+ and order the dual \mathfrak{B}^* by the dual cone \mathfrak{B}_+^* . We prove that, if \mathfrak{B} is orthogonally generated, each $f \in \mathfrak{B}^*$ has an orthogonal, and norm-unique, Jordan decomposition $f = f_+ - f_-$ with $f_{\pm} \in \mathfrak{B}^*$,

$$\|f\| = \|f_+\| + \|f_-\|,$$

if, and only if, the norm on \mathfrak{B} has the order theoretic property

$$\|a\| = \inf\{\lambda \geq 0; -\lambda u \leq a \leq \lambda v \text{ for some } u, v \in \mathfrak{B}_1\},$$

when \mathfrak{B}_1 is the unit ball of \mathfrak{B} . Various characterizations of the canonical half-norm associated with \mathfrak{B}_+ are also given.

0. Introduction. Let \mathfrak{B} be a Banach space with a positive cone \mathfrak{B}_+ i.e., a norm-closed proper convex cone, and introduce the dual cone \mathfrak{B}_+^* , in the dual \mathfrak{B}^* of \mathfrak{B} , by

$$\mathfrak{B}_+^* = \{f \in \mathfrak{B}^*; f(a) \geq 0, a \in \mathfrak{B}_+\}.$$

It follows that \mathfrak{B}_+^* is a norm-closed convex cone and if \mathfrak{B}_+ is weakly generating in the sense that $\mathfrak{B} = \overline{\mathfrak{B}_+ - \mathfrak{B}_+}$, where the bar denotes the closure, then \mathfrak{B}_+^* is proper. We shall call \mathfrak{B}_+ *orthogonally generating* if every $a \in \mathfrak{B}$ admits a decomposition $a = a_1 - a_2$ with $a_i \in \mathfrak{B}_+$ ($i = 1, 2$) and

$$\|a_1 + a_2\| = \|a_1 - a_2\|.$$

Clearly, every Banach lattice and the hermitian part of a C^* -algebra have orthogonally generating positive cones with $a_1 = a_+$ and $a_2 = a_-$ where a_{\pm} denote the usual positive and negative components of a .

In general, the cones \mathfrak{B}_+ and \mathfrak{B}_+^* define order relations on \mathfrak{B} and \mathfrak{B}^* respectively. If $a, b \in \mathfrak{B}$, one sets $a \geq b$ whenever $a - b \in \mathfrak{B}_+$. Similarly, if $f, g \in \mathfrak{B}^*$, one sets $f \geq g$ whenever $f - g \in \mathfrak{B}_+^*$.

The main purpose of this note is to determine conditions under which a general $f \in \mathfrak{B}^*$ has an orthogonal norm-unique Jordan decomposition, i.e., a decomposition of the form $f = f_+ - f_-$ with $f_{\pm} \in \mathfrak{B}_+^*$ such that

- (1) (Jordan decomposition) $\|f\| = \|f_+\| + \|f_-\|$;
- (2) (Orthogonality) $\|f_+ + f_-\| = \|f_+ - f_-\|$;

(3) (Norm-uniqueness) If $f = g_1 - g_2$ is another decomposition with the property (1), then

$$\|f_+\| = \|g_1\| \quad \text{and} \quad \|f_-\| = \|g_2\|.$$

Our principal result is the following:

THEOREM 1. *If \mathfrak{B}_+ is orthogonally generating, the following conditions are equivalent:*

1. *For every $a \in \mathfrak{B}$*

$$\|a\| = \inf\{\lambda \geq 0; -\lambda u \leq a \leq \lambda v, u, v \in \mathfrak{B}_1\},$$

where \mathfrak{B}_1 denotes the unit ball of \mathfrak{B} .

2. *Every $f \in \mathfrak{B}^*$ has an orthogonal norm-unique Jordan decomposition.*

3. *If $a = a_1 - a_2$ is an orthogonal decomposition of $a \in \mathfrak{B}$, then*

$$\|a\| = \|a_1\| \vee \|a_2\| = N(a) \vee N(-a)$$

where N is the canonical half-norm associated with \mathfrak{B}_+ .

Before giving the definition of half-norms, we note that condition 1 is easily verified if \mathfrak{B} is the hermitian part of a C^* -algebra. First set

$$\|a\|_1 = \inf\{\lambda \geq 0; -\lambda u \leq a \leq \lambda v, u, v \in \mathfrak{B}_1\}$$

and note that $\|a\|_1 \leq \|a\|$. Next adjoin an identity element $\mathbf{1}$ if necessary, and remark that in principle this reduces $\|\cdot\|_1$. But, if $-\lambda u \leq a \leq \lambda v$ with $u, v \in \mathfrak{B}_1$, then $(\mathbf{1} - u) \leq (\mathbf{1} + a/\lambda) \leq \mathbf{1} + v$ and $0 \leq \mathbf{1} + a/\lambda \leq 2\mathbf{1}$. Therefore, $\|a\| \leq \lambda$ and $\|a\| = \|a\|_1$.

The situation is quite different for order complete Banach lattices. Theorem 1 then implies (see [7], Example 1.5) that \mathfrak{B}^* has such a Jordan decomposition if, and only if, \mathfrak{B} is an AM-space.

The proof of Theorem 1 is based upon the notion of a *half-norm*, i.e., a function N over \mathfrak{B} with the properties:

(N1) $0 \leq N(a) \leq k\|a\|$ for some $k > 0$,

(N2) $N(a_1 + a_2) \leq N(a_1) + N(a_2)$,

(N3) $N(\lambda a) = \lambda N(a)$ for all $\lambda > 0$,

(N4) $N(a) \vee N(-a) = 0$ if, and only if, $a = 0$.

The existence of a half-norm over \mathfrak{B} is equivalent to the existence of a positive cone \mathfrak{B}_+ in \mathfrak{B} . In fact, if N is a half-norm on \mathfrak{B} , then

$$\mathfrak{B}_+ = \{a \in \mathfrak{B}; N(-a) = 0\}$$

is a positive cone. Conversely, if \mathfrak{B}_+ is a positive cone in \mathfrak{B} , then

$$N(a) = \inf\{\|a + b\|; b \in \mathfrak{B}_+\}$$

defines a half-norm over \mathfrak{B} . Following Arendt, Chernoff and Kato [2] we call this latter half-norm the *canonical half-norm* associated with \mathfrak{B}_+ . Note that it automatically satisfies

$$0 \leq N(a) \leq \|a\|.$$

Half-norms are particularly useful for studying positive semigroups [2], [3], [9]. We derive various properties of half-norms in §2, after discussing the Jordan decomposition property in §1.

1. The Jordan decomposition. Throughout this section, let \mathfrak{B} be a Banach space ordered by a positive cone \mathfrak{B}_+ and let N be a half-norm associated with \mathfrak{B}_+ , i.e., N is such that

$$\mathfrak{B}_+ = \{a; N(-a) = 0\}.$$

LEMMA 2. *Let f be a linear functional on \mathfrak{B} . If there exists a constant $\alpha > 0$ such that*

$$f(a) \leq \alpha N(a) \quad \text{for all } a \in \mathfrak{B}$$

then f is positive and continuous. Conversely, if N is the canonical half-norm associated with \mathfrak{B}_+ and $f \in \mathfrak{B}^$ is positive, then*

$$f(a) \leq \|f\| N(a) \quad \text{for all } a \in \mathfrak{B}.$$

Proof. If $f(a) \leq \alpha N(a)$ for all $a \in \mathfrak{B}$, then $-f(a) \leq \alpha N(-a)$, and f is obviously positive. But by condition (N1),

$$|f(a)| \leq \alpha N(a) \vee N(-a) \leq \alpha k \|a\|$$

i.e., f is continuous. Conversely, if $f \in \mathfrak{B}^*$ is positive and N is canonical, we choose $b_n \in \mathfrak{B}_+$ such that $\|a + b_n\| < N(a) + 1/n$. Then,

$$f(a) \leq f(a + b_n) \leq \|f\| \|a + b_n\| \leq \|f\| \left(N(a) + \frac{1}{n} \right).$$

Hence, $f(a) \leq \|f\| N(a)$.

We denote by \mathfrak{B}_N^* the set of all $f \in \mathfrak{B}^*$ such that $f(a) \leq N(a)$ for all $a \in \mathfrak{B}$. The importance of this set is due to the following N -extension theorem

LEMMA 3. *For every $a \in \mathfrak{B}$, there exists $f \in \mathfrak{B}_N^*$ such that $f(a) = N(a)$.*

Proof. We may assume that $N(a) \neq 0$. Let \mathfrak{N} be the linear space spanned by a and define a linear functional g on \mathfrak{N} by

$$g(\xi a) = \xi N(a) \quad \text{for all } \xi \in \mathbf{R}.$$

Then, it is easy to see that

$$g(b) \leq N(b) \quad \text{for all } b \in \mathfrak{M}.$$

It now follows from the subadditivity and homogeneity of N that g has a linear extension f to \mathfrak{B} satisfying the properties of the lemma (see, for example, [4], pages 65–66).

We remark that this lemma implies

$$N(a) = \sup\{f(a); f \in \mathfrak{B}_N^*\}.$$

Next, we define the conjugate N^* of N by

$$N^*(f) = \sup\{f(a); a \in \mathfrak{B}_+, \|a\| \leq 1\}$$

for every $f \in \mathfrak{B}^*$. Then, N^* has the following properties;

- (N1)* $0 \leq N^*(f) \leq \|f\|$
- (N2)* $N^*(f + g) \leq N^*(f) + N^*(g)$,
- (N3)* $N^*(\lambda f) = \lambda N^*(f)$ for $\lambda \geq 0$.

In order that N^* is a half-norm on \mathfrak{B}^* it must also satisfy the condition

$$(N4)* \quad N^*(f) \vee N^*(-f) = 0 \text{ if, and only if, } f = 0.$$

For this we need an assumption on the positive cone \mathfrak{B}_+ . The positive cone \mathfrak{B}_+ is said to be *generating* if every $a \in \mathfrak{B}$ has a decomposition $a = a_1 - a_2$ with $a_i \in \mathfrak{B}_+$ ($i = 1, 2$). Ando [1], has proved that when \mathfrak{B}_+ is generating there exists a constant $\rho > 0$ such that each $a \in \mathfrak{B}$ has a decomposition $a = a_1 - a_2$ with $a_i \in \mathfrak{B}_+$ ($i = 1, 2$), and

$$\|a_1\| \vee \|a_2\| \leq \rho \|a\|.$$

When this is the case, we shall say that \mathfrak{B}_+ is ρ -*generating*.

LEMMA 4. *When \mathfrak{B}_+ is generating, N^* is a half-norm on \mathfrak{B}^* and, for $f \in \mathfrak{B}^*$, f is positive if, and only if, $N^*(-f) = 0$.*

If \mathfrak{B}_+ is ρ -generating then

$$\|f\| \leq \rho(N^*(f) + N^*(-f)).$$

Proof. If $N^*(f) + N^*(-f) = 0$, we have $f(a) = 0$ when $a \geq 0$ or $a \leq 0$. Since \mathfrak{B}_+ is generating, this implies $f = 0$ and, hence, N^* is a half-norm on \mathfrak{B}^* . It is obvious that $N^*(-f) = 0$ if f is positive. The converse follows from

$$-f(a) \leq N^*(-f)\|a\| \quad \text{for } a \geq 0.$$

Now, to prove the last statement, assume that $\alpha > N^*(f) + N^*(-f)$ and choose α_i ($i = 1, 2$) such that $\alpha = \alpha_1 + \alpha_2$, $N^*(f) < \alpha_1$ and $N^*(-f) < \alpha_2$.

Then, for every $a \in \mathfrak{B}$ and its decomposition $a = a_1 - a_2$ with $a_i \in \mathfrak{B}_+$ ($i = 1, 2$),

$$f(a) = f(a_1) - f(a_2) < \alpha_1 \|a_1\| + \alpha_2 \|a_2\|$$

and

$$-f(a) = f(a_2) - f(a_1) < \alpha_1 \|a_2\| + \alpha_2 \|a_1\|.$$

Therefore,

$$|f(a)| < \alpha_1 \rho \|a\| + \alpha_2 \rho \|a\| < \alpha \rho \|a\|,$$

and, hence, $\|f\| \leq \alpha \rho$.

We remark that, if every element of \mathfrak{B} admits a Jordan decomposition one has $\|f\| = N^*(f) + N^*(-f)$.

Now, we start the proof of Theorem 1. We begin with a result of Grosberg and Krein [5].

LEMMA 5. (Grosberg-Krein). *If N is the canonical half-norm associated with \mathfrak{B}_+ the following two conditions are equivalent;*

- (1) $\|a\| = N(a) \vee N(-a)$ for all $a \in \mathfrak{B}$.
- (2) Every element of \mathfrak{B}^* admits a Jordan decomposition.

Proof. Assume that the condition 1 holds and set $P = \{f \in \mathfrak{B}^*; \|f\| \leq 1, f \geq 0\}$. Then, since $\mathfrak{B}_N^* \subset P$, we can conclude from Lemma 3 that the polar P^0 of P coincides with the closed unit ball \mathfrak{B}_1 of \mathfrak{B} . Hence, the closed unit ball \mathfrak{B}_1^* of \mathfrak{B}^* coincides with the bipolar P^{00} . Therefore Grothendieck's argument [6] leads us to condition 2. Conversely, if condition 2 holds and $N(a) \vee N(-a) < 1$, we choose an arbitrary $f \in \mathfrak{B}^*$ such that $\|f\| = 1$. Then, for $f_{\pm} \geq 0$ such that $f = f_+ - f_-$ and $\|f\| = \|f_+\| + \|f_-\|$, we have

$$|f_+(a)| \leq \|f_+\| \quad \text{and} \quad |f_-(a)| \leq \|f_-\|$$

In fact, since N is canonical, we can find $b, c \in \mathfrak{B}_+$ such that

$$\|a + b\| < 1 \quad \text{and} \quad \|-a + c\| < 1.$$

Then,

$$f_+(a) \leq f_+(a + b) \leq \|f_+\| \quad \text{and} \quad -f_+(a) \leq f_+(-a + c) \leq \|f_+\|.$$

Therefore, $|f_+(a)| \leq \|f_+\|$. Similarly, $|f_-(a)| \leq \|f_-\|$. Then, $|f(a)| \leq |f_+(a)| + |f_-(a)| \leq \|f\| = 1$ and, hence, $\|a\| \leq 1$.

It was proved in [7], Lemma 1.1, that the canonical half-norm N associated with \mathfrak{B}_+ satisfies

$$N(a) = \inf\{\lambda \geq 0; a \leq \lambda u, u \in \mathfrak{B}_1\}.$$

Therefore, condition 1 in Theorem 1 is another expression of $\|a\| = N(a) \vee N(-a)$. Hence, by the Groshberg-Krein theorem, Lemma 5, every element of \mathfrak{B}^* admits a Jordan decomposition. We are going to show that this decomposition is orthogonal and norm-unique. Note that, if \mathfrak{B}_+ is orthogonally generating, it is 1-generating. In fact, if $a = a_1 - a_2$ is an orthogonal decomposition, then, since $\|a\| = \|a_1 + a_2\|$,

$$\begin{aligned} 2\|a_1\| &= \|(a_1 + a_2) + (a_1 - a_2)\| \\ &\leq \|a_1 + a_2\| + \|a_1 - a_2\| = 2\|a\|, \end{aligned}$$

which implies $\|a_1\| \leq \|a\|$. Similarly, $\|a_2\| \leq \|a\|$. Therefore, the following two lemmas, together with Lemma 5, prove that condition 1 implies condition 2 in Theorem 1.

LEMMA 6. *Assume that \mathfrak{B}_+ is 1-generating and $f = f_1 - f_2$ is a Jordan decomposition of $f \in \mathfrak{B}^*$. Then $\|f_1\| = N^*(f)$ and $\|f_2\| = N^*(-f)$.*

Proof. By Lemma 4, we have

$$\|f\| \leq N^*(f) + N^*(-f).$$

On the other hand, we have $N^*(f) \leq \|f_1\|$, because

$$f(a) = f_1(a) - f_2(a) \leq f_1(a) \leq \|f_1\|$$

if $a \geq 0$ and $\|a\| \leq 1$. Similarly, $N^*(-f) \leq \|f_2\|$. Then, since $\|f\| = \|f_1\| + \|f_2\|$, we must have $N^*(f) = \|f_1\|$ and $N^*(-f) = \|f_2\|$.

LEMMA 7. *Assume that \mathfrak{B}_+ is orthogonally generating and $f = f_1 - f_2$ is a Jordan decomposition of $f \in \mathfrak{B}^*$. Then f_1 and f_2 are orthogonal, i.e.,*

$$\|f_1 + f_2\| = \|f_1 - f_2\|$$

Proof. Let $a = a_1 - a_2$ be an orthogonal decomposition of $a \in \mathfrak{B}$. Then

$$\pm f(a) \leq (f_1 + f_2)(a_1 + a_2).$$

Hence,

$$|f(a)| \leq \|f_1 + f_2\| \|a_1 + a_2\| \leq \|f_1 + f_2\| \|a\|$$

which implies $\|f\| \leq \|f_1 + f_2\|$. On the other hand, it follows from the definition of Jordan decomposition that $\|f_1 + f_2\| \leq \|f_1\| + \|f_2\| = \|f\|$. Therefore f_1 and f_2 are orthogonal.

Next, we prove that condition 2 implies condition 3 in Theorem 1. First, we note that we have

$$\|a\| = N(a) \vee N(-a)$$

by Lemma 5. Now, let $a = a_1 - a_2$ be an orthogonal decomposition. Then, as we have shown above, we have $\|a_i\| \leq \|a\|$ ($i = 1, 2$). On the other hand, since $a \leq a_1$, we have $N(a) \leq N(a_1) \leq \|a_1\|$ and, similarly, $N(-a) \leq \|a_2\|$. Hence,

$$\|a\| \geq \|a_1\| \vee \|a_2\| \geq N(a) \vee N(-a) = \|a\|.$$

That condition 3 implies condition 1 in Theorem 1 is trivial.

2. Half-norms. Let \mathfrak{B} be an ordered Banach space with a positive cone \mathfrak{B}_+ . The equality

$$N(a) = \inf\{\|a + b\|; b \in \mathfrak{B}_+\} = \inf\{\lambda \geq 0; a \leq \lambda u, u \in \mathfrak{B}_1\}$$

referred to in §1, gives an order theoretic characterization of the canonical half-norm N associated with \mathfrak{B}_+ .

The next theorem gives a criterion for another order theoretic characterization of N .

THEOREM 8. *The following conditions are equivalent:*

- (1) $N(a) = \inf\{\lambda \geq 0; a \leq \lambda u, u \in \mathfrak{B}_1 \cap \mathfrak{B}_+\}$,
- (2) *For each $\epsilon > 0$ and $a \in \mathfrak{B}$ there is a decomposition*

$$a = a_+ - a_- \quad \text{with } a_{\pm} \in \mathfrak{B} \text{ and } \|a_+\| \leq (1 + \epsilon)\|a\|.$$

Proof. Assume that condition 1 holds. If $\epsilon > 0$ and $a \in \mathfrak{B}$, there is $u \in \mathfrak{B}_1 \cap \mathfrak{B}_+$ such that $a \leq N(a)(1 + \epsilon)u$. Hence, $a = a_+ - a_-$ with $a_+ = N(a)(1 + \epsilon)u$ and $a_- = a_+ - a$. But, $\|a_+\| \leq N(a)(1 + \epsilon) \leq (1 + \epsilon)\|a\|$. Conversely, assume that condition 2 holds. If $a \leq \lambda u$ with $u \in \mathfrak{B}_+$ and u has a decomposition $u = u_+ - u_-$ with $u_{\pm} \in \mathfrak{B}_+$ and $\|u_+\| \leq 1 + \epsilon$, then $a \leq \lambda u_+$ and,

$$N(a) < \inf\{\lambda \geq 0; a \leq \lambda u, u \in \mathfrak{B}_1 \cap \mathfrak{B}_+\} \leq (1 + \epsilon)N(a).$$

Since this estimate is valid for all $\epsilon > 0$, one has the desired identification.

It was proved in [7], Proposition 1.6, that condition 2 with $\varepsilon = 0$ is implied by the following three equivalent conditions:

(i) There is a $u \in \mathfrak{B}_1$ such that

$$\{a: \|u - a\| < 1\} \subset \mathfrak{B}_+,$$

(ii) \mathfrak{B}_1 has a maximal element u ,

(iii) there is a $u \in \mathfrak{B}_1$ such that $N = N_u$, where

$$N_u(a) = \inf\{\lambda \geq 0; a \leq \lambda u\}.$$

COROLLARY 9. *If $(\mathfrak{B}, \mathfrak{B}_+)$ is the dual of an ordered Banach space \mathfrak{B}_* with positive cone \mathfrak{B}_{*+} and if N is the canonical half-norm associated with \mathfrak{B}_+ , then the following conditions are equivalent:*

(1) $N(a) = \inf\{\lambda \geq 0; a \leq \lambda u, u \in \mathfrak{B}_1 \cap \mathfrak{B}_+\}$,

(2) each $a \in \mathfrak{B}$ has a decomposition $a = a_+ - a_-$ with $a_{\pm} \in \mathfrak{B}_+$ and $\|a_+\| \leq \|a\|$.

Proof. In view of Theorem 8, we only need to show that condition 1 implies condition 2. Now, if condition 1 holds, it follows from Theorem 9 that for $\varepsilon > 0$ and $a \in \mathfrak{B}$ there is a $u_{\varepsilon} \in \mathfrak{B}_1 \cap \mathfrak{B}_+$ such that $a \leq N(a)(1 + \varepsilon)u_{\varepsilon}$. But $\mathfrak{B}_1 \cap \mathfrak{B}_+$ is weak* compact and hence u_{ε} has a weak* limit point u . Therefore

$$N(a)u(\omega) = \lim N(a)(1 + \varepsilon)u_{\varepsilon}(\omega) \geq a(\omega)$$

for all $\omega \in \mathfrak{B}_{*+}$ and $a \leq N(a)u$ by the definition of a dual cone. Now, $a = a_+ - a_-$ with $a_+ = N(a)u \in \mathfrak{B}_+$, $a_- = a_+ - a \in \mathfrak{B}_+$, and $\|a_+\| \leq N(a) \leq \|a\|$.

If \mathfrak{B} is either a Banach lattice or the hermitian part of a C^* -algebra, then each $a \in \mathfrak{B}$ has a canonical decomposition $a = a_+ - a_-$ into positive and negative components $a_{\pm} \in \mathfrak{B}_+$ [8], [4]. In both cases, however $\|a_{\pm}\| \leq \|a\|$ and hence the canonical half-norm has the order theoretic characterization given by condition 1 of Theorem 8.

REFERENCES

- [1] T. Ando, *On fundamental properties of a Banach space with a cone*, Pacific J. Math., **12** (1962), 1163–1169.
- [2] W. Arendt, P. R. Chernoff and T. Kato, *A generalization of dissipativity and positive semigroups*, J. Operator Theory, **8** (1982), 167–180.
- [3] O. Bratteli, T. Digernes and D. W. Robinson, *Positive semigroups on ordered Banach spaces*, J. Operator Theory, (to appear).
- [4] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics*, Vol. I, Springer-Verlag (1979).

- [5] J. Grosberg et M. Krein, *Sur la décomposition des fonctionnelles en composantes positives*, C. R. (Doklady) Acad. Sci., URSS (N.S.) **25** (1939), 723–726.
- [6] A. Grothendieck, *Un resultat sur le dual d'une C^* -algebra*, J. de Math., **36** (1957), 97–108.
- [7] D. W. Robinson and S. Yamamuro, *Addition of an identity to an ordered Banach space*, J. Australian M.S. (to appear).
- [8] H. H. Schaefer, *Banach Lattices and Positive Operators*, Springer-Verlag (1974).
- [9] S. Yamamuro, *Notes on Locally convex spaces with ACK-calibrations*, unpublished manuscript.

Received May 13, 1982.

AUSTRALIAN NATIONAL UNIVERSITY
CANBERRA, A.C.T. 2600, AUSTRALIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)
University of California
Los Angeles, CA 90024

HUGO ROSSI
University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR OGUS
University of California
Berkeley, CA 94720

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, CA 90089-1113

R. FINN and H. SAMELSON
Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906–1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$132.00 a year (6 Vol., 12 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics ISSN 0030-8730 is published monthly by the Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1984 by Pacific Journal of Mathematics

Pacific Journal of Mathematics

Vol. 110, No. 2

October, 1984

Robert A. Bekes , The range of convolution operators	257
Dennis K. Burke and Sheldon Davis , Subsets of ${}^\omega\omega$ and generalized metric spaces	273
Giovanni Emmanuele , A remark on a paper: "Common fixed points of nonexpansive mappings by iteration"	283
I. Erdélyi and Sheng-Wang Wang , On strongly decomposable operators ...	287
Gerhard Gierz , Injective Banach lattices with strong order units	297
Maurizio Letizia , Quotients by complex conjugation of nonsingular quadrics and cubics in \mathbf{P}_C^3 defined over \mathbf{R}	307
P. H. Maserick and Franciszek Hugon Szafraniec , Equivalent definitions of positive definiteness	315
Costel Peligrad and S. Rubinstein , Maximal subalgebras of C^* -crossed products	325
Derek W. Robinson and Sadayuki Yamamuro , Hereditary cones, order ideals and half-norms	335
Derek W. Robinson and Sadayuki Yamamuro , The Jordan decomposition and half-norms	345
Richard Rochberg , Interpolation of Banach spaces and negatively curved vector bundles	355
Dale Rolfsen , Rational surgery calculus: extension of Kirby's theorem	377
Walter Iaan Seaman , Helicoids of constant mean curvature and their Gauss maps	387
Diana Shelstad , Endoscopic groups and base change \mathbf{C}/\mathbf{R}	397
Jerrold Norman Siegel and Frank Williams , Numerical invariants of homotopies into spheres	417
Alladi Sitaram , Some remarks on measures on noncompact semisimple Lie groups	429
Teruhiko Soma , Atoroidal, irreducible 3-manifolds and 3-fold branched coverings of S^3	435
Jan de Vries , On the G -compactification of products	447
Hans Weber , Topological Boolean rings. Decomposition of finitely additive set functions	471